

Health-Promotion Interventions Targeting Multiple Behaviors: A Meta-Analytic Review of General and Behavior-Specific Processes of Change

Angela L. Zhang¹, Sicong Liu², Benjamin X. White³, Xi C. Liu³, Marta Durantini², Man-pui Sally Chan²,
Wenhao Dai², Yubo Zhou¹, Melody Leung², Qijia Ye², Devlin O’Keefe², Lidia Palmese², and
Dolores Albarracín^{1, 2}

¹ Department of Psychology, University of Pennsylvania

² Annenberg School for Communication, University of Pennsylvania

³ Department of Psychology, University of Illinois Urbana-Champaign

Although health-promotion interventions that recommend changes across multiple behavioral domains are a newer alternative to single-behavior interventions, their general efficacy and their mechanisms of change have not been fully ascertained. This comprehensive meta-analysis (6,878 effect sizes from 803 independent samples from 364 research reports, $N = 186,729$ participants) examined the association between the number of behavioral recommendations in multiple-behavior interventions and behavioral and clinical change across eight domains (i.e., diet, smoking, exercise, HIV [Human Immunodeficiency Virus] prevention, HIV testing, HIV treatment, alcohol use, and substance use). Results showed a positive, linear effect of the number of behavioral recommendations associated with behavioral and clinical change across all domains, although approximately 87% of the samples included between 0 and 4 behavioral recommendations. This linear relation was mediated by improvements in the psychological well-being of intervention recipients and, in several domains (i.e., HIV, alcohol use, and drug use), suggested behavioral cuing. However, changes in information, motivation, and behavioral skills did not mediate the impact of the number of recommendations on behavioral and clinical change. The implications of these findings for theory and future intervention design are discussed.

Public Significance Statement

The present meta-analysis examined the effects of multiple-behavior interventions and processes of change. A higher number of recommendations is associated with more positive behavioral and clinical outcomes. Analyses of potential mediators suggest two key mechanisms of change underlying this relation: (a) cuing of one behavior by another and (b) improvements in psychological well-being. This work advances our knowledge of the optimal number of recommendations to include in health-promotion interventions as well as the mechanisms through which multiple recommendations operate. Thus, it can inform the design of future interventions to increase their efficacy in facilitating positive health outcomes.

Keywords: multiple-behavior intervention, efficacy, meta-analysis, mechanism of change, public health

Supplemental materials: <https://doi.org/10.1037/bul0000427.supp>

This article was published Online First June 24, 2024.

Angela L. Zhang  <https://orcid.org/0000-0001-8553-5941>

Dolores Albarracín  <https://orcid.org/0000-0002-9878-942X>

Angela L. Zhang and Sicong Liu contributed equally to this article and share first authorship. The data supporting the findings reported in this study are openly available in the Open Science Framework at https://osf.io/qsx57/?view_only=6fffd7080d844e98cbd2d0834ed61e9, and the study was pre-registered at https://osf.io/mq5kr/?view_only=54d04d57b5ef45aaa480a47259cb91cd. The authors declare no competing interests. An original version of this work was part of Benjamin X. White’s dissertation. Dolores Albarracín received funding from Grants R01 AI147487, DP1 DA048570, and R01 MH094241 from the National Institutes of Health and from Grant NSF 2031972 from the National Science Foundation.

This work is licensed under a Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License (CC BY-NC-ND 4.0; <https://creativecommons.org/licenses/by-nc-nd/4.0>). This license permits

copying and redistributing the work in any medium or format for noncommercial use provided the original authors and source are credited and a link to the license is included in attribution. No derivative works are permitted under this license.

Angela L. Zhang played a supporting role in data curation and validation and an equal role in writing—original draft and writing—review and editing. Sicong Liu played a lead role in formal analysis, a supporting role in data curation and validation, and an equal role in writing—original draft and writing—review and editing. Benjamin X. White played a supporting role in data curation and writing—original draft. Xi C. Liu played a supporting role in data curation and validation. Marta Durantini played a supporting role in data curation and validation. Man-pui Sally Chan played a lead role in data curation and a supporting role in validation. Wenhao Dai played a supporting role in data curation and validation. Yubo Zhou played a supporting role in data curation and validation. Melody Leung played a supporting role in data curation and validation. Qijia Ye played a supporting role in data curation and validation. Devlin O’Keefe played a supporting role in

continued

Clusters of unhealthy behaviors contribute to the most prevalent diseases in the United States and most areas of the world (Farnham et al., 2010; Heron, 2019; Office of Disease Prevention and Health Promotion, 2018). As a result, the most promising behavioral-health programs have moved from attempting to change a single behavior to attempting to change behavioral clusters such as improving lifestyle (King et al., 2015). For example, interventions targeting both diet and exercise have decreased cardiovascular risk (Salas-Salvadó et al., 2019), delayed onset of type 2 diabetes (Gong et al., 2019), and improved cognitive functioning (Ngandu et al., 2015). Other interventions have been successful in simultaneously decreasing both substance use and risky sexual behaviors (Mouttapa et al., 2009; Tucker et al., 2017). Prior syntheses of the efficacy of these programs (Dai et al., 2020; Sunderrajan et al., 2021; Wilson et al., 2015), however, have looked at a small number of domains (i.e., one to three) and left critical questions unanswered. One of these questions is whether higher numbers of recommendations (i.e., exhortations to perform specific health behaviors) increase behavioral and clinical change across diverse behavioral domains. Whereas the number of recommendations in lifestyle programs has shown to facilitate change only up to a point, the number of recommendations in drug and alcohol programs as well as HIV (human immunodeficiency virus) programs appears to increase clinical and behavioral change in a linear fashion. Therefore, the optimal number of behavioral recommendations across the board and the processes underlying the efficacy of multiple-behavior interventions remain unclear.

Another limitation of the literature is the absence of a process framework about multiple-behavior change mechanisms and evidence testing those mechanisms. In this article, we introduce a conceptualization in which multiple behavioral recommendations can trigger either general or behavior-specific processes of behavioral change. General processes of behavioral change tend to occur spontaneously and apply to different behavioral domains, thus being relatively efficient. These processes include *cuing*, a mechanism in which changes in one behavior cue changes in other behaviors (for related processes, see Bargh et al., 1992; Weingarten et al., 2016; Wood, 2017) such as alcohol drinking cuing consumption of calorie-dense snacks (Carels et al., 2014). General processes of change also comprise changes in psychological well-being, including reductions in depression and anxiety symptoms (Worley et al., 2012) and increases in self-control (Hofmann et al., 2014), all of which cut across behaviors. These general processes are likely to occur by simply facilitating multiple-behavior change efficiently via the introduction of new cues to positive behaviors (Albarracín, 2021; Bargh, 1994) or via a reduction in stress and consequent engagement in unhealthy behaviors to reduce stress. Consequently, these processes may lead to interventions with higher numbers of behavioral recommendations exerting a linear impact on change in behavioral and clinical outcomes. Prior work examining the benefits of one behavior for other behaviors has referred to these changes as “spill-over” (Mata et al., 2009) or “transfer” (Lippke et al., 2012) effects.

In contrast, behavior-specific processes of change require thinking about the specific behaviors recommended in an intervention. For

example, interventions that make more recommendations may increase information and behavioral skills acquisition, such as knowledge about cardiovascular health and grocery shopping skills, which can in turn facilitate regular exercise and a healthy diet (Kiernan et al., 2013). Moreover, multiple-behavior interventions may improve motivation by strengthening attitudes, norms, and intentions to perform each recommended behavior (Albarracín et al., 2001). Nevertheless, processing information about each behavior may require cognitive capacity and motivation to think about the behaviors (Albarracín, 2002), thus limiting how many recommendations can be introduced (Wilson et al., 2015). Consequently, if these processes dominate, they may result in a curvilinear effect of the number of behavioral recommendations on behavioral and clinical change due to each recommendation incurring costs that diminish benefits as the number of recommendations increases. Compensatory effects (e.g., indulging in unhealthy snacks now with the intention of exercising more later; Knäuper et al., 2004) have after all been identified in several domains. For example, physically active people consume more alcohol, people who frequently consume alcohol engage in more physical activity (C. R. Nigg et al., 2009), and participating in an exercise intervention can lead to increasing energy intake (Martin et al., 2019). Similarly, workers who switch from a sedentary occupation to a more physically active one compensate by exercising less in their leisure time, whereas the opposite pattern is found for workers who switch from a physically active occupation to a sedentary one (Nooijen et al., 2018).

In this study, we conducted a meta-analysis of multiple-behavior interventions to assess the extent to which cuing and increases in psychological well-being, information, motivation, and behavioral skills help to explain the effects of multiple behavioral recommendations on behavioral and clinical change. We first evaluated the shape of the relation between the number of recommendations and intervention efficacy, considering both behavioral outcomes (self-report measures of behavior; e.g., self-report scales) and clinical outcomes (e.g., blood pressure, cholesterol, sexually transmitted infection [STI] test results). This synthesis included a comprehensive data set spanning eight prominent behavioral intervention domains: (a) exercise, (b) diet, (c) smoking, (d) HIV prevention, (e) HIV testing, (f) HIV treatment, (g) alcohol use, and (h) substance use. These domains are of particular interest to behavioral change researchers because they represent some of the leading modifiable risk factors associated with the highest burdens of disease, death, and economic costs both in the United States and globally (Bolnick et al., 2020; Mokdad et al., 2018; Murray et al., 2020; Vos et al., 2020), and the literatures that cover them have been used to test prior models of behavioral change (e.g., see Carpenter, 2010; McEachan et al., 2011). We thus examined behavioral cuing and changes in psychological well-being, information, motivation, and behavioral skills in these diverse and important areas.

Single-Behavior Interventions and Their Shortcomings

The first generation of interventions developed to change behavior focused on changing one behavior deemed important for

data curation and validation. Lidia Palmese played a supporting role in data curation and validation. Dolores Albarracín played a lead role in conceptualization, funding acquisition, and supervision and a supporting role in data curation, validation, writing—original draft, and writing—review and editing.

Correspondence concerning this article should be addressed to Dolores Albarracín, Department of Psychology, University of Pennsylvania, 202 South 36th Street, Philadelphia, PA 19104, United States. Email: dalba@upenn.edu

a population. Early HIV prevention interventions in the 1980s, when the disease was first identified, focused on increasing condom use (Wolitski et al., 2005), and many alcohol- and substance-use treatment programs have focused on only one problem behavior (Murthy, 2017). Weight loss interventions have generally been more mixed in that some recommend both exercise and diet (e.g., Parsons et al., 2005), but many still recommend solely changes in either diet or physical activity, with early meta-analyses noting that about 70% of studies recommended a single behavior as the target for change (Conn et al., 2002).

One limitation of single-behavior interventions has been that they routinely produce only small-to-medium effect sizes in behavioral or clinical change ($d = 0.3$, interpreted as a change of 0.3 standard deviations). For example, a meta-analysis of interventions to increase physical activity found that the average intervention changed physical activity by about a quarter of a standard deviation ($d = 0.27$; Rhodes, Janssen, et al., 2017). Interventions to curb alcohol ($d = 0.28$; Dotson et al., 2015); quit smoking ($d = 0.34$; Wu et al., 2006); and prevent HIV ($d = 0.40$; Albarracín et al., 2005) have all shown medium effect sizes as well.

The limitations of single-behavior interventions are perhaps not surprising given the reality of the challenges people experience with respect to changing their health behaviors. Several observational studies noted that, rather than people performing a single unhealthy behavior, they often have clusters of unhealthy practices (Keller et al., 2011; Lippke et al., 2012; Ritchwood et al., 2015). For lifestyle behaviors, there are positive correlations between exercise and diet: $r = .16$ to $.26$; between exercise and smoking: $r = .21$; and between diet and smoking: $r = .11$ (Keller et al., 2011; Lippke et al., 2012). A meta-analysis of observational studies of adolescent sexual health behaviors also found moderate associations between use of alcohol and/or substances and risky sexual behaviors: $r = .22$; between alcohol use and unprotected sex: $r = .11$; between substance use and unprotected sex: $r = .18$; between alcohol use and multiple partners: $r = .25$; and between substance use and multiple partners: $r = .25$ (Ritchwood et al., 2015). What these observational studies suggest is that single-behavior interventions may be ill equipped to address these patterns of correlated behaviors.

Multiple-Behavior Interventions

Noting the limitations of single-behavior interventions, behavioral change researchers began developing and testing multiple-behavior interventions (Noar et al., 2008; Prochaska et al., 2008; Smedley & Syme, 2001). A multiple-behavior intervention is characterized by recommending change in at least two health-behavior domains either simultaneously or sequentially. Where a single-behavior intervention may recommend that clients increase their physical activity (one behavioral domain), a multiple-behavior intervention may recommend changes in lifestyle, such as increasing physical activity, reducing fat intake, and quitting smoking (three behavioral domains). Initially, interventions targeted multiple behaviors because populations had multiple risk factors, expecting a similar amount of change in each behavior and expecting additive effects on clinical outcomes such as improvements in cardiovascular health (Hjermann et al., 1981; Kjelsberg & Multiple Risk Factor Intervention Trial Research Group, 1982; Wilhelmsen et al., 1986). Yet, other studies, such as Spring et al. (2009), have shown that combining smoking and weight control advice not only improves cardiovascular health but also leads to

greater smoking abstinence and less weight gain compared to recommending smoking cessation alone. Likewise, interventions that combine drug and alcohol risk reduction with sexual risk reduction lead to larger decreases in sexually transmitted infections than do interventions targeting only sexual risk reduction (Bryan et al., 2018).

Initial tests of whether recommending change in multiple behaviors leads to greater behavioral change for each recommended behavior have been promising. Across three meta-analyses, multiple-behavior interventions outperformed single-behavior ones in how much behavioral change they produced on average (Dai et al., 2020; Sunderrajan et al., 2021; Wilson et al., 2015). Still, the shape of the impact of the number of behavioral recommendations on average clinical and behavioral change remains a question. Both Dai et al. (2020) and Sunderrajan et al. (2021) noted a positive, linear effect of the number of recommendations, such that each recommendation added about $d = 0.07$ of improvement in behavioral and clinical outcomes on average. In contrast, Wilson et al. (2015) noted that studies with two to three recommendations outperformed those with only one ($d = 0.33$ vs. $d = 0.17$), but that there was no difference between single- and multiple-behavior programs once there were four or more recommendations ($d = 0.33$ vs. $d = 0.19$). Yet, these prior syntheses have concerned relatively circumscribed domains, whereas our meta-analysis tried to clarify whether the number of recommendations exerts linear or curvilinear effects on behavioral and clinical change across eight domains of public health significance.

Theoretical Framework: General and Behavior-Specific Processes of Behavioral Change

Despite this important body of past literature, prior meta-analytic work has predominantly focused on demonstrating the efficacy of multiple-behavior interventions in specific health areas rather than examining the likely mechanisms of behavioral change across health behaviors. To fill this gap, we propose that multiple recommendations can operate through either general or behavior-specific processes of behavioral change by which changes in one behavior potentiate changes in other behaviors.

General Processes

General processes of behavioral change are those that generalize across behavioral domains because they do not involve thinking about the specific content of a behavioral recommendation. As a result, these processes require minimal cognitive capacity and motivation to implement the recommended changes and can thus be executed efficiently. One example of a general process is behavioral cuing. Prior research on classical conditioning (Pavlov, 1927) and priming (Bargh et al., 1992; Dijksterhuis et al., 2000) has demonstrated that certain stimuli can provide automatic cues for behavior. More recently, research has shown that habitual behaviors (i.e., habits), defined as behaviors that occur with a high level of automaticity (i.e., low or no cognitive effort), are tied to consistent environmental cues (Verplanken & Orbell, 2003; Wood & Neal, 2007). For example, people reflecting on their behavior report going to the bathroom to brush their teeth with no thought once they start their bedtime routine (McCloskey & Johnson, 2019). Behavioral cuing may occur as a result of habit strength, as habituation of one behavior (e.g., exercise) may free up cognitive resources needed to self-regulate another behavior (e.g., eating;

Fleig et al., 2011). In line with this hypothesis, exercise has been found to be associated with healthy eating habits via increases in exercise habit strength (Fleig et al., 2011, 2014). All of these studies thus hint at the possibility that changes in one behavior, such as exercise, may seamlessly produce changes in other behaviors, such as alcohol use.

Changes in psychological well-being (e.g., decreases in stress, anxiety, and depression) represent another general process of behavioral change. Prior research (e.g., Appelhans et al., 2012; Thoits, 2010) has shown that experiencing low psychological well-being (e.g., anxiety, stress, or depression) increases risky behaviors such as binge eating, alcohol use, substance use, and risky sex. Conversely, engagement in unhealthy behaviors, such as smoking, has been implicated as a risk factor for psychological disorders (Firth et al., 2020). Accordingly, a higher number of behavioral recommendations may produce behavioral change through increases in psychological well-being. For example, increases in physical activity have been found to decrease anxiety and depression (Ashdown-Franks et al., 2020; Marcolina, 2007), and antidepressant treatment has been shown to improve exercise performance (Özerbil et al., 2006). More generally, behavioral activation has been shown to lift depression, possibly through dopaminergic mechanisms (Salamone et al., 2015) that may come into play when multiple behaviors are recommended.

In the case of multiple-behavior interventions, general processes would suggest a linear relation between the number of recommendations in an intervention and resulting behavioral or clinical change. For the proposed mechanism of behavioral cuing, this premise is supported by multitasking research findings that people swapping from one task to another (a scenario similar to doing one thing and being cued to do another) perform at the same level when the task swaps automatically and when they are told to prepare to swap tasks (Rogers & Monsell, 1995). Furthermore, the cost of swapping remains stable regardless of how many different behaviors are being swapped (Cho et al., 2015; Kiesel et al., 2010), again implying automaticity in performing multiple tasks. As a result, executing multiple behaviors may occur without posing significant attentional demands and lead to linear effects of the number of recommendations on behavioral change. The same might be the case for changes in emotional well-being, as promoting well-being typically frees cognitive resources and should influence other behaviors in an effortless fashion. Nevertheless, changes in well-being should be accompanied by changes in stress and/or depression, whereas cuing implies that changes in one behavior mediate changes in another behavior.

Behavior-Specific Processes

Behavior-specific processes are likely to involve changes in information, motivation, and behavioral skills pertinent to each behavior (Fisher et al., 2006; Fisher & Fisher, 1992). Thus, these specific processes are likely to require cognitive capacity, motivation, and time because they necessitate thinking about the specific content of a behavioral recommendation. Conscious and effortful processes have been a consistent predictor of behavioral change in multiple areas. Studies of student success, for example, have shown that students who are more engaged with lecture material perform better in both remembering material and applying it properly (Bakker et al., 2015; Robinson, 2013). Furthermore, models of goal pursuit

have suggested that tasks that require effort may enhance pursuit and resulting change, showing benefits to behavior-specific processes of change (Bandura, 1977; Lee et al., 2015).

One limitation of behavior-specific processes of behavioral change is that effortful processes can lead to underperformance when resources are overtaxed. For example, when learning new information, confusion and conflict stemming from receiving too much information can reduce responsiveness (Harkins & Petty, 1981) and, of course, recall (Hall et al., 2015; McDonald et al., 2017). People may also devalue hard-to-implement behaviors (Lee et al., 2015) or burn out when their resources are challenged (Angrave & Charlwood, 2015; Irie et al., 2001). The primary implication of these findings is that recommending a higher number of behaviors may decrease change for each behavior, resulting in a curvilinear relation in which change flattens after a certain number of recommendations is made.

The impact of behavior-specific processes of change is likely to be apparent in changes in information (Fisher et al., 2006; Fisher & Fisher, 1992). Models of health behavior have consistently argued that behavior is influenced by people's beliefs, many of which are based on knowledge about the behavior and its outcomes (Ajzen, 1991; Glanz & Bishop, 2010). For example, people's understanding of skin cancer risk and facts regarding the benefits of using sunscreen correlates with more sunscreen use (Grubbs & Tabano, 2000), and parental knowledge about vaccines correlates with vaccine uptake in young children (Smith et al., 2017). Interventions with more behavioral recommendations may thus provide unique information that is applicable to the target behaviors and exert an influence through changes in information.

Multiple-behavior interventions may also affect the motivation to enact specific behaviors, another key determinant and mediator of behavioral change that comprises attitudes, social norms, and intentions (Ajzen, 1991; Rich et al., 2015; Topa & Moriano, 2010). In a meta-analysis of condom use interventions, attitudes and beliefs predicted intentions to use condoms in future encounters (Albarracín et al., 2005). Furthermore, prior work has noted the potential for motivation to spillover between health behaviors. For example, engagement in physical activity predicts better regulation of eating behaviors, and this relation is mediated, in part, by increased intrinsic motivation to exercise (Mata et al., 2009). Similarly, exercise is associated with greater motivation to quit smoking, which, in turn, is associated with higher odds of staying abstinent (LaRowe et al., 2023).

The number of behavioral recommendations may also promote behavioral change by improving behavioral skills. Behavioral skills, people's ability to implement routines that promote successful performance of specific behaviors, are key to ensuring behavioral change. For example, interventions targeting HIV prevention, treatment adherence, and weight loss have found that skills in monitoring one's own behavior and making corrections if goals are not being met are both important to changing behavior (Albarracín et al., 2005; Burke et al., 2011; Venditti et al., 2014). Furthermore, a recent metareview of the health-behavior-change literature identified skills training as one of the intervention features most often associated with reduced engagement in risk-taking behaviors, such as smoking, unhealthy eating, and alcohol use (Protogerou et al., 2020). Recommendations to change multiple behaviors may thus facilitate the acquisition and development of behavioral skills.

All in all, we expected changes in information, motivation, and behavioral skills to reflect behavior-specific processes that require attention to each behavior and may thus pose limits in terms of the number of behaviors that are of benefit. We recognize that knowledge about one behavior may be relevant to other behaviors, changing attitudes toward one behavior can produce unintended changes in other attitudes (Fishbach et al., 2004; Glanz & Bishop, 2010; McDonald et al., 2017; Pintrich, 2000; Stedry & Kay, 1966), and some skills cut across domains. Nevertheless, reviewing information, behavioral outcomes, and skills necessary to quit smoking is sufficiently time consuming to prevent also focusing on information, behavioral outcomes, and skills about using condoms. Therefore, even if information, motivation, and behavioral skills could generalize over time, the process is likely to be more demanding than changes in one behavior cuing changes in other behaviors or improvements in psychological well-being reducing the need to manage stress by smoking or having risky sex.

The Present Meta-Analysis

We conducted a comprehensive meta-analysis of multiple-behavior interventions to assess the effects of the number of recommendations included in an intervention on behavioral and clinical improvement. Interventions spanned the domains of physical exercise, diet, smoking, HIV prevention, HIV testing, HIV treatment, alcohol use, and substance use. In total, we synthesized changes in 58 different behavioral and clinical outcomes into effect sizes that represent behavioral improvements in exercise; diet; smoking; HIV prevention, testing, and/or treatment; alcohol use; or substance use. In addition, we synthesized changes in measures that represent changes in psychological well-being (e.g., depression and anxiety), health information (e.g., exercise and diet), motivation (e.g., perceived importance and attitude), and behavioral skills (e.g., stress management and decision-making skills), all of which were considered possible mechanisms of change. Although the effect sizes could also be calculated by comparing treatment groups to control groups, a problem associated with this between-group comparison approach is that different trials may employ different types of control groups (i.e., passive vs. active) and systematically affect the computed effect sizes (Karlsson & Bergmark, 2015). Specifically, a given trial with a passive control is likely to result in larger between-group comparison effect size computations than another trial with an active control, which can be regarded as receiving a minimum amount of treatment (see, Brookmeyer et al., 2016).

The main analyses took advantage of recently developed metaregression models of correlated and hierarchical effects with three levels to account for nonindependent observation from both data hierarchy (i.e., measures nested under trial conditions/samples that were nested under trials) and correlated sampling errors (e.g., multiple measures from a given trial condition). One focus of the analyses was to examine the strength and linearity of the effect of the number of behavioral recommendations on behavioral and clinical change. As the number of recommendations has generally accrued naturalistically in the literature (Cochran & Chambers, 1965), we used propensity-score modeling as an optimal way of controlling for differences among conditions with different numbers of recommendations and improving confidence in our inferences (Austin, 2011).

To explicate the effect associated with the number of behavioral recommendations, we tested hypotheses involving both general and

behavior-specific processes, including (a) changes in some outcome behaviors cuing changes in another outcome behavior and (b) changes in the mediators of psychological well-being, information, motivation, and behavioral skills. The behavioral cuing effect and changes in psychological well-being may coexist with linear changes in the number of recommendations, as these processes do not require substantial cognitive resources to function. In contrast, mediation through information, motivation, and/or behavioral skills may be associated with a curvilinear relation between the number of recommendations and behavioral and clinical change, as these processes require greater cognitive resources to take effect. The meta-analysis was preregistered at https://osf.io/mq5kr/?view_only=54d04d57b5ef45aaa480a47259cb91cd. For a list of deviations from the preregistration and coverage of preregistered analyses not included in this article, see Supplemental Material A.

Method

Review and Inclusion Criteria

Eight different behavioral research domains were preidentified to be included in the search: (a) exercise, (b) diet, (c) smoking, (d) HIV prevention, (e) HIV testing, (f) HIV treatment, (g) alcohol-use, and (h) other substance-use interventions. We conducted searches using MEDLINE, APA PsycInfo, EBSCO, Scopus, the Web of Science Core Collection, Journal Storage, and Crossref. In addition to these database searches, we conducted searches of conference abstracts and ProQuest for unpublished literature. When reports were in languages other than English, they were translated and screened.

Behavioral Domain Searches

The following keywords were used to find intervention reports: “intervention,” “health education,” “persuasion,” “recommendation,” “treatment,” “educational program,” “rehabilitation,” “counseling outcomes,” “treatment outcomes,” “treatment effectiveness evaluation,” “treatment compliance,” “health promotion,” “behavioral change,” and “randomized trial.” Within each of the eight domains (exercise, diet, smoking, HIV prevention, HIV testing, HIV treatment, alcohol use, and substance use), these general keywords were combined with unique keywords to identify domain-specific intervention reports, as explained below.

1. To identify *diet interventions*, we used the following keywords: “binge eating,” “body image,” “bulimia,” “caloric intake,” “craving,” “diet,” “dietary restraint,” “eating behavior,” “eating disorders,” “fat intake,” “food intake,” “fruit intake,” “metabolism disorders,” “healthy nutrition,” “obesity,” “sugar intake,” “vegetable intake,” “weight control,” “weight loss,” and “healthy eating.”
2. To identify *exercise interventions*, we used the following keywords: “aerobic exercise,” “body image,” “physical activity,” “sport training,” “strength training,” “weight control,” “weight loss,” “lack of exercise,” “walking,” “gymnastics,” “going to gym,” “running,” “biking,” “work out,” and “physical inactivity.”
3. To identify *smoking interventions*, we used the keywords (a) “tobacco” and (b) “smoking.”

4. To identify *HIV prevention, HIV testing, and HIV treatment interventions*, we used the keywords “HIV,” “AIDS,” “STI/sexually transmitted disease,” “condom use,” “circumcision,” “alcohol use,” “drug use,” and “adherence.”
5. Finally, to identify *alcohol-use and substance-use interventions*, we used the keywords (a) “alcohol,” (b) “drug,” and (c) “substance.”

The specific database search queries are shown in [Supplemental Material B](#). The literature search yielded a total of 42,065 studies with the cutoff being July 27, 2022.

Inclusion Criteria


Figure 1 presents a graphic depiction of the search and selection strategy. Reviewers first completed multiple rounds of training in order to establish high reliability in screening records for eligibility.

Once the training and searches were complete, each reviewer was assigned a subset of records to independently screen, with each record screened by one reviewer. All records that were initially selected for inclusion and coded were later rechecked by a second reviewer to ensure the inclusion criteria were correctly applied. We applied the following inclusion criteria (Fleiss' $\kappa = 0.70$ for inclusion/exclusion):

1. *Presence of at least two groups*. Reports had to include both a multiple-behavior intervention group and a comparison group that we could use as a reference. Comparison groups were separated into three categories: (a) passive controls (no recommendations); (b) single-behavior interventions (i.e., interventions making one recommendation); and (c) other multiple-behavior interventions (i.e., interventions making two or more recommendations).

Figure 1

Flowchart for Study Screening and Inclusion



Note. HIV = human immunodeficiency virus.

2. *Presence of two or more behavioral recommendations in the interventions and account of behavioral recommendations.* Reports had to include a description of the target intervention with enough details to verify whether multiple recommendations were included in at least one study condition. As an example, in an included report, [Amaro et al. \(2007\)](#) recommended that participants both reduce their substance use and reduce their sexual risk through abstinence. We excluded reports that did not provide a clear enough description of interventions to determine the number of recommendations (5%).
3. *Presence of behavioral recommendations from two or more domains.* Reports had to include an intervention that included recommendations in at least two behavioral domains which included physical activity, diet, smoking, HIV prevention, HIV testing, HIV treatment, alcohol use, and drug use. For example, [Kalichman et al. \(2014\)](#) included behavioral recommendations to reduce alcohol consumption and increase condom use (i.e., one alcohol use recommendation and one HIV prevention recommendation). As another example, [Dushay et al. \(2001\)](#) recommended participants to test for HIV (one testing behavior) and use condoms (one preventive behavior). In contrast, an intervention that recommended people to reduce their fat intake and reduce their sugar intake would not be included because it would have concerned only the dietary domain.
4. *Presence of at least one behavioral or clinical outcome.* Reports had to include at least one behavioral outcome or one clinical outcome. We excluded reports that included only information on behavioral intentions, attitudes, social norms, or other nonbehavioral or nonclinical outcomes.
5. *Presence of appropriate statistics to estimate improvement over time.* Reports had to include enough statistical information to calculate effect sizes representing improvement or worsening over time. Thus, reports had to include outcome values at both the baseline and at least one posttest. When information was reported only for a delayed posttest, reports were excluded (1%). Studies that reported only analyses with statistical controls or other statistical corrections were also excluded.

After applying our inclusion criteria and excluding studies that appeared to meet the criteria, such as secondary reports (e.g., [Hershow et al., 2020](#)) or those focusing on merely one health domain (e.g., [Cunningham et al., 2020](#)), 364 studies contributing 803 groups and 6,819 effect sizes were included in our analyses (see [Supplemental Material C](#), for study group characteristics and [Supplemental Material D](#), for citations of included studies). The groups from a given study typically represent different intervention conditions. However, when reports stratified results by demographics or risk factors (e.g., high- vs. low-risk drinkers; [Dumas et al., 2017](#)), we separated effect sizes for different groups.

Coding of Effect Sizes

All coding and data extraction were conducted via an internally produced online data entry system using a coding and entry manual

developed by the research team. One researcher extracted the means and standard deviations or proportions for each available study outcome. A second researcher then rechecked this data entry for each report.

Our effect size of interest was within-group improvement over time (d) for all intervention-relevant behaviors and clinical outcomes: If a study reported multiple outcomes, then multiple effect sizes were calculated. If there were multiple posttests, then multiple delayed effect sizes were calculated. To calculate d for continuous outcomes, the mean of the pretest measure was subtracted from the mean of the posttest measure, and this difference was then divided by the pooled standard deviation of the means ([Borenstein et al., 2009](#)). To calculate d for proportional outcomes, an odds ratio was calculated by dividing the odds of the behavior at the posttest by the odds of the behavior at the pretest and then converting it into a d by taking the natural log and dividing it by 1.81 ([Borenstein et al., 2009](#)). In cases where the reported proportions were 0s or 1s, 0.005 was added to 0s or subtracted from 1s, respectively, to reduce extreme odds ratio calculations ([Sweeting et al., 2004](#)). All effect sizes were corrected for small-sample size bias ([Hedges & Olkin, 1985](#)) and coded such that positive effect sizes reflect health improvement (e.g., increased testing rates and decreases in unprotected sex). As the correlation between pretest and posttest observations was unknown, we assumed that $r = 0.5$ in calculations ([Morris, 2008](#)). We used corrected estimates and standard errors for studies reporting results from three reports of cluster-randomized trials ([Hedges, 2007](#)). Previous analyses from other domains of the larger data set have found that models are robust to changes in the assumed correlation ([Wilson et al., 2015](#)).

We applied these calculation rules to obtain effect sizes for multiple posttests as well. If a study had three measurements, such as a baseline, an immediate posttest, and a delayed posttest, we calculated two effect sizes: a pretest versus immediate posttest effect size and a pretest versus delayed posttest effect size. We used pretests as the comparison throughout, as change relative to the baseline reflects improvement at each point. Each effect size has an associated time variable indicating whether the comparison group is from an immediate posttest or a delayed posttest as well as a measure of how many days postintervention the measurement was obtained.

The data set included 58 different outcomes, which were categorized as being either behavioral or clinical. Behavioral outcomes are executed by individuals and are observable through direct measurements, whereas clinical outcomes require biomedical samples that are analyzed and tested in a laboratory setting and do not reflect behavior in a direct way. Examples of behavioral outcomes included attending counseling sessions, using condoms, adhering to a medication regimen or a diet, performing physical activity, and smoking. Examples of clinical outcomes were blood glucose level, drug urine analysis, HIV viral load, and death.

The outcomes could also be categorized according to the domains (see [Supplemental Material E](#)). Relevant to lifestyle interventions, in the domain of *diet*, commonly measured outcomes were energy intake (e.g., kcal/week); the intake of specific nutrition components (e.g., carbohydrate, protein, vitamin, water); number of meals per day; compliance with dietary recommendations (e.g., fruit and vegetable intake); and presence of unhealthy eating or overeating. Commonly measured outcomes related to *physical activity* were the presence of daily exercise behaviors, total hours of weekly physical activity, presence of occupational physical activity, compliance with

exercise recommendations, sedentary behaviors, self-monitoring of pulse and blood pressure, energy expenditure in physical activity, and number of hours watching TV. Commonly measured outcomes related to *smoking* were whether one was smoking and the number of cigarettes smoked per day.

In the area of HIV interventions, commonly measured outcomes related to *sexual risk* were whether participants had engaged in risky sexual behaviors, such as not using condoms during sexual intercourse or having multiple sexual partners. Commonly measured outcomes related to *testing* were whether one was tested for HIV or sexually transmitted infections (STIs), and commonly measured outcomes related to *treatment* were whether one received treatment for HIV or STIs, problems adhering to medications, and changes in biomedical indices (e.g., viral load) for receiving antiretroviral therapy.

In the area of alcohol- and substance-use interventions, commonly measured outcomes related to *alcohol use* were the amount of alcohol consumption during a temporal unit (e.g., day); the frequency of excessive alcohol use (e.g., number of excessive drinking days in a month); the number of days being drunk; and the proportion of abstinence days. Commonly measured outcomes related to *substance use* were the amount of substance use (e.g., heroin, opioids, methadone, cocaine, crack) during a temporal unit (e.g., in the past week); the frequency of excessive drug use; the frequency of ecstasy use; the proportion of abstinence days; and the number of times one had sexual intercourse under the influence of drugs.

We calculated change in all the behavioral and clinical outcomes measured in a study (see [Supplemental Material F](#) for sample reports). For example, if a lifestyle intervention recommended exercise and diet but also measured smoking, all three behavioral changes were recorded. In addition, the data set included effect sizes for improvements in different measures corresponding to psychological well-being. Thus, if an intervention condition reported a measure of depression and a measure of coping behavior, which are two psychological well-being measures, the average of the two effect sizes was used to gauge changes in psychological well-being, which was used as a mediator.

Psychological Well-Being

We obtained effect sizes for the following measures related to well-being: (a) depression and mania, reverse-scored through our effect-size calculation (Calgary Depression Scale; [Addington et al., 1993](#); Center for Epidemiological Studies-Depression; [Radloff, 1977](#); Patient Health Questionnaire-9; [Kroenke et al., 2001](#); Clinical Global Impressions Scale-Bipolar Version-Overall and Mania + Depression subscales; [Spearing et al., 1997](#); Montgomery-Åsberg Depression Rating Scale; [Montgomery & Åsberg, 1979](#); Edinburgh Postnatal Depression Scale; [Cox et al., 1987](#); Hamilton Depression Rating Scale; [Hamilton, 1960](#); Beck Depression Inventory; [Beck et al., 1961](#); mental health subscale of the Addiction Severity Index; [McLellan et al., 1992](#)); (b) psychological quality of life (Medical Outcomes Study 36-item short-form health survey; [McHorney et al., 1993](#); Health Promoting Lifestyle scale; [Walker & Hill-Polerecky, 1996](#); EuroQol Visual Analogue Scale; [EuroQol Group, 1990](#); Acceptance and Action Questionnaire-II; [Bond et al., 2011](#); self-reported mindfulness; self-reported trauma); (c) anxiety and stress,

reverse-scored through our effect-size calculation (Symptom Checklist-90; [Derogatis et al., 1973](#); anxiety subscale of the Depression Anxiety Stress Scale-21; [Henry & Crawford, 2005](#); self-report items); (d) social quality of life (Social Functioning item from the Medical Outcomes Study 36-item short-form health survey; [Ware & Sherbourne, 1992](#); community engagement, social support, and social function items; self-reported victimization); (e) positive mood (Positive and Negative Affect Schedule; [Watson et al., 1988](#); Profile of Mood States; [McNair et al., 1971](#)); and (f) coping (high self-reported positive reinterpretation; low self-reported behavioral disengagement).

As mentioned previously, the coding of effect sizes took the form of within-group growth over time. The advantage of this approach as opposed to between-group comparisons is that different trials may employ different types of control groups (i.e., passive vs. active; see [Karlsson & Bergmark, 2015](#)) and that control and intervention groups in our meta-analysis generally have different numbers of recommendations (e.g., 0 and 3 in one study vs. 1 and 2 in another). One advantage of the within-group growth approach is to allow each group to serve as its own control over time and different interventions to be compared as a function of the number of recommendations. Passive and active control groups could receive the coding of 0 and 1, respectively, regarding the number of behavioral recommendations, and different interventions could be coded in terms of the number of behavioral recommendations as well.

Coding of the Number of Recommendations

As was done in prior analyses of earlier, specific subsets of these data ([Dai et al., 2020](#); [Sunderrajan et al., 2021](#); [Wilson et al., 2015](#)), we coded the number of recommendations as the number of specific, observable behaviors participants received for general health promotion in the intervention programs. The recommendations are behavioral goals for intervention recipients to follow. These recommendations were extracted from authors' descriptions of the intervention methodology. For example, [Go et al. \(2015\)](#) encouraged participants to reduce (a) their sexual risk behavior and (b) their injection drug use. Therefore, that condition was coded as having two behavioral recommendations. In another study, [Kypri and McAnally \(2005\)](#) reported an intervention that included four recommendations, namely (a) increasing physical activity, (b) reducing alcohol use, (c) increasing fruit intake, and (d) increasing vegetable intake. The intercoder reliability coefficient for the coding of the number of recommendations was acceptable ($r = .81$).

Coding of Other Moderators

We coded a variety of moderators that can be used to describe the sample and analyze possible covariates. The interrater reliability for each appears in [Table 1](#).

Report Characteristics

We coded (a) publication year; (b) first authors' institution (e.g., college, hospital, research center); (c) first authors' institutional area (e.g., psychology, public health, medicine); (d) source type (e.g., journal article, dissertation); (e) language of the intervention;

Table 1
Description of the Data

Descriptive variable	Variable level	%
General report characteristics		
Academic affiliation ($\kappa = .93$)	University/college	76
	Research center	12
	Hospital/health service	9
	Centers for Disease Control and Prevention	1
	Medical school	2
Institutional area ($\kappa = 1$)	Psychology	8
	Community/public health	20
	Medicine	50
	Other	22
Intervention characteristics		
Recruitment context ($\kappa = 1$)	Hospital/health clinic	54
	Educational	8
	Nonhealth care	25
	Multiple	13
Delivery context ($\kappa = 1$)	Health clinic	75
	Community	8
	Schools	9
	Street	1
	Mass media	2
	Multiple contexts	5
Communication context ($\kappa = .97$)	Face-to-face	87
	Software	3
	Internet	5
	Multiple contexts	5
Delivery format ($\kappa = 1$)	Groups	29
	Individuals	58
	Both	13
Facilitator ($\kappa = .91$)	Doctor/nurse	51
	Public health educator	21
	Clinical psychologist	9
	Community leader	1
	Multiple	18

Note. κ = intercoder reliability for categorical variables. Academic affiliation = the academic affiliation of the first author; Institutional area = the subject area of the first author; Recruitment context = the context for recruiting participants; Exposure setting = the setting of intervention delivery; Delivery medium = the medium used for delivering intervention; Delivery format = the format of delivering intervention; Facilitator = the person delivering the intervention; Informational strategies = the intervention provided informational education to facilitate behavioral change; Motivational strategies = the intervention included motivational strategies to facilitate behavioral change; Instruction in behavioral skills = the intervention included teaching behavioral skills to facilitate behavioral change. The percentage values for levels of each variable were based on all the available observations from a total of 667 independent study samples.

(f) country of the intervention (as well as state or city if provided); and (g) research design.

Sample Features

We recorded the following sample characteristics: (a) the percentage of males in each group; (b) the percentage of male participants who identified as gay or bisexual; (c) the percentage of participants of European, African, Latin, Asian, and Native American descent; (d) the mean age; (e) the country or countries where the study was conducted; and (f) the proportion of the sample with a preexisting

health condition, defined as a previous diagnosis of a relevant disease (e.g., diabetes in lifestyle domain studies).

Method Features

For each study, we recorded factors related to the design and implementation of the intervention, including (a) whether participants were randomly assigned to the intervention and control groups; (b) the mean number of days between the intervention and each posttest; (c) whether participants were recruited in a clinical or nonclinical setting; (d) whether an expert source/facilitator was present; and (e) whether the intervention was delivered to a group, to individuals, or to a combination of the two. We also recorded (f) whether the primary exposure format was face to face or another format such as video; (g) whether the exposure setting was clinical (e.g., a clinic) or nonclinical (e.g., a school); (h) whether the study used intention-to-treat analyses; (i) the total time participants spent participating in the intervention; and (j) whether the authors described the intervention as culturally appropriate, tailored, or neither.

Analyses

General Approach

All the analyses were conducted in R with the α level set at .05. The primary unit of analysis was the effect size d representing behavioral and clinical improvements over time and weighted by its precision, which is the inverse of the variance (Borenstein et al., 2009). Metaregression was adopted as the primary analytical approach. Given that the present data involved 6,819 effect sizes from 803 independent samples/groups from 364 studies and that each independent sample was associated with more than eight effect sizes on average, our primary analysis involved the model of correlated and hierarchical effects with three levels (CHE+; Pustejovsky & Tipton, 2022) using the R packages Metafor (Viechtbauer, 2019) and ClubSandwich (Pustejovsky, 2021) with other analyses (i.e., publication bias, propensity-score modeling, and mediation testing) performed using the packages Robumeta (Soma et al., 2021), ordinal (Christensen, 2011), Psych (Revelle, 2015), and Lavaan (Rosseel, 2012).

The CHE+ model was recently developed to merge the hierarchical approach with the robust variance estimation (RVE) approach in metaregression. This model assumes a more generalized form than both hierarchical and RVE approaches by accounting for effect-size dependence arising from both a nested data structure (e.g., measures nested under trial conditions/samples that are themselves nested under trials) and correlated error variances among the effect sizes from a given sample (e.g., sampling error). In addition, unlike the RVE approach, which relies on moment-of-methods estimation, the CHE+ model uses the restricted maximum likelihood (REML) estimation to obtain parameter estimates. Both empirical and simulated analyses have suggested that the CHE+ models can reach more precise (i.e., smaller standard errors) and accurate (i.e., estimates being closer to the true values) parameter estimates than RVE models (Pustejovsky & Tipton, 2022). All these features made the CHE+ model a good option for the current research.

In specifying a given CHE+ model, in addition to the fixed-effects model part, we included three random-effects parameters at default to account for random variability of the true effects at the level of study (Level 1), sample (Level 2), and measure (Level 3). We also specified the correlation parameter of sampling error (i.e., $\rho = 0.6$) with smoothing of the marginal variances in the RVE-like part of the CHE+ model and used small-sample corrections (McCaffrey & Bell, 2003; Pustejovsky & Tipton, 2018). These general strategies were adjusted according to specific research questions in analyses, which included (a) publication bias determination, (b) covariate identification, (c) propensity-score models, and (d) tests of the main hypotheses.

Publication Bias Analyses

To detect publication bias or selective reporting in the data set, we applied multiple methods, including (a) the funnel plot for direct visualization (Light & Pillemer, 1984); (b) the Egger's Sandwich test to quantify the asymmetry by detecting whether effect sizes are correlated with sample size (Pustejovsky & Rodgers, 2019); and (c) the selection models to determine whether effect sizes were relatively overrepresented in certain range(s) of p -values (Vevea & Hedges, 1995). In the funnel plot, all effect sizes were aggregated to the sample level prior to generating the plots. In the Egger Sandwich test, the effect sizes were not aggregated, and statistical dependence was handled with both the RVE (with $\rho = 0.6$) approach and the hierarchical approach due to a lack of past research with the CHE+ version of Egger's Sandwich test. In these tests, we used the variance term in predicting effect sizes (Rodgers & Pustejovsky, 2021) and calculated a modified version of the variance according to Pustejovsky and Rodgers (2019) to mitigate artifactual association with effect sizes. The selection model relied on data aggregated to the sample level and examined the relative probability density of observing one-sided p values using the following p value cutoff points: .001, .025, .05, .1, .5, and 1. Given that past research has shown the predictive power of the number of behavioral recommendations on effect sizes, we also included the number of behavioral recommendations as a covariate in the fixed-effects part to improve the fit of the selection model.

Controlling for Important Covariates

To control for known covariates that may predict behavioral and clinical outcomes, we conducted CHE+ models to analyze univariate associations between our effect sizes and the following exploratory moderators: (a) whether the effect size was based on clinical or behavioral outcomes, (b) whether the exposure to the intervention was as an individual or as part of a group, (c) whether the intervention was delivered in a clinic or not, (d) whether the intervention facilitator was a professionally trained expert, (e) whether the intervention involved motivational strategies, (f) whether the effect size was calculated using proportional outcomes (and later transformed into Cohen's d metric), (g) a socioeconomic status variable indicating the proportion of participants completing high school education, (h) the standardized duration of the intervention program, and (i) the standardized time interval between baseline and posttest measurement. Most of these covariates were suggested by previous meta-analyses synthesizing domain-specific evidence (cf. Dai et al., 2020; Sunderrajan et al., 2021; Wilson et al., 2015) and, if

they predicted effect sizes, were included in the CHE+ models that tested how the number of behavioral recommendations affected behavioral and clinical change.

Propensity Score Model

To further improve inferences about the effects of the number of behavioral recommendations while controlling for possible covariate differences among studies with different recommendation numbers, the CHE+ models included a scalar version of the propensity score as another covariate (Joffe & Rosenbaum, 1999). The propensity scores were estimated from a generalized linear mixed-effects model, wherein linear combinations of available covariates were introduced as predictors of the number of behavioral recommendations via a logistic link function. The number of behavioral recommendations was treated as an ordinal outcome including zero recommendations, one recommendation, two recommendations, three recommendations, and four or more recommendations.¹ The propensity-score model was based on 803 condition-level observations and included a random effect at the trial level. Overall, developing the propensity-score model was an iterative process. The model selection decision was based on criteria that included (a) the degree of balancing covariates among the number of behavioral recommendations on each quintile of the propensity score and (b) the area of common support regarding the propensity-score distribution. Also, preference was given to (c) models with a relatively simple linear combination of covariates (rather than interactions and quadratics; see Joffe & Rosenbaum, 1999) and (d) models preserving observations during the trimming process. The procedures generally followed the recommendations outlined in Bergstra et al. (2019).

Analyses Testing Main Hypotheses

Using data associated with the untrimmed trial conditions from the propensity-score model and those from all the available observations, we implemented the CHE+ model to examine associations between the number of behavioral recommendations and behavioral and clinical improvements. We hypothesized that tapping a general process could result in a monotonic association between the number of behavioral recommendations and behavioral and clinical change, mathematically expressed by a linear term in the CHE+ model. In contrast, the number of behavioral recommendations may predict improvements only up to a point, approximating an inverted- U shape mathematically expressed by a quadratic term in the CHE+ model. To avoid collinearity between the linear and quadratic terms of the number of behavioral recommendations, we centered the linear term before squaring it to generate the quadratic term.

We then implemented two CHE+ models. The first model included the linear term and all the covariates, and the second included the linear and quadratic terms along with the same set of covariates. As a reminder, the covariates included an indicator variable for the type of effect size (i.e., clinical outcome vs. behavioral outcome) in the model; the covariates associated with intervention features; a modified version of effect-size variance;

¹ The propensity score model represents the only occasion where the number of recommendations was categorized for the sake of modeling. In all other models, the number of recommendations was treated as a continuous variable.

and the propensity score estimates. If clinical outcomes were shown to be different from behavioral ones, we would then explore the relation with the number of behavioral recommendations separately for each outcome.

Evaluation and Sources of Heterogeneity

To gauge the heterogeneity of our effects, we obtained Q , I^2 , and 95% prediction interval (PI) of the base model, which is a CHE+ model with only the intercept term in the fixed-effects part. We computed I^2 by dividing the estimated study-level variance (i.e., σ^2_{τ}) by the total variance estimated from all three levels of data hierarchy and computed 95% PIs using the estimated average effects and the study-level variance in the base model. I^2 is a statistic that indicates the proportion of heterogeneity, whereas the 95% PI is an absolute description of effect-size dispersion (Borenstein et al., 2017). When substantial heterogeneity was identified, we performed exploratory analyses to investigate sources of heterogeneity.

Analyses of Our Hypotheses

We began to test mechanisms by examining cuing, which assumes that more behavioral recommendations can potentiate each other when one behavior cues other behaviors. For these analyses, we aggregated effect sizes to the sample level and created the categories of (a) exercise, (b) diet, (c) smoking, (d) HIV (including prevention, testing, and treatment), (e) alcohol use, and (f) drug use. Because the literatures tend to intervene on exercise, diet, and smoking together (e.g., an intervention to reduce cardiovascular disease), and similarly on HIV, alcohol use, and drug use together, these behaviors present two literature groups, including lifestyle (subsuming exercise, diet, and smoking) as well as HIV, alcohol, and substance use. We tested the cuing effect among behaviors within the lifestyle literature and within the HIV, alcohol, and substance-use literatures. In these analyses, specific behaviors measured at the immediate posttest were set to predict other behaviors measured at the delayed posttest, while controlling for potential selection bias and the interval of time between the baseline and posttest measurements. Because each sample generated only one observation, the cuing-testing models included two random effects that corresponded to the study and sample levels, respectively. For instance, in the model testing whether exercise would cue diet and smoking, changes associated with exercise at the immediate posttest as well as time interval were included in the two-level metaregression model to predict changes in diet and smoking at the delayed posttests.

We also tested whether the effects of the number of recommendations on behavioral and clinical changes were mediated by changes in psychological well-being, information, motivation, and behavioral skills. We followed the three-step tradition in testing an “X-M-Y” mediational chain. The steps included testing (a) X-Y (i.e., the number of behavioral recommendations predicting effect sizes); (b) X-M (i.e., the number of behavioral recommendations predicting the mediator); and (c) XM-Y (i.e., both the number of behavioral recommendations and the mediator predicting behavioral and clinical improvements). We focused on the product of the recommendation-mediator path coefficient in Step X-M and the mediator-behavioral/clinical change path coefficient in Step XM-Y because such a product indicates mediation strength (i.e., indirect effect) without involving

the measurement unit of the mediator (Hayes & Rockwood, 2017). Three CHE+ models were fit to immediate and delayed changes and each CHE+ model’s fixed-effects part corresponding to a given step of the mediational chain with a covariate indicating the elapse of time (in days) between the intervention and the dependent measure. When a given mediator showed a meaningful magnitude of mediation strength (i.e., indirect effect), we further conducted tests of corresponding indirect effect(s) using Sobel tests (Sobel, 1982) and path analyses with both bootstrapped confidence intervals from 5,000 iterations (Shrout & Bolger, 2002) and Huber–White robust standard error (Huber, 1967; White, 1980).

Transparency and Openness

The preregistration of the review’s protocol is at https://osf.io/mq5kr/?view_only=54d04d57b5ef45aaa480a47259cb91cd. We prepared the protocol by following the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols checklist and reported our work by following the PRISMA guidelines (see [Supplemental Material L](#)). The R code and data supporting the findings reported in this study are openly available in the Open Science Framework at https://osf.io/qsx57/?view_only=6ffffd7080d844e98cbd2d0834ed61e9. The present article includes deviations from the preregistration. The major deviation involves using the more recent and more generalized metaregression models of correlated and hierarchical effects (CHE) instead of the preregistered robust variance estimation (RVE). In addition, a couple of questions focusing on the refined effect of specific recommendation combinations, sample characteristics, and methodological factors on intervention efficacy could not be fully tested and reported due to the constraints arising from both data availability and the scope of the present article. Details about specific deviations are described in [Supplemental Material A](#).

Results

Description of Included Studies, Samples, and Effect Sizes

[Table 1](#) presents a description of our data. Overall, we included a total of 6,878 effect sizes (with positive values indicating improvement in the direction of healthy behavior and clinical outcomes) from 803 independent groups from 364 studies that sampled 186,729 participants. The effect sizes included 4,282 immediate postintervention effect sizes from 781 independent groups from 355 studies with an average trial-onset-to-measurement time of 215 ($SD = 292$) days and 2,596 delayed postintervention effect sizes from 378 independent groups from 171 studies with an average trial-onset-to-measurement time of 385 ($SD = 267$) days. The large majority (96%) involved random group assignment. The median group contributed five effect sizes, and the average group contributed 8.6 effect sizes. The average number of recommendations across all groups was 2.50, and the average number of recommendations among multiple-behavior interventions was 3.37. All the studies were reported in English except for one study reported in Korean and another study reported in Italian. Nearly all (99%) were from journal articles having a median publication year of 2016 (range: [1979, 2022]), with the remainder coming from doctoral dissertations. Sample sizes varied widely ($M_{\text{size}} = 513$, $SD_{\text{size}} = 1,295$), and the average sample was racially/ethnically

diverse, with an average racial/ethnic background of 46% non-Hispanic White, 27% African, 16% Asian, 13% Latinx, 1% Native American, and 14% other or multiple. The average sample age was 40 ($SD = 16$) years old and had good representation of males and females (57% female, $SD = 31\%$), with 45% completing high school or an equivalent. Among the 364 included studies, most were conducted in the United States (51.4%), with much smaller proportions elsewhere: United Kingdom (5.5%), The Netherlands (3.6%), Australia (3%), Brazil (2.5%), Canada (2.5%), China (2.5%), Japan (2.2%), Germany (1.9%), India (1.9%), and Spain (1.9%). [Supplemental Material G](#) lists details of study distribution over countries and how each country is categorized into Western versus non-Western countries ([World Population Review, 2023](#)).

Assessment of Bias

Figure 2 displays funnel plots regarding the effect-size distribution according to the precision measure of standard error after aggregating effect sizes within each sample. A visual inspection suggests no clear asymmetry of effect-size distribution. Beyond the funnel plots, results from the RVE and hierarchical variants of the Egger's Sandwich test suggested no evidence of effect-size distribution asymmetry in either the hierarchical version ($B = -0.26$, $SE = 0.13$, $p = .98$) or the RVE version ($B = -0.09$, $SE = 0.11$, $p = .78$) of the test.


In addition, results from the selection model showed that, compared to the unadjusted model, the adjusted model fit significantly better ($p < .001$) by including the relative probability density weights associated with the p value intervals with higher weights suggesting a stronger tendency to report findings for particular ranges of p values.

Specifically, the following weights (which are all significant at $p < .001$) were estimated: $B = 2.56$ for p between .001 and .025; $B = 2.57$ for p between .025 and .05; $B = 2.23$ for p between .05 and .1; $B = 3.65$ for p between .1 and .5; and $B = 0.98$ for p between .5 and 1. Because the first p value interval (i.e., from 0 to .001) was used as a reference by constraining it to have a weight of 1, this pattern of results did not suggest a clear bias toward including more effect sizes associated with relatively smaller p values, as the weights are generally balanced among all categories. In sum, the assessments did not evidence bias in the present data, although we included the modified variance ([Pustejovsky & Rodgers, 2019](#)) as a covariate in all the subsequent meta-regressions to enhance our estimates of the moderators of interest.

Tests to Identify Important Covariates

Results from the preliminary CHE+ models suggested that clinical outcomes were associated with smaller effect sizes than behavioral outcomes ($B = -0.09$, $SE = 0.03$, $p = .004$). In addition, among important covariates related to method features, individual or group exposure to interventions ($B = 0.02$, $SE = 0.03$, $p = .36$) and interventions including an expert facilitator ($B = 0.03$, $SE = 0.03$, $p = .30$) did not significantly moderate behavioral and clinical improvement. The socioeconomic variable indicating the proportion of participants completing high school education ($B = 7e-4$, $SE = 6e-4$, $p = .24$) did not significantly predict behavioral and clinical improvement either. In contrast, interventions that took place in a clinical (vs. nonclinical) setting ($B = 0.07$, $SE = 0.03$, $p = .02$), interventions that used motivational strategies ($B = 0.30$, $SE = 0.04$,

Figure 2
Funnel Plot of Aggregated Sample-Level Effect Sizes



Note. The plot was generated using aggregated sample-level effect sizes (to avoid data nonindependence issue). The dotted vertical line corresponds to the random-effects model estimate (0.32), and the funnel shape corresponds to the 95% pseudoconfidence interval.

$p < .001$), longer intervention durations ($B = 0.06$, $SE = 0.02$, $p = .01$), a shorter time interval between baseline and posttest ($B = -0.03$, $SE = 0.01$, $p = .005$), and proportional outcomes ($B = 0.27$, $SE = 0.08$, $p < .001$) were associated with greater behavioral and clinical improvements. As a result, we included these significant confounds in the CHE+ models that tested the impact and processes of the number of behavioral recommendations. Yet, these covariates could not be included in the analyses of cuing and mediational processes because they led to very small numbers of observations (i.e., $k < 70$; for recommendations about minimum sample sizes, see Tanner-Smith & Tipton, 2014).

Estimates of Propensity Scores

The selection of variables for the propensity-score model was based on the criterion that the variables must be considered as “baseline covariates” and cannot be affected by the number of behavioral recommendations, which is the “treatment.” Checking covariate balance among five categories of the number of recommendations required dividing the trial conditions into quintiles based on the propensity score, resulting in $5 \times 5 = 25$ combinations. Fifteen out of 132 categorical covariates were selected given low missing rates (i.e., $< 8\%$) and sufficient observations (i.e., > 100) in each category of the covariate. Table 2 shows more details about the covariates. The final propensity-score model consisted of regressing

Table 2
Covariates Included in the Propensity Score Model

Covariate no.	Covariate	Category no.
Covariate 1	Whether the intervention targeted a particular gender group	2
Covariate 2	Whether participants self-selected into the study	2
Covariate 3	Whether the intervention used attitudinal arguments	2
Covariate 4	Whether the intervention used informational arguments	2
Covariate 5	Whether the intervention taught behavioral skills	2
Covariate 6	Whether the intervention showed what to do when facing behavioral barriers	2
Covariate 7	Whether the intervention used self-monitoring prompts	2
Covariate 8	Whether the study included at least one follow-up before 6 months	2
Covariate 9	Whether the intervention aimed to improve lifestyle behaviors	2
Covariate 10	Whether the intervention aimed to improve HIV behaviors	2
Covariate 11	Whether the intervention aimed to improve substance-use behaviors	2
Covariate 12	Whether the intervention occurred in North America	2
Covariate 13	Whether the study included more than one posttest	2
Covariate 14	Time of the publication (categorized into pre-2003, 2003–2010, post-2010)	3
Covariate 15	Percentage of males in the sample (categorized into 4 percentage quarters)	4

Note. Covariate no. corresponds to those shown in Supplemental Material H. HIV = human immunodeficiency virus.


the number of behavioral recommendations on the main effects of these covariates with a random effect at the trial level. Figure 3 illustrates the contrast prior to and after trimming the trial conditions given the estimated propensity scores. Further details regarding the covariate balance conditional on propensity-score quintiles can be found in Supplemental Material H. Overall, 203 trial conditions were trimmed off in identifying the area of common support with the propensity score ranging from $[-6, 0]$. This process reduced the total number of trial conditions from 803 to 600, leading to 120 trial conditions in each final quintile of the propensity score.

Effect of the Number of Recommendations in an Intervention

Given potential differences between behavioral and clinical effect sizes, we first tested the linear versus nonlinear hypothesis separately for behavioral and clinical improvements. Table 3 shows the CHE+ model results, indicating a positive, linear effect but not a quadratic effect ($p > .13$) of the number of behavioral recommendations on both behavioral and clinical improvements. Table 4 shows the main CHE+ model results, indicating that the linear form ($B = 0.03$, $SE = 0.01$, $p < .001$) but not the quadratic form ($B = -2e-3$, $SE = 1e-3$, $p = .12$) of the number of behavioral recommendations predicted effect sizes after controlling for the confounds. Figure 4 illustrates the positive linear effect of the number of behavioral recommendations on behavioral and clinical improvements. In addition, the CHE+ models that did not include the propensity scores as a covariate (and thus used 781 trial conditions) generated a larger estimate of the positive, linear effect of the number of behavioral recommendations ($B = 0.04$, $SE = 0.007$, $p < .001$), suggesting that the propensity scores were able to attenuate an upward bias in such an estimation. The parallel effects of the number of behavioral recommendations on behavioral and clinical outcomes supported treating the two types of outcomes together in subsequent analyses. This strategy was also desirable to increase the sample size available for each analysis.

Finally, we performed three sensitivity analyses. The first sensitivity analysis followed the difference-in-differences approach by using between-group effect sizes. Results consistently supported the linear ($B = 0.03$, $SE = 0.01$, $p = .001$) but not the quadratic term ($B = -0.002$, $SE = 0.001$, $p = .12$) of the number of behavioral recommendations (see Supplemental Material I). The second sensitivity analysis gauged whether the present findings are robust against the concern that psychology findings are overly reliant on WEIRD (Western, Educated, Industrialized, Rich, and Democratic, see Henrich et al., 2010) samples. Results suggested that the linear effect of the number of recommendations was significant after including an indicator variable for whether participants resided in a Western versus non-Western country as well as a socioeconomic status variable indicating the proportion of participants completing high school education (see Supplemental Material J). The last sensitivity analysis helped test the robustness of findings in Table 4 against three indicator variables related to study quality, including (a) whether the study randomized participants/clusters into conditions, (b) whether the study included a posttest at least 6 months after the baseline measurement, and (c) whether the study involved an intent-to-treat analysis. All results showed consistent findings regarding the linear effect of the number of recommendations (see Supplemental Material K).

Figure 3
Propensity Score Distributions Prior to and After Trimming



Note. Propensity score distributions are shown for each level of the treatment (i.e., number of behavioral recommendations) prior to (Panel A) and after (Panel B) trimming for the area of common support (i.e., ranges on X-axis shared by all the distributions). *k* represents the number of trial conditions in each distribution. See the online article for the color version of this figure.

Evaluation and Sources of Heterogeneity

We evaluated effect-size heterogeneity of the model in Table 4. Results supported the presence of a small-to-moderate level of heterogeneity, $\tau = 0.30$, $Q(2,742) = 70770.62$, $p < .001$, $I^2 = 28.59\%$. The 95% PI $[-0.29, 0.88]$ showed a wide range of effects in the literature we synthesized.

To explore the sources of heterogeneity, we identified seven additional study-level predictor variables in addition to the number of recommendations, three related to the study population and four related to the study design. Table 5 shows three nested models fit to the same data, including (a) an average-effect model, (b) a model that adds the number of recommendations to the average-effect model, and (c) a model adding the seven additional predictors. Table 5 also shows the percentage of reduction in heterogeneity for each model after adding the predictor variables to the average-effect model. As shown, the number of recommendations alone accounted for 7.23% of the estimated heterogeneity. Moreover, adding the additional seven predictors to the model with number of recommendations accounted for up to 29.31% of heterogeneity. In particular, studies that included a higher number of recommendations ($B = 0.03$, $SE = 0.01$, $p = .002$), studies with longer intervention duration ($B = 0.05$, $SE = 0.02$, $p = .01$), and studies conducted in non-Western countries ($B = -0.19$, $SE = 0.09$, $p < .05$), were

associated with larger effect sizes, which suggested some important sources of heterogeneity.

Tests of Potential Mechanisms

We tested possible behavioral cuing as well as well-being and motivation as mediators of the linear effect of the number of recommendations included in an intervention.

Behavioral Cuing

Table 6 shows the results from the CHE+ models regarding behavioral cuing. These analyses considered, after controlling for potential selection bias and the time of measurement for both the cuing and the cued behaviors, the degree to which a specific behavior (e.g., exercise) predicts other behaviors in the study (e.g., diet and smoking), investigating whether changes in that specific behavior potentiate changes in the other behaviors. Results indicated that, after controlling for selection bias and the time of measurement, changes in HIV ($B = 0.70$, $SE = 0.18$, $p = .01$), alcohol use ($B = 0.48$, $SE = 0.10$, $p < .001$), and drug use ($B = 0.35$, $SE = 0.10$, $p = .02$) showed significant cuing effects, whereas changes in exercise ($B = 0.27$, $SE = 0.22$, $p = .26$), diet ($B = 0.14$, $SE = 0.06$, $p = .10$), and smoking ($B = 0.02$, $SE = 0.34$, $p = .97$) did

Table 3*Effect of Number of Recommendations on Behavioral and Clinical Outcomes*

Variable	Behavioral outcome				Clinical outcome			
	Linear (<i>n</i> = 233; <i>k</i> = 402; <i>s</i> = 1,501)		Curvilinear (<i>n</i> = 233; <i>k</i> = 402; <i>s</i> = 1,501)		Linear (<i>n</i> = 186; <i>k</i> = 344; <i>s</i> = 1,616)		Curvilinear (<i>n</i> = 186; <i>k</i> = 344; <i>s</i> = 1,616)	
	<i>B</i> (<i>SE</i>)	<i>p</i>	<i>B</i> (<i>SE</i>)	<i>p</i>	<i>B</i> (<i>SE</i>)	<i>p</i>	<i>B</i> (<i>SE</i>)	<i>p</i>
Fixed effects								
Intercept	0.28 (0.1)	.006**	0.37 (0.1)	.001***	0.11 (0.08)	.19	0.2 (0.09)	.03*
Effect-size variance	−0.42 (0.29)	.16	−0.4 (0.29)	.18	−0.02 (0.21)	.92	−0.02 (0.2)	.91
Propensity score	−0.04 (0.02)	.04*	−0.03 (0.02)	.11	−0.02 (0.01)	.16	−0.01 (0.01)	.62
Number of recommendations	0.02 (0.01)	.04*	0.03 (0.01)	.009**	0.02 (0.01)	.01*	0.04 (0.01)	.001***
Squared number of recommendations			−0.005 (0.003)	.16			−0.003 (0.001)	.13
Random effects								
σ_1^2 (studies)	0.16		0.16		0.08		0.07	
σ_2^2 (samples)	0.00		0.00		0.00		0.00	
σ_3^2 (measures)	0.21		0.21		0.14		0.14	

Note. *n* is the number of studies; *k* is the number of unique groups; *s* is the total number of effect sizes. σ_1^2 is the random component (variance) at the study level; σ_2^2 is the random component (variance) at the sample level; σ_3^2 is the random component (variance) at the measure level. Passive controls in the trials correspond to the number of recommendation being 0's and active controls 1's. *SE* = standard error.

* $p < .05$. ** $p < .01$. *** $p < .001$.

not demonstrate cuing. The overall pattern suggested significant cuing across behaviors in the HIV and substance-use literatures.

Changes in Well-Being

Along with cuing, changes in well-being, but not motivation, could explain the linear effects of the number of behavioral

recommendations on behavioral and clinical change. Table 7 summarizes the results from the CHE+ model testing psychological well-being, information, motivation, and behavioral skills, respectively, as mediators of the effect of the number of behavioral recommendations on behavioral and clinical improvement. As shown, only psychological well-being and motivation showed promise, as the number of behavioral recommendations predicted improvement in

Table 4*Effects of the Number of Recommendations*

Variable	Linear model (<i>n</i> = 291, <i>k</i> = 501, <i>s</i> = 2,743)		Curvilinear model (<i>n</i> = 291, <i>k</i> = 501, <i>s</i> = 2,743)	
	<i>B</i> (<i>SE</i>)	<i>p</i>	<i>B</i> (<i>SE</i>)	<i>p</i>
Fixed effects				
Intercept	0.08 (0.09)	.34	0.16 (0.09)	.08
Effect-size variance	0.10 (0.22)	.65	0.1 (0.22)	.65
Propensity score	−0.02 (0.02)	.22	−0.02 (0.02)	.30
Number of recommendations	0.02 (0.01)	.009**	0.03 (0.01)	.001***
Squared number of recommendations			−2e−3 (1e−3)	.12
Motivation ^a	0.11 (0.08)	.14	0.10 (0.08)	.22
Clinic context ^a	0.01 (0.04)	.82	2e−3 (0.04)	.95
Clinical outcome ^a	−0.07 (0.04)	.13	−0.07 (0.04)	.13
Proportion outcome ^a	0.36 (0.1)	.001***	0.36 (0.1)	.001**
Duration ^b	0.03 (0.02)	.30	0.02 (0.02)	.39
Time ^b	−0.03 (0.02)	.08	−0.03 (0.02)	.09
Random effects				
σ_1^2 (studies)		0.10		0.09
σ_2^2 (samples)		0.00		0.00
σ_3^2 (measures)		0.21		0.21


Note. Motivation = intervention contained motivational strategies; Clinical context = whether the intervention took place in a clinical or nonclinical setting; Clinical outcomes = whether the effect size is clinical or behavioral; Proportion outcome = The calculation of effect size for the outcome is based on proportions (vs. mean and standard deviations); Duration = the duration of the intervention program; Time = the time interval between baseline and posttest measurement. *n* is the number of studies; *k* is the number of unique groups; *s* is the total number of effect sizes. σ_1^2 is the random component (variance) at the study level; σ_2^2 is the random component (variance) at the sample level; σ_3^2 is the random component (variance) at the measure level. Passive controls in the trials correspond to the number of recommendation being 0's and active controls 1's. *SE* = standard error.

^a Indicator variable. ^b Standardized variable.

** $p < .01$. *** $p < .001$.

Figure 4

The Effect of Number of Recommendations on Clinical and Behavioral Improvements



Note. The red regression line (with the shade representing 95% CI) is based on the CHE+ model shown in the left panel of Table 4, while keeping other covariates constant at their grand means. The effect sizes are plotted with different colors to indicate the associated number of behavioral recommendations and with different sizes to indicate weight (i.e., inverse variance). CI = confidence interval; CHE = correlated and hierarchical effects. See the online article for the color version of this figure.

psychological well-being ($B = 0.12$, $SE = 0.05$, $p = .02$) and motivation ($B = 0.07$, $SE = 0.03$, $p = .05$), and well-being ($B = 0.35$, $SE = 0.18$, $p = .107$) and motivation ($B = 0.47$, $SE = 0.13$, $p = .01$) also seemed to predict behavioral/clinical improvement when the

number of recommendations was included. Conversely, in the follow-up tests of the indirect effect (i.e., product of the two paths), only well-being was evidenced as a mediator in all three tests in Table 8, including the Sobel test ($Z = 3.35$, $p < .001$); the path analytic model using bootstrapped CI ($B = 0.05$, $SE = 0.02$, $p < .001$); and the path analysis using robust standard error ($B = 0.05$, $SE = 0.02$, $p < .001$). Motivation was not a mediator ($p > .18$). These results suggested that adding one recommendation to the number of behavioral recommendations would lead to a $d = 0.042$ increment in behavioral and clinical improvement at the posttest via improvements in psychological well-being.

Discussion

Employing the recent CHE+ models to account for the complex dependence structure among effect sizes from 364 study reports, the present meta-analysis investigated how and why the number of behavioral recommendations in an intervention affected clinical and behavioral changes in eight domains related to lifestyle, HIV, and substance use. Results indicated a positive, linear (but not curvilinear) effect of the number of recommendations on behavioral and clinical changes after controlling for confounding factors from several sources, including selective reporting (i.e., modified variance of the effect size); conceptually important covariates (e.g., whether the effect size is associated with a behavioral or clinical measure); and 15 other covariates empirically derived and controlled for by introducing the propensity score (see Table 4 and Figure 4). These results thus offer both stronger and more generalizable causal inference about the impact of behavioral recommendations than

Table 5
Exploratory Models for Identifying Sources of Heterogeneity

Variable	Average-effect model		Model with number of recommendations as a predictor ($n = 139$, $k = 233$, $s = 1,238$)		Model with all predictor	
	B (SE)	p	B (SE)	p	B (SE)	p
Fixed effects						
Intercept	0.28 (0.03)	.001***	0.16 (0.04)	.001***	0.47 (0.11)	.001***
Number of recommendations			0.04 (0.01)	.001***	0.03 (0.01)	.002**
Duration ^b					0.05 (0.02)	.01*
Western countries ^a					−0.19 (0.09)	.05*
HIV ^a					−0.12 (0.12)	.30
Substance use ^a					−0.17 (0.08)	.06
Mean age ^b					0.04 (0.03)	.13
Percent males ^b					0.02 (0.03)	.48
Year of study onset ^b					0.04 (0.03)	.17
Sample size ^b					−0.14 (0.07)	.12
Random effects						
σ_1^2 (studies)	0.0788		0.0731		0.0557	
σ_2^2 (samples)	0.0000		0.0000		0.0000	
σ_3^2 (measures)	0.1924		0.1912		0.1918	
Reduction of heterogeneity			7.23%		29.31%	

Note. Western country = whether the residence of participants is in western versus non-Western countries; HIV = whether the study is from the HIV domain or not; Mean age = the mean age of the study sample; Percent males = the percentage of males in the study sample; Year of study onset = the year when the study began; Sample size = the study sample size; Duration = the duration of the intervention program. σ_1^2 is the random component (variance) at the study level; σ_2^2 is the random component (variance) at the sample level; σ_3^2 is the random component (variance) at the measure level. Passive controls in the trials correspond to the number of recommendation being 0's and active controls 1's. SE = standard error; HIV = human immunodeficiency virus.

^a Indicator variable. ^b Standardized variable.

* $p < .05$. ** $p < .01$. *** $p < .001$.

Table 6*Tests of Behavioral Change Cuing Other Behavioral Changes*

Variable	Exercise (<i>n</i> = 29, <i>k</i> = 55)			Diet (<i>n</i> = 27, <i>k</i> = 54)			Smoking (<i>n</i> = 11, <i>k</i> = 19)			HIV (<i>n</i> = 16, <i>k</i> = 35)			AU (<i>n</i> = 18, <i>k</i> = 38)			DU (<i>n</i> = 19, <i>k</i> = 41)		
	<i>B</i> (<i>SE</i>)	<i>p</i>		<i>B</i> (<i>SE</i>)	<i>p</i>		<i>B</i> (<i>SE</i>)	<i>p</i>		<i>B</i> (<i>SE</i>)	<i>p</i>		<i>B</i> (<i>SE</i>)	<i>p</i>		<i>B</i> (<i>SE</i>)	<i>p</i>	
Fixed effects																		
Intercept	0.16 (0.15)	.28		0.11 (0.08)	.19		0.43 (0.54)	.47		0.48 (0.26)	.10		0.16 (0.08)	.09		0.29 (0.14)	.06	
Effect-size variance	0.01 (0.44)	.98		0.06 (0.33)	.86		−1.75 (2.54)	.52		−0.06 (0.85)	.95		−0.35 (0.3)	.33		−0.33 (0.32)	.37	
Effect size of specific behavior	0.27 (0.22)	.26		0.14 (0.06)	.10		0.02 (0.34)	.97		0.7 (0.18)	.01*		0.48 (0.1)	.001***		0.35 (0.1)	.02*	
Cueing behavior time ^a	0 (0.09)	.96		0.03 (0.04)	.40		0.11 (0.09)	.29		−0.28 (0.26)	.34		−0.1 (0.12)	.44		0.2 (0.09)	.07	
Cued behavior time ^a	−0.04 (0.1)	.68		−0.14 (0.07)	.08		−0.18 (0.22)	.47		0.73 (0.42)	.18		0.02 (0.08)	.81		−0.34 (0.07)	.04*	
Random effects																		
σ_1^2 (studies)	0.20			0.05			0.29			0.09			0.04			0.11		
σ_2^2 (samples)	0.04			0.01			0.04			0.00			0.00			0.00		

Note. Cueing behavior time = time interval between baseline and the cueing behavior measured at immediate posttest; Cued behavior time = time interval between baseline and the cued behavior measured at delayed posttest; Domains include lifestyle (exercise, diet, smoking) as well as one for HIV, AU, and DU. *n* is the number of studies; *k* is the number of unique groups; *s* is the total number of effect sizes. σ_1^2 is the random component (variance) at the study level; σ_2^2 is the random component (variance) at the sample level. Passive controls in the trials correspond to the number of recommendation being 0's and active controls 1's. HIV = human immunodeficiency virus; AU = alcohol use; DU = drug use; *SE* = standard error.

^aStandardized variable.

* *p* < .05. *** *p* < .001.

previous work (cf. Dai et al., 2020; Meader et al., 2017; C. R. Nigg & Long, 2012; Prochaska & Prochaska, 2011; Sunderrajan et al., 2021; Wilson et al., 2015). In addition, the findings about behavioral cuing and psychological well-being point to general mechanisms that have not been previously identified and are key to designing future programs. We discuss these findings and their implications in further detail subsequently.

The Effect of the Number of Behavioral Recommendations

We developed a propensity-score model to better isolate the impact of the number of behavioral recommendations, which could covary with other factors (Cochran & Chambers, 1965). The propensity-score model allows researchers with observational data to approximately equate trial conditions, such as, in our case, groups with different numbers of behavioral recommendations (Austin, 2011). Overall, the propensity-score model suggested some differences across trial conditions receiving zero, one, and higher numbers of behavioral recommendations (Figure 3A). Trimming trial conditions served to reduce but not fully eliminate the differences across the numbers of recommendations (Figure 3B). Likewise, comparing the CHE+ model parameters with and without covarying propensity scores (0.03 vs. 0.04) revealed an upward bias when propensity scores were not considered, which can result in false positive findings or exaggerated efficacy claims. Nevertheless, the propensity-score-adjusted estimate still suggested that each behavioral recommendation contributes to positive behavioral and clinical change. For example, recommending four behaviors would result in approximately 0.10 *SDs* of clinical and behavioral improvement over not recommending any behavioral recommendations.

Figure 4 illustrates the positive linear effect of the number of recommendations on behavioral and clinical outcomes. As shown, although the number of observations is smaller as the number of recommendations exceeds four, no clear pattern of a curvilinear relation appeared in our data. This finding of a linear effect of the

number of recommendations on behavioral and clinical outcomes is consistent with those of prior meta-analyses of HIV (Sunderrajan et al., 2021) and substance-use (Dai et al., 2020) interventions, although one meta-analysis from the lifestyle domain suggested an inverted-U relationship (see Wilson et al., 2015). Regarding the causes of such mixed findings, despite the inclusion of different domains and study reports, the differences may be due to different analytic methods. Specifically, Wilson et al. (2015) used an analysis of variance (ANOVA)-like approach by grouping recommendation numbers into four categories (i.e., 0, 1, 2–3, 4+) and observed a small efficacy decrease from the 2 to 3 category to the 4+ category, whereas other meta-analyses supporting the linear pattern employed the multivariate metaregression approach to directly test the effect of the number of recommendations. We think this linear-versus-nonlinear issue will be clarified with future intervention studies including a relatively high number of recommendations beyond four. The current body of evidence supports a linear instead of a curvilinear effect of the number of recommendations on behavioral and clinical outcomes.

The linear effect of the number of recommendations can also help to understand heterogeneity in the synthesized literature. Based on the estimated I^2 and 95% PI, between-study heterogeneity constitutes 28.59% of the total variance, and the expected intervention effect can range from −0.29, which is a small-to-moderate negative effect, to 0.88, which is a large positive effect. To explore the sources of this heterogeneity, we considered the number of recommendations and seven additional predictors related to study populations and methodologies. As shown in Table 5, we found that the number of recommendations alone accounted for up to 7.23% of the synthesis heterogeneity, and that higher numbers of recommendations, longer intervention duration, and involving participants from non-Western countries were associated with greater intervention efficacy. These initial findings thus suggested three methodological factors contributing to heterogeneity, which may help guide future study designs (Melsen et al., 2014).

Although the linear effect of the number of recommendations was robust to the inclusion of several research-quality factors in

Table 7*Mediation Analysis for Psychological Well-Being, Information, Motivation, and Behavioral Skill*

Variable	Psychological well-being (<i>n</i> = 43, <i>k</i> = 91, <i>s</i> = 136)		Information (<i>n</i> = 16, <i>k</i> = 32, <i>s</i> = 41)		Motivation (<i>n</i> = 19, <i>k</i> = 39, <i>s</i> = 56)		Behavioral skill (<i>n</i> = 27, <i>k</i> = 55, <i>s</i> = 74)	
	<i>B</i> (<i>SE</i>)	<i>p</i>	<i>B</i> (<i>SE</i>)	<i>p</i>	<i>B</i> (<i>SE</i>)	<i>p</i>	<i>B</i> (<i>SE</i>)	<i>p</i>
Step 1: <i>X</i> → <i>Y</i>								
Fixed effects								
Intercept	0.03 (0.08)	.73	0.01 (0.2)	.96	0.02 (0.12)	.84	0 (0.09)	.98
Effect-size variance	0.33 (0.25)	.20	0.73 (0.76)	.38	0.25 (0.44)	.58	0.52 (0.37)	.21
Time ^a	0.02 (0.06)	.74	−0.11 (0.06)	.27	−0.05 (0.02)	.13	−0.04 (0.02)	.23
Number of recommendations	0.06 (0.01)	.001**	0.05 (0.03)	.22	0.05 (0.02)	.05	0.05 (0.01)	.006**
Random effects								
σ_1^2 (studies)	0.04		0.10		0.07		0.06	
σ_2^2 (samples)	0.01		0.06		0.02		0.01	
σ_3^2 (measures)	0.01		0.00		0.00		0.00	
Step 2: <i>X</i> → <i>M</i>								
Fixed effects								
Intercept	−0.14 (0.17)	.42	−0.01 (0.12)	.97	−0.04 (0.12)	.76	0.15 (0.07)	.04*
Effect-size variance	0.49 (0.33)	.15	1.22 (0.62)	.1	0.17 (0.49)	.74	−0.41 (0.27)	.18
Time ^a	0 (0)	.36	−0.01 (0.03)	.67	0 (0)	.67	0 (0)	.63
Number of recommendations	0.12 (0.05)	.02*	0.08 (0.04)	.09	0.07 (0.03)	.05	0.03 (0.02)	.09
Random effects								
σ_1^2 (studies)	0.01		0.00		0.02		0.03	
σ_2^2 (samples)	0.23		0.15		0.03		0.01	
σ_3^2 (measures)	0.00		0.00		0.00		0.00	
Step 3: <i>X</i> + <i>M</i> → <i>Y</i>								
Fixed effects								
Intercept	0.05 (0.07)	.44	0.02 (0.19)	.92	0.03 (0.11)	.77	−0.03 (0.1)	.76
Effect-size variance	0.22 (0.23)	.35	0.65 (0.76)	.43	0.28 (0.41)	.52	0.64 (0.39)	.16
Time ^a	0.01 (0.05)	.8	−0.11 (0.06)	.28	−0.05 (0.02)	.13	−0.04 (0.02)	.24
Number of recommendations	0.02 (0.01)	.12	0.04 (0.05)	.41	0.01 (0)	.19	0.04 (0.01)	.01*
Mediator	0.35 (0.18)	.11	0.06 (0.19)	.79	0.47 (0.13)	.01**	0.28 (0.15)	.1
Random effects								
σ_1^2 (studies)	0.01		0.11		0.09		0.07	
σ_2^2 (samples)	0.02		0.06		0.00		0.01	
σ_3^2 (measures)	0.01		0.00		0.00		0.00	
Indirect effect								
Mediation strength	0.042		.005		.033		.008	

Note. *n* is the number of studies; *k* is the number of unique groups; *s* is the total number of effect sizes; *X* is the independent variable; *M* is the mediator variable; *Y* is the outcome variable; Time = time interval in days between baseline and outcome measure. σ_1^2 is the random component (variance) at the study level; σ_2^2 is the random component (variance) at the sample level; σ_3^2 is the random component (variance) at the measure level. Passive controls in the trials correspond to the number of recommendation being 0's and active controls 1's. *SE* = standard error.

^a Standardized variable.

* *p* < .05. ** *p* < .01.

a sensitivity analysis (see [Supplemental Material K](#)), other analyses may be attempted in the future. For example, although we examined whether a study used randomization, we did not capture whether that randomization used rigorous procedures (e.g., random sequence generation and allocation sequence concealment; [Dechartres et al., 2016](#)), nor did we measure other factors related to potential bias such as double blinding ([Page et al., 2016](#)) or trial preregistration ([Odutayo et al., 2017](#)). Thus, we recommend researchers to additionally consider these important research-quality factors in future syntheses of this literature.

The General and Behavior-Specific Processes of Change

The present meta-analysis was structured around testing a framework of general processes, which do not require a substantial amount of cognitive capacity or implementation motivation, and behavior-specific processes, which do require more cognitive

resources and motivation to take effect. Specifically, we proposed that, in the case of multiple-behavior interventions, general processes should lead to a positive effect of higher numbers of recommendations because the effects of these recommendations occur efficiently. In contrast, behavior-specific processes should be associated with a curvilinear relation between the number of behavioral recommendations and behavioral and clinical change, as the increasing demand on cognitive resources should lead to diminishing returns at high numbers of recommendations.

Accordingly, we found linear associations between the number of recommendations and behavioral and clinical improvements across all tests ([Tables 3 and 4](#)). That is, there was no evidence of weakening or reversals in the positive effects of the number of recommendations when a study recommended a higher number of behaviors within the range of the present investigation. Yet, even though the number of behavioral recommendations ranged from 0 to 17, 87% of the conditions had four or fewer recommendations.

Table 8*Test of Indirect Effects Through Psychological Well-Being and Motivation*

Test of indirect effect	Psychological well-being (<i>s</i> = 139)		Motivation (<i>s</i> = 58)	
	Z/B (SE)	<i>p</i>	Z/B (SE)	<i>p</i>
Sobel test	3.35	.001***	1.10	.27
Path analysis w/bootstrap CI	0.05 (0.01)	.001***	0.009 (0.008)	.26
Path analysis w/robust SE	0.05 (0.01)	.001***	0.009 (0.006)	.18

Note. *s* is the number of effect-size observations. CI = confidence interval; SE = standard error; Z = Z value.

****p* < .001.

This overall linear association thus supports the notion that general processes of behavioral change explain the impact of the number of recommendations on change in behavioral and clinical outcomes.

This linear relation also challenges the notion that compensatory effects between health behaviors would reduce the efficacy of multiple-behavior interventions as the number of recommendations increases. Although prior studies have noted compensatory effects between behaviors such as alcohol use and physical activity (C. R. Nigg et al., 2009) and eating behaviors and physical activity (Martin et al., 2019; C. Nigg et al., 2021), these studies did not examine compensatory effects in the context of multiple-behavior interventions, instead examining behavior when there was either no intervention or only a single-behavior intervention. Thus, it is possible that compensatory effects are less likely to arise in multiple-behavior interventions. As multiple-behavior interventions include specific recommendations targeting each health behavior, recipients may be less likely to use improvements in one behavior to excuse declines in another behavior.

Another finding supporting general processes of behavioral change concerns the association between changes in one behavior and changes in other behaviors. We found that the number of recommendations exerted an effect on changes in one behavior via changes in other behaviors (Table 6). As outlined in the introduction, this finding is consistent with the hypothesis of behavioral cuing, where change in one behavior cues engagement or disengagement with other behaviors. However, changes in alcohol and drug use were the strongest predictors of change in other behaviors, potentially due to reductions in substance use not only limiting cuing for other risk behaviors but also reducing disinhibition and increasing cognitive capacity to follow health recommendations.

Finally, we examined the mediator of improvement in psychological well-being to explain the effect of the number of behavioral recommendations on behavioral and clinical change. Evidence from multiple tests (Tables 6 and 7) demonstrated that psychological well-being consistently played a positive mediating role, which captures reductions in anxiety/stress, trauma, and depression, as well as improvements in coping, mood, and psychosocial quality of life. This finding suggests that simultaneously recommending multiple behavioral changes may have the power to induce a spillover effect, whereby engaging in healthy behavior in one domain leads to improvements in psychological well-being that then facilitate healthy behavior in other domains, thus reinforcing adherence to behavioral recommendations in the long term. In contrast, of the

three behavior-specific mechanisms that we tested (i.e., information, motivation, and behavioral skills), none mediated the relation between the number of behavioral recommendations and behavioral and clinical change, again suggesting that it is primarily general processes of behavioral change that underly this relation.

Limitations

Several limitations of this meta-analysis are worth mentioning. First, the current research questions are limited by the kind of information searched, reported, and publicly available. Despite the overall data set being both sizeable and comprehensive in reporting primary behavioral and clinical outcomes, we have searched and included studies that target eight domains of health behaviors and only approximately 12.4% of the included studies measured psychological well-being. As such, we chose to test mediation using measures of psychological well-being and behavioral or clinical outcomes taken at the same posttest, because using only instances in which psychological well-being was measured at an earlier posttest than the behavioral or clinical outcome would have drastically reduced the number of available effect sizes from 136 to only 45. Nonetheless, testing mediation in this fashion leaves open the possibility of a reverse causal relation (i.e., changes in behavioral or clinical outcomes causing improved psychological well-being). Furthermore, some of the included analyses are near the lower bounds of the recommended number of observations for appropriate statistical power (Tanner-Smith et al., 2016) or concessions were made in the inclusion of confounds to preserve the number of observations. Yet, given the comprehensive nature of this meta-analysis, our analyses are likely the best initial test of mechanisms in these domains of health behavior, but future costly interventions are required to verify the present findings.

Second, it is possible that the search terms we used to conduct the behavioral domain searches lacked certain keywords used in a subset of articles. Specifically, the search terms we used to identify smoking interventions did not include the word “nicotine,” while the search terms we used to identify alcohol-use and substance-use interventions did not include the names of common substances such as “cocaine,” “heroin,” and “marijuana.” To address this issue, we re-ran our diet, exercise, and smoking literature search in several databases both with and without the search term “nicotine” to test whether adding this word to our original list of search terms would significantly increase the number of resulting studies. We found that, compared to using our original search terms, adding “nicotine” increased the number of search results from 1,407 to 1,414 (0.5% increase) in APA PsycInfo; 811–814 (0.4% increase) in Scopus; and 17,641–17,689 (0.3% increase) in Web of Science. Similarly, we re-ran our HIV, alcohol use, and drug use literature search in several databases both with and without adding the words “cocaine,” “heroin,” and “marijuana” to our original search terms. We found that, compared to using our original search terms, adding “cocaine,” “heroin,” and “marijuana” increased the number of search results from 5,102 to 5,490 (7.6% increase) in APA PsycInfo; 1,871–1,895 (1.3% increase) in Scopus; and 39,857–41,509 (4.1% increase) in Web of Science. Thus, although it is possible that a few relevant articles were missed during our original search, based on the above evidence, our original search terms captured the vast majority of eligible studies.

Third, the analyses targeted to investigate mechanisms of behavioral change would be optimal if well-being were fully reported in all studies. Recommended methods for tests of mediation in meta-analysis are either to calculate a joint indirect effect using meta-analytic structural equation modeling or to calculate a cross-lagged effect size that accounts for the covariance of behaviors (Cheung, 2015). Unfortunately, both approaches require complete information to calculate a variance-covariance matrix (e.g., the covariance between two different outcomes reported in each study; Cheung, 2015). Across the wide variety of studies included, the rate of correlation-matrix or covariance reporting was so low that neither of these approaches was feasible. The regression-based and path analysis models (Table 8) presented here provide less precise estimates of indirect effects over time. A better but yet unavailable approach would be a model that helps directly test the effect of the number of recommendations on outcomes at the delayed posttest through changes in the mediators at the immediate posttest. We thus encourage researchers to report correlation matrices for all outcomes whenever possible and make de-identified data from their studies available in a more routine fashion.

Fourth, an inherent problem of all meta-analyses is that these analyses reflect effects aggregated to the sample level. Behavioral change is inherently a process that occurs within an individual, and therefore, behavioral change may not be well represented in aggregated data (e.g., Simpson's Paradox; Simpson, 1951). The contribution of this meta-analysis, however, is to assess whether a common mechanism occurs with sufficient frequency in each intervention population to produce the expected average effects. Regardless, these results should be cautiously replicated with measures of behavioral and clinical change among individual participants.

Fifth, while our analyses suggest that the linear effect of the number of recommendations is robust to whether interventions were conducted in Western or non-Western countries and that studies in non-Western nations were associated with larger effect sizes, it should be noted that our review was comprised primarily of samples taken in Western contexts, with under 20% of the 364 included studies having been conducted in non-Western countries (see Supplemental Material G). Furthermore, our use of English keywords and reliance on databases that cater primarily to English-language content likely precluded us from finding additional relevant non-Western studies. Thus, our findings regarding Western versus non-Western countries should be viewed as preliminary pending replication with additional non-Western studies, and we recommend that researchers conducting future meta-analyses of this literature expand their searches to include non-English-language keywords in Google Scholar to capture a more comprehensive and diverse set of reports in this domain.

Finally, the primary analysis in our work relied on within-group growth effect sizes. Although this approach has several benefits for answering our research questions, the lack of control through group comparison within studies is a limitation. We therefore developed the propensity-score model to help make the groups receiving different numbers of behavioral recommendations more comparable at baseline and included the propensity scores as a covariate in the CHE+ models. Future meta-analytic research can explore means of utilizing between-group comparisons within studies to better understand the mechanisms associated with the impact of the number of behavioral recommendations. The network meta-analysis approach represents such a possibility (Cipriani et al., 2013).

Future Directions

One question in this meta-analysis is whether there is evidence of decreases in recommendation benefits as they approach a high number, as predicted if behavior-specific processes are at work and the number of recommendations is sufficiently taxing. Although little evidence of a curvilinear association between the number of recommendations and behavioral and clinical change using typical criteria has been identified, a substantial range restriction in the data set exists, with the majority of studies presenting only zero to four recommendations and most studies topping out at five recommendations. This range of recommendations may not overtax mental resources (Cowan, 2010), or changes in several behaviors may operate efficiently as proposed. A much-needed line of research is multiple experimental tests of whether providing five or more recommendations would still produce consistent linear benefits.

Another important research direction is to elucidate differences between sequential and concurrent behavioral change and how each is associated with the mechanisms proposed here. A meta-analysis is unable to distinguish these two approaches to behavioral change given the overall large scale of time between measurements in comparison to the underlying behavioral process (i.e., 8–12 weeks), and sequential versus concurrent change is a relatively understudied area of multiple-behavior-change research that has produced mixed results. Individual interventions have found that sequential change may be beneficial, such as a weight loss intervention that supported a sequential approach to diet and smoking cessation (Spring et al., 2004). In contrast, another that directly compared the two approaches found a small advantage for concurrent over sequential change in achieving long-term behavioral change (20.3% vs. 16.9%; Hyman et al., 2007). A more recent systematic review found only six multiple-behavior interventions that directly compared sequential versus concurrent behavioral change, and there was no evidence strongly supporting one approach over the other when synthesizing the available evidence (James et al., 2016). The question of which approach is better and how variations in this approach relate to the number and content of behavioral recommendations is still open with multiple implications. An additional question is whether both approaches operate through the same or different psychosocial processes. Both questions invite future work, such as experimentally inducing simultaneous/sequential change and naturalistically examining each approach to behavioral change.

The current project focused on investigating two general mechanisms (i.e., behavioral cuing vs. mediation via psychological well-being) underpinning behavioral change in research on multiple behavioral recommendations. One addition to the framework worth exploring is the role of personality change in response to interventions (Roberts et al., 2017). A person acquiring and applying behavioral skills to multiple behaviors may exhibit general changes in self-control, and examining whether or not this occurs requires further analyses to see if changes generalize to behaviors that were not addressed in the intervention. The current measure of psychological well-being encompasses items like anxiety and negative affect that are likely associated with the personality measure of neuroticism (Cattell & Scheier, 1961). Furthermore, people may respond to multiple recommendations differently depending on their personality, with more conscientious people organizing and pursuing multiple behaviors more effectively. As such, future experimental work

examining individual differences in multiple-behavior interventions would be an interesting and needed line of research.

Conclusions

To conclude, providing multiple behavioral recommendations is positively associated with changes in behavioral and clinical outcomes. Leveraging the CHE+ meta-regression model and a comprehensive data set, the present meta-analysis provided strong evidence that more recommendations are linearly associated with more behavioral change when the number of recommendations is within the range of zero–four behavioral recommendations, which are most frequently investigated. It further identified evidence of general mechanisms, including behavioral cuing among outcome behaviors and changes in psychological well-being, to explain the advantage of multiple behavioral recommendations. Hopefully, this research will inspire further endeavors to help people achieve healthier lives and inspire practitioners to recommend multiple behaviors in critical areas of human health.

References

- References marked with an asterisk have studies included in the present meta-analysis.
- *Abrantes, A. M., Bloom, E. L., Strong, D. R., Riebe, D., Marcus, B. H., Desaulniers, J., Fokas, K., & Brown, R. A. (2014). A preliminary randomized controlled trial of a behavioral exercise intervention for smoking cessation. *Nicotine & Tobacco Research*, 16(8), 1094–1103. <https://doi.org/10.1093/ntr/ntu036>
 - *Adam, L. M., Jarman, M., Barker, M., Manca, D. P., Lawrence, W., & Bell, R. C. (2020). Use of healthy conversation skills to promote healthy diets, physical activity and gestational weight gain: Results from a pilot randomised controlled trial. *Patient Education and Counseling*, 103(6), 1134–1142. <https://doi.org/10.1016/j.pec.2020.01.001>
 - Addington, D., Addington, J., & Maticka-Tyndale, E. (1993). Assessing depression in schizophrenia: The Calgary Depression Scale. *British Journal of Psychiatry*, 163(S22), 39–44. <https://doi.org/10.1192/S0007125000292581>
 - *Ahmadi, M., Laumeier, I., Ihl, T., Steinicke, M., Ferse, C., Endres, M., Grau, A., Hastrup, S., Poppert, H., Palm, F., Schoene, M., Seifert, C. L., Kandil, F. I., Weber, J. E., von Weitzel-Mudersbach, P., Wimmer, M. L. J., Algra, A., Amarenco, P., Greving, J. P., ... Audebert, H. J. (2020). A support programme for secondary prevention in patients with transient ischaemic attack and minor stroke (INSPIRE-TMS): An open-label, randomised controlled trial. *Lancet Neurology*, 19(1), 49–60. [https://doi.org/10.1016/S1474-4422\(19\)30369-2](https://doi.org/10.1016/S1474-4422(19)30369-2)
 - Ajzen, I. (1991). The theory of planned behavior. *Organizational Behavior and Human Decision Processes*, 50(2), 179–211. [https://doi.org/10.1016/0749-5978\(91\)90020-T](https://doi.org/10.1016/0749-5978(91)90020-T)
 - Albarracín, D. (2002). Cognition in persuasion: An analysis of information processing in response to persuasive communications. *Advances in Experimental Social Psychology*, 34, 61–130. [https://doi.org/10.1016/S0065-2601\(02\)80004-1](https://doi.org/10.1016/S0065-2601(02)80004-1)
 - Albarracín, D. (2021). *Action and inaction in a social world: Predicting and changing attitudes and behavior*. Cambridge University Press. <https://doi.org/10.1017/9781108878357>
 - Albarracín, D., Gillette, J. C., Earl, A. N., Glasman, L. R., Durantini, M. R., & Ho, M.-H. (2005). A test of major assumptions about behavior change: A comprehensive look at the effects of passive and active HIV-prevention interventions since the beginning of the epidemic. *Psychological Bulletin*, 131(6), 856–897. <https://doi.org/10.1037/0033-2909.131.6.856>
 - Albarracín, D., Johnson, B. T., Fishbein, M., & Muellerleile, P. A. (2001). Theories of reasoned action and planned behavior as models of condom use: A meta-analysis. *Psychological Bulletin*, 127(1), 142–161. <https://doi.org/10.1037/0033-2909.127.1.142>
 - *Albrecht, A. E., Marcus, B. H., Roberts, M., Forman, D. E., & Parisi, A. F. (1998). Effect of smoking cessation on exercise performance in female smokers participating in exercise training. *The American Journal of Cardiology*, 82(8), 950–955. [https://doi.org/10.1016/S0002-9149\(98\)00511-6](https://doi.org/10.1016/S0002-9149(98)00511-6)
 - *Alghamdi, R. Q. (2017). A randomized controlled trial of a 12-week intensive lifestyle intervention program at a primary care obesity clinic for adults in western Saudi Arabia. *Saudi Medical Journal*, 38(8), 837–845. <https://doi.org/10.15537/smj.2017.8.20553>
 - *Allen, J. K., Blumenthal, R. S., Margolis, S., Young, D. R., Miller, E. R., III, & Kelly, K. (2002). Nurse case management of hypercholesterolemia in patients with coronary heart disease: Results of a randomized clinical trial. *American Heart Journal*, 144(4), 678–686. [https://doi.org/10.1016/S0002-8703\(02\)00141-2](https://doi.org/10.1016/S0002-8703(02)00141-2)
 - *Amaro, H., Larson, M., Zhang, A., Acevedo, A., Dai, J., & Matsumoto, A. (2007). Effects of trauma intervention on HIV sexual risk behaviors among women with co-occurring disorders in substance abuse treatment. *Journal of Community Psychology*, 35(7), 895–908. <https://doi.org/10.1002/jcop.20188>
 - *Amoah, J., Said, S., Rampal, L., Manaf, R., Ibrahim, N., Owusu-Agyei, S., & Poku Asante, K. (2021). Effects of a school-based intervention to reduce cardiovascular disease risk factors among secondary school students: A cluster-randomized, controlled trial. *PLOS ONE*, 16(11), Article e0259581. <https://doi.org/10.1371/journal.pone.0259581>
 - *Anderson, A. S., Chong, H. Y., Craigie, A. M., Donnan, P. T., Gallant, S., Hickman, A., McAdam, C., McKell, J., McNamee, P., Macaskill, E. J., Mutrie, N., O'Carroll, R. E., Rauchhaus, P., Sattar, N., Stead, M., & Treweek, S. (2021). A novel approach to increasing community capacity for weight management a volunteer-delivered programme (ActWELL) initiated within breast screening clinics: A randomised controlled trial. *The International Journal of Behavioral Nutrition and Physical Activity*, 18(1), Article 34. <https://doi.org/10.1186/s12966-021-01099-7>
 - *Anderson, A. S., Dunlop, J., Gallant, S., Macleod, M., Miedzybrodzka, Z., Mutrie, N., O'Carroll, R. E., Stead, M., Steele, R. J. C., Taylor, R. S., Vinnicombe, S., & Berg, J. (2018). Feasibility study to assess the impact of a lifestyle intervention ('LivingWELL') in people having an assessment of their family history of colorectal or breast cancer. *BMJ Open*, 8(2), Article e019410. <https://doi.org/10.1136/bmjopen-2017-019410>
 - *Anderson, D., Mizzari, K., Kain, V., & Webster, J. (2006). The effects of a multimodal intervention trial to promote lifestyle factors associated with the prevention of cardiovascular disease in menopausal and postmenopausal Australian women. *Health Care for Women International*, 27(3), 238–253. <https://doi.org/10.1080/07399330500506543>
 - *Andersson, C. (2020). Personalized normative feedback interventions targeting hazardous alcohol use and alcohol-related risky sexual behavior in Swedish university students: A randomized controlled replication trial. *Addictive Behaviors Reports*, 12, Article 100300. <https://doi.org/10.1016/j.abrep.2020.100300>
 - *Andreae, S. J., Andreae, L. J., Cherrington, A. L., Richman, J. S., Johnson, E., Clark, D., & Safford, M. M. (2021). Peer coach delivered storytelling program improved diabetes medication adherence: A cluster randomized trial. *Contemporary Clinical Trials*, 104, Article 106358. <https://doi.org/10.1016/j.cct.2021.106358>
 - Angrave, D., & Charlwood, A. (2015). What is the relationship between long working hours, over-employment, under-employment and the subjective well-being of workers? Longitudinal evidence from the UK. *Human Relations*, 68(9), 1491–1515. <https://doi.org/10.1177/0018726714559752>
 - *Anstey, K. J., Cherbuin, N., Kim, S., McMaster, M., D'Este, C., Lautenschlager, N., Rebok, G., McRae, I., Torres, S. J., Cox, K. L., & Pond, C. D. (2020). An internet-based intervention augmented with a diet

- and physical activity consultation to decrease the risk of dementia in at-risk adults in a primary care setting: Pragmatic randomized controlled trial. *Journal of Medical Internet Research*, 22(9), Article e19431. <https://doi.org/10.2196/19431>
- Appelhans, B. M., Whited, M. C., Schneider, K. L., Ma, Y., Oleski, J. L., Merriam, P. A., Waring, M. E., Olendzki, B. C., Mann, D. M., Ockene, I. S., & Pagoto, S. L. (2012). Depression severity, diet quality, and physical activity in women with obesity and depression. *Journal of the Academy of Nutrition and Dietetics*, 112(5), 693–698. <https://doi.org/10.1016/j.jand.2012.02.006>
- *Arad, A. D., DiMenna, F. J., Thomas, N., Tamis-Holland, J., Weil, R., Geliebter, A., & Albu, J. B. (2015). High-intensity interval training without weight loss improves exercise but not basal or insulin-induced metabolism in overweight/obese African American women. *Journal of Applied Physiology*, 119(4), 352–362. <https://doi.org/10.1152/jappphysiol.100306.2015>
- *Arent, S. M., Walker, A. J., Pellegrino, J. K., Sanders, D. J., McFadden, B. A., Ziegenfuss, T. N., & Lopez, H. L. (2018). The combined effects of exercise, diet, and a multi-ingredient dietary supplement on body composition and adipokine changes in overweight adults. *Journal of the American College of Nutrition*, 37(2), 111–120. <https://doi.org/10.1080/07315724.2017.1368039>
- Ashdown-Franks, G., Firth, J., Carney, R., Carvalho, A. F., Hallgren, M., Koyanagi, A., Rosenbaum, S., Schuch, F. B., Smith, L., Solmi, M., Vancampfort, D., & Stubbs, B. (2020). Exercise as medicine for mental and substance use disorders: A meta-review of the benefits for neuropsychiatric and cognitive outcomes. *Sports Medicine*, 50(1), 151–170. <https://doi.org/10.1007/s40279-019-01187-6>
- *Atlantis, E., Chow, C. M., Kirby, A., & Fiatarone Singh, M. A. (2006). Worksite intervention effects on physical health: A randomized controlled trial. *Health Promotion International*, 21(3), 191–200. <https://doi.org/10.1093/heapro/dal012>
- Austin, P. C. (2011). An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivariate Behavioral Research*, 46(3), 399–424. <https://doi.org/10.1080/00273171.2011.568786>
- *Avants, S., Arthur Margolin, A., Usubiaga, M., & Doebrick, C. (2004). Targeting HIV-related outcomes with intravenous drug users maintained on methadone: A randomized clinical trial of a harm reduction group therapy. *Journal of Substance Abuse Treatment*, 26(2), 67–78. [https://doi.org/10.1016/S0740-5472\(03\)00159-4](https://doi.org/10.1016/S0740-5472(03)00159-4)
- *Bachanas, P., Kidder, D., Medley, A., Pals, S. L., Carpenter, D., Howard, A., Antelman, G., DeLuca, N., Muhenje, O., Sheriff, M., Somi, G., Katuta, F., Cherutich, P., & Moore, J. (2016). Delivering prevention interventions to people living with HIV in clinical care settings: Results of a cluster randomized trial in Kenya, Namibia, and Tanzania. *AIDS and Behavior*, 20(9), 2110–2118. <https://doi.org/10.1007/s10461-016-1349-2>
- *Bae, J. W., Woo, S. I., Lee, J., Park, S. D., Kwon, S. W., Choi, S. H., Yoon, G. S., Kim, M. S., Hwang, S. S., & Lee, W. K. (2021). mHealth interventions for lifestyle and risk factor modification in coronary heart disease: Randomized controlled trial. *JMIR mHealth and uHealth*, 9(9), Article e29928. <https://doi.org/10.2196/29928>
- *Bahromov, M., & Weine, S. (2011). HIV prevention for migrants in transit: Developing and testing TRAIN. *AIDS Education and Prevention*, 23(3), 267–280. <https://doi.org/10.1521/aeap.2011.23.3.267>
- Bakker, A. B., Sanz Vergel, A. I., & Kuntze, J. (2015). Student engagement and performance: A weekly diary study on the role of openness. *Motivation and Emotion*, 39(1), 49–62. <https://doi.org/10.1007/s1031-014-9422-5>
- *Bandini, L. G., Eliasziw, M., Ditttrich, G. A., Curtin, C., Maslin, M., Must, A., Boutelle, K. N., & Fleming, R. K. (2021). A family-based weight loss randomized controlled trial for youth with intellectual disabilities. *Pediatric Obesity*, 16(11), Article e12816. <https://doi.org/10.1111/ijpo.12816>
- Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change. *Psychological Review*, 84(2), 191–215. <https://doi.org/10.1037/0033-295X.84.2.191>
- *Baranowski, T., Baranowski, J., Cullen, K., Thompson, D. I., Nicklas, T., Zakeri, I. E., & Rochon, J. (2003). The fun, food, and fitness project (FFFP): The Baylor GEMS Pilot Study. *Ethnicity and Disease*, 13(Suppl. 1), S30–S39.
- Bargh, J. A. (1994). The four horsemen of automaticity: Awareness, intention, efficiency, and control in social cognition. In R. S. Wyer, Jr., & T. K. Srull (Eds.), *Handbook of social cognition: Basic processes; Applications* (pp. 1–40). Lawrence Erlbaum Associates.
- Bargh, J. A., Chaiken, S., Govender, R., & Pratto, F. (1992). The generality of the automatic attitude activation effect. *Journal of Personality and Social Psychology*, 62(6), 893–912. <https://doi.org/10.1037/0022-3514.62.6.893>
- *Barnes, C., Hall, A., Nathan, N., Sutherland, R., McCarthy, N., Pettet, M., Brown, A., & Wolfenden, L. (2021). Efficacy of a school-based physical activity and nutrition intervention on child weight status: Findings from a cluster randomized controlled trial. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 153, Article 106822. <https://doi.org/10.1016/j.ypmed.2021.106822>
- *Barroso, M., Zomeño, M. D., Díaz, J. L., Pérez-Fernández, S., Martí-Lluch, R., Cerdón, F., Ramos, R., Cabezas, C., Salvador, G., Castell, C., Schröder, H., & Grau, M. (2020). Control of cardiovascular risk factors with tailored recommendations: A randomized controlled trial. *Preventive Medicine*, 141, Article 106302. <https://doi.org/10.1016/j.ypmed.2020.106302>
- *Baviskar, M. P., Rangari, S., Mishra, S., & Mohanta, B. S. (2021). Assessment of a group-based comprehensive diabetes management program to improve glycemic control, quality of life and self-care behavior in patients with type 2 diabetes mellitus in a primary healthcare setting of a metropolitan city in India: CDMP MUM Trial. *International Journal of Diabetes in Developing Countries*, 41(1), 156–163. <https://doi.org/10.1007/s13410-020-00828-1>
- *Bayne-Smith, M., Fardy, P. S., Azzollini, A., Magel, J., Schmitz, K. H., & Agin, D. (2004). Improvements in heart health behaviors and reduction in coronary artery disease risk factors in urban teenaged girls through a school-based intervention: The PATH program. *American Journal of Public Health*, 94(9), 1538–1543. <https://doi.org/10.2105/AJPH.94.9.1538>
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, 4(6), 561–571. <https://doi.org/10.1001/archpsyc.1961.01710120031004>
- *Becker, D. M., Yanek, L. R., Johnson, W. R., Jr., Garrett, D., Moy, T. F., Reynolds, S. S., Blumenthal, R. S., Vaidya, D., & Becker, L. C. (2005). Impact of a community-based multiple risk factor intervention on cardiovascular risk in black families with a history of premature coronary disease. *Circulation*, 111(10), 1298–1304. <https://doi.org/10.1161/01.CIR.0000157734.97351.B2>
- *Befort, C. A., VanWormer, J. J., Desouza, C., Ellerbeck, E. F., Gajewski, B., Kimminau, K. S., Greiner, K. A., Perri, M. G., Brown, A. R., Pathak, R. D., Huang, T. T., Eiland, L., & Drincic, A. (2021). Effect of behavioral therapy with in-clinic or telephone group visits vs in-clinic individual visits on weight loss among patients with obesity in rural clinical practice: A randomized clinical trial. *JAMA: Journal of the American Medical Association*, 325(4), 363–372. <https://doi.org/10.1001/jama.2020.25855>
- *Berendsen, B. A. J., Hendriks, M. R. C., Rutten, G. M., Kremers, S. P. J., Savelberg, H. H. C. M., & Schaper, N. C. (2020). The added value of frequent physical activity group sessions in a combined lifestyle intervention: A cluster randomised trial in primary care. *Preventive Medicine Reports*, 20, Article 101204. <https://doi.org/10.1016/j.pmedr.2020.101204>
- Bergstra, S. A., Sepriano, A., Ramiro, S., & Landewé, R. (2019). Three handy tips and a practical guide to improve your propensity score models. *RMD Open*, 5(1), Article e000953. <https://doi.org/10.1136/rmdopen-2019-000953>
- *Bijlholt, M., Ameye, L., Van Uytsel, H., Devlieger, R., & Bogaerts, A. (2021). The INTER-ACT E-health supported lifestyle intervention

- improves postpartum food intake and eating behavior, but not physical activity and sedentary behavior—A randomized controlled trial. *Nutrients*, 13(4), Article 1287. <https://doi.org/10.3390/nu13041287>
- *Bilger, M., Shah, M., Tan, N. C., Tan, C. Y. L., Bundoc, F. G., Bairavi, J., & Finkelstein, E. A. (2021). Process- and outcome-based financial incentives to improve self-management and glycemic control in people with type 2 diabetes in Singapore: A randomized controlled trial. *The Patient: Patient-Centered Outcomes Research*, 14(5), 555–567. <https://doi.org/10.1007/s40271-020-00491-y>
- Bolnick, H. J., Bui, A. L., Bulchis, A., Chen, C., Chapin, A., Lomsadze, L., Mokdad, A. H., Millard, F., & Dieleman, J. L. (2020). Health-care spending attributable to modifiable risk factors in the USA: An economic attribution analysis. *The Lancet. Public Health*, 5(10), e525–e535. [https://doi.org/10.1016/S2468-2667\(20\)30203-6](https://doi.org/10.1016/S2468-2667(20)30203-6)
- Bond, F. W., Hayes, S. C., Baer, R. A., Carpenter, K. M., Guenole, N., Orcutt, H. K., Waltz, T., & Zettle, R. D. (2011). Preliminary psychometric properties of the Acceptance and Action Questionnaire-II: A revised measure of psychological inflexibility and experiential avoidance. *Behavior Therapy*, 42(4), 676–688. <https://doi.org/10.1016/j.beth.2011.03.007>
- Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009). *Introduction to meta-analysis*. Wiley. <https://doi.org/10.1002/9780470743386>
- Borenstein, M., Higgins, J. P. T., Hedges, L. V., & Rothstein, H. R. (2017). Basics of meta-analysis: I^2 is not an absolute measure of heterogeneity. *Research Synthesis Methods*, 8(1), 5–18. <https://doi.org/10.1002/jrsm.1230>
- *Brawner, B. M., Jemmott, L. S., Wingood, G., Lozano, A. J., & Hanlon, A. L. (2019). Project GOLD: A pilot randomized controlled trial of a novel psychoeducational HIV/STI prevention intervention for heterosexually-active black youth. *Research in Nursing & Health*, 42(1), 8–28. <https://doi.org/10.1002/nur.21930>
- *Broekhuizen, K., van Poppel, M. N. M., Koppes, L. L., Kindt, I., Brug, J., & van Mechelen, W. (2012). Can multiple lifestyle behaviours be improved in people with familial hypercholesterolemia? Results of a parallel randomised controlled trial. *PLOS ONE*, 7(12), Article e50032. <https://doi.org/10.1371/journal.pone.0050032>
- Brookmeyer, K. A., Hogben, M., & Kinsey, J. (2016). The role of behavioral counseling in sexually transmitted disease prevention program settings. *Sexually Transmitted Diseases*, 43(2), S102–S112. <https://doi.org/10.1097/OLQ.0000000000000327>
- *Brown, J. C., Sarwer, D. B., Troxel, A. B., Sturgeon, K., DeMichele, A. M., Denlinger, C. S., & Schmitz, K. H. (2021). A randomized trial of exercise and diet on body composition in survivors of breast cancer with overweight or obesity. *Breast Cancer Research and Treatment*, 189(1), 145–154. <https://doi.org/10.1007/s10549-021-06284-7>
- *Brown, L. K., Chernoff, M., Kennard, B. D., Emslie, G. J., Lypen, K., Buisson, S., Weinberg, A., Whiteley, L. B., Traite, S., Krotje, C., Harrieff, L., Townley, E., Bunch, A., Purswani, M., Shaw, R., Spector, S. A., Agwu, A., Shapiro, D. E., & the IMPAACT 2002 team. (2021). Site-randomized controlled trial of a combined cognitive behavioral therapy and a medication management algorithm for treatment of depression among youth living with HIV in the United States. *Journal of Acquired Immune Deficiency Syndromes*, 88(5), 497–505. <https://doi.org/10.1097/QAI.00000000000002790>
- *Bruckert, E., Giral, P., Paillard, F., Ferrières, J., Schlienger, J. L., Renucci, J. F., Abdennbi, K., Durack, I., Chadarevian, R., & the PEGASE group. (2008). Effect of an educational program (PEGASE) on cardiovascular risk in hypercholesterolaemic patients. *Cardiovascular Drugs and Therapy*, 22(6), 495–505. <https://doi.org/10.1007/s10557-008-6137-4>
- Bryan, A. D., Magnan, R. E., Gillman, A. S., Yeater, E. A., Feldstein Ewing, S. W., Kong, A. S., & Schmiede, S. J. (2018). Effect of including alcohol and cannabis content in a sexual risk-reduction intervention on the incidence of sexually transmitted infections in adolescents: A cluster randomized clinical trial. *JAMA Pediatrics*, 172(4), Article e175621. <https://doi.org/10.1001/jamapediatrics.2017.5621>
- *Burke, H. M., Chen, M., Murray, K., Bezuidenhout, C., Ngwepe, P., Bernholc, A., & Medina-Marino, A. (2020). The effects of the integration of an economic strengthening and HIV prevention education programme on the prevalence of sexually transmitted infections and savings behaviours among adolescents: A full-factorial randomised controlled trial in South Africa. *BMJ Global Health*, 5(4), Article e002029. <https://doi.org/10.1136/bmjgh-2019-002029>
- Burke, L. E., Wang, J., & Sevvick, M. A. (2011). Self-monitoring in weight loss: A systematic review of the literature. *Journal of the American Dietetic Association*, 111(1), 92–102. <https://doi.org/10.1016/j.jada.2010.10.008>
- *Burke, V., Beilin, L. J., Cutt, H. E., Mansour, J., Wilson, A., & Mori, T. A. (2005). Effects of a lifestyle programme on ambulatory blood pressure and drug dosage in treated hypertensive patients: A randomized controlled trial. *Journal of Hypertension*, 23(6), 1241–1249. <https://doi.org/10.1097/01.hjh.0000170388.61579.4f>
- *Burke, V., Giangulio, N., Gillam, H. F., Beilin, L. J., Houghton, S., & Milligan, R. A. K. (1999). Health promotion in couples adapting to a shared lifestyle. *Health Education Research*, 14(2), 269–288. <https://doi.org/10.1093/her/14.2.269>
- *Bush, T., Lovejoy, J., Javitz, H., Torres, A. J., Wassum, K., Tan, M. M., & Spring, B. (2018). Simultaneous vs. sequential treatment for smoking and weight management in tobacco quitlines: 6 and 12 month outcomes from a randomized trial. *BMC Public Health*, 18(1), Article 678. <https://doi.org/10.1186/s12889-018-5574-7>
- *Butryn, M. L., Martinelli, M. K., Crane, N. T., Godfrey, K., Roberts, S. R., Zhang, F., & Forman, E. M. (2020). Counselor surveillance of digital self-monitoring data: A pilot randomized controlled trial. *Obesity*, 28(12), 2339–2346. <https://doi.org/10.1002/oby.23015>
- *Byrnes, H. F., Miller, B. A., Bourdeau, B., Johnson, M. B., Buller, D. B., Berteletti, J., & Rogers, V. A. (2019). Prevention of alcohol and other drug overuse among nightclub patrons: A randomized trial of a group-based mobile intervention at nightclubs. *Journal of Studies on Alcohol and Drugs*, 80(4), 423–430. <https://doi.org/10.15288/jsad.2019.80.423>
- *Cahill, A. G., Haire-Joshu, D., Cade, W. T., Stein, R. I., Woolfolk, C. L., Moley, K., Mathur, A., Schechtman, K., & Klein, S. (2018). Weight control program and gestational weight gain in disadvantaged women with overweight or obesity: A randomized clinical trial. *Obesity*, 26(3), 485–491. <https://doi.org/10.1002/oby.22070>
- *Cakir, H., & Pinar, R. (2006). Randomized controlled trial on lifestyle modification in hypertensive patients. *Western Journal of Nursing Research*, 28(2), 190–209. <https://doi.org/10.1177/0193945905283367>
- *Calsyn, D., Hatch-Maillette, M., Tross, S., Doyle, S., Crits-Christoph, P., Song, Y., Harrer, J., Lalos, G., & Berns, S. (2009). Motivational and behavioral skills training HIV/STI sexual risk reduction groups for men. *Journal of Substance Abuse Treatment*, 37(2), 138–150. <https://doi.org/10.1016/j.jsat.2008.11.008>
- *Cambien, F., Richard, J. L., Ducimetiere, P., Warnet, J. M., & Kahn, J. (1981). The Paris cardiovascular risk factor prevention trial. Effects of two years of intervention in a population of young men. *Journal of Epidemiology and Community Health*, 35(2), 91–97. <https://doi.org/10.1136/jech.35.2.91>
- *Camolas, J., Santos, O., Moreira, P., & do Carmo, I. (2017). INDIVIDUO: Results from a patient-centered lifestyle intervention for obesity surgery candidates. *Obesity Research & Clinical Practice*, 11(4), 475–488. <https://doi.org/10.1016/j.orcp.2016.08.003>
- *Campbell, M. K., Tessaro, I., DeVellis, B., Benedict, S., Kelsey, K., Belton, L., & Sanhueza, A. (2002). Effects of a tailored health promotion program for female blue-collar workers: Health works for women. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 34(3), 313–323. <https://doi.org/10.1006/pmed.2001.0988>
- Carels, R. A., Burmeister, J. M., Koball, A. M., Oehlhof, M. W., Hinman, N., LeRoy, M., Bannon, E., Ashrafioun, L., Storfer-Isser, A., Darby, L. A., & Gumble, A. (2014). A randomized trial comparing two approaches to

- weight loss: Differences in weight loss maintenance. *Journal of Health Psychology*, 19(2), 296–311. <https://doi.org/10.1177/1359105312470156>
- *Carey, M. P., Rich, C., Norris, A. L., Krieger, N., Gavarkovs, A. G., Kaplan, C., Guthrie, K. M., & Carey, K. B. (2020). A brief clinic-based intervention to reduce alcohol misuse and sexual risk behavior in young women: Results from an exploratory clinical trial. *Archives of Sexual Behavior*, 49(4), 1231–1250. <https://doi.org/10.1007/s10508-020-01635-9>
- Carpenter, C. J. (2010). A meta-analysis of the effectiveness of health belief model variables in predicting behavior. *Health Communication*, 25(8), 661–669. <https://doi.org/10.1080/10410236.2010.521906>
- *Carpenter, C. L., Kapur, K., Ramakrishna, P., Pamujula, S., Yadav, K., Giovanni, J. E., Julian, O., Ekstrand, M. L., Sinha, S., & Nyamathi, A. M. (2022). Lean mass improvement from nutrition education and protein supplementation among Rural Indian women living with HIV/AIDS: Results from cluster randomized factorial trial at 18-month follow-up. *Nutrients*, 14(1), Article 179. <https://doi.org/10.3390/nu14010179>
- *Carrico, A. W., Antoni, M. H., Durán, R. E., Ironson, G., Penedo, F., Fletcher, M. A., Klimas, N., & Schneiderman, N. (2006). Reductions in depressed mood and denial coping during cognitive behavioral stress management with HIV-Positive gay men treated with HAART. *Annals of Behavioral Medicine*, 31(2), 155–164. https://doi.org/10.1207/s15324796abm3102_7
- *Castro-Conde, A., Abeytua, M., Arrarte Esteban, V. I., Caravaca Pérez, P., Dalmau González-Gallarza, R., Garza Benito, F., Hidalgo Urbano, R. J., Torres Marqués, J., Vidal-Pérez, R., & Nuñez-Gil, I. J. (2021). Feasibility and results of an intensive cardiac rehabilitation program. Insights from the MxM (Más por Menos) randomized trial. *Revista Española de Cardiología*, 74(6), 518–525. <https://doi.org/10.1016/j.rec.2020.03.029>
- Cattell, R. B., & Scheier, I. H. (1961). *The meaning and measurement of neuroticism and anxiety*. Ronald.
- *Celio, M. A., Mastroleo, N. R., Barnett, N. P., Colby, S. M., Kahler, C. W., Operario, D., & Monti, P. M. (2019). Mechanisms of behavior change in a brief dual-target motivational intervention: Reduction in alcohol use mediates intervention effects on risky sex. *Psychology of Addictive Behaviors*, 33(4), 349–359. <https://doi.org/10.1037/ad0000461>
- *Chaiyasoot, K., Sarasak, R., Pheungruang, B., Dawilai, S., Pramyothin, P., Boonyasiri, A., Supapuang, O., Jassil, F. C., Yamwong, P., & Batterham, R. L. (2018). Evaluation of a 12-week lifestyle education intervention with or without partial meal replacement in Thai adults with obesity and metabolic syndrome: A randomised trial. *Nutrition & Diabetes*, 8(1), Article 23. <https://doi.org/10.1038/s41387-018-0034-0>
- *Cheng, L., Sit, J. W. H., Choi, K. C., Chair, S. Y., Li, X., Wu, Y., Long, J., & Tao, M. (2018). Effectiveness of a patient-centred, empowerment-based intervention programme among patients with poorly controlled type 2 diabetes: A randomised controlled trial. *International Journal of Nursing Studies*, 79, 43–51. <https://doi.org/10.1016/j.ijnurstu.2017.10.021>
- *Chernoff, R. A., & Davison, G. C. (2005). An evaluation of a brief HIV/AIDS prevention intervention for college students using normative feedback and goal setting. *AIDS Education and Prevention*, 17(2), 91–104. <https://doi.org/10.1521/aeap.17.3.91.62902>
- Cheung, M. W. L. (2015). *Meta-analysis: A structural equation modeling approach*. Wiley. <https://doi.org/10.1002/9781118957813>
- Cho, K. W., Altarriba, J., & Popiel, M. (2015). Mental juggling: When does multitasking impair reading comprehension? *Journal of General Psychology*, 142(2), 90–105. <https://doi.org/10.1080/00221309.2014.1003029>
- *Choi, H. K., & Kim, H. O. (2020). Effect of lifestyle intervention program for overweight and obesity pregnant women. *Journal of Korean Academy of Nursing*, 50(3), 459–473. <https://doi.org/10.4040/jkan.19228>
- *Choo, E. K., Zlotnick, C., Strong, D. R., Squires, D. D., Tapé, C., & Mello, M. J. (2016). BSAFER: A web-based intervention for drug use and intimate partner violence demonstrates feasibility and acceptability among women in the emergency department. *Substance Abuse*, 37(3), 441–449. <https://doi.org/10.1080/08897077.2015.1134755>
- *Chossis, I., Lane, C., Gache, P., Michaud, P. A., Pécoud, A., Rollnick, S., & Daepfen, J. B. (2007). Effect of training on primary care residents' performance in brief alcohol intervention: A randomized controlled trial. *Journal of General Internal Medicine*, 22(8), 1144–1149. <https://doi.org/10.1007/s11606-007-0240-2>
- Christensen, R. H. B. (2011). A tutorial on fitting cumulative link mixed models with clmm2 from the ordinal package. *Analysis*, 1(1), 1–10.
- *Christian, J., Bessesen, D., Byers, T., Christian, K., Goldstein, M., & Bock, B. (2008). Clinic-based support to help overweight patients with type 2 diabetes increase physical activity and lose weight. *Archives of Internal Medicine*, 168, 141–146. <https://doi.org/10.1001/archinternmed.2007.13>
- Cipriani, A., Higgins, J. P., Geddes, J. R., & Salanti, G. (2013). Conceptual and technical challenges in network meta-analysis. *Annals of Internal Medicine*, 159(2), 130–137. <https://doi.org/10.7326/0003-4819-159-2-201307160-00008>
- Cochran, W. G., & Chambers, S. P. (1965). The planning of observational studies of human populations. *Journal of the Royal Statistical Society. Series A (General)*, 128(2), 234–266. <https://doi.org/10.2307/2344179>
- *Cohen, T. R., Hazell, T. J., Vanstone, C. A., Rodd, C., & Weiler, H. A. (2016). A family-centered lifestyle intervention for obese six- to eight-year-old children: Results from a one-year randomized controlled trial conducted in Montreal, Canada. *Canadian Journal of Public Health*, 107(4–5), e453–e460. <https://doi.org/10.17269/CJPH.107.5470>
- Conn, V. S., Valentine, J. C., & Cooper, H. M. (2002). Interventions to increase physical activity among aging adults: A meta-analysis. *Annals of Behavioral Medicine*, 24(3), 190–200. https://doi.org/10.1207/S15324796ABM2403_04
- *Cordes, J., Thünker, J., Regenbrecht, G., Zielasek, J., Correll, C. U., Schmidt-Kraepelin, C., Lange-Asschenfeldt, C., Agelink, M. W., Kahl, K. G., Gaebel, W., Klimke, A., & Hauner, H. (2014). Can an early weight management program (WMP) prevent olanzapine (OLZ)-induced disturbances in body weight, blood glucose and lipid metabolism? Twenty-four- and 48-week results from a 6-month randomized trial. *The World Journal of Biological Psychiatry*, 15(3), 229–241. <https://doi.org/10.3109/15622975.2011.592546>
- *Cornman, D. H., Schmiede, S. J., Bryan, A., Benziger, T. J., & Fisher, J. D. (2007). An information-motivation-behavioral skills (IMB) model-based HIV prevention intervention for truck drivers in India. *Social Science & Medicine*, 64(8), 1572–1584. <https://doi.org/10.1016/j.socscimed.2006.11.011>
- *Coutmans, J. M. J., Bolman, C. A. W., Oenema, A., & Lechner, L. (2022). The effects of a web-based computer-tailored diet and physical activity intervention based on self-determination theory and motivational interviewing: A randomized controlled trial. *Internet Interventions*, 28, Article 100537. <https://doi.org/10.1016/j.invent.2022.100537>
- *Courcoulas, A. P., Gallagher, J. W., Neiberg, R. H., Eagleton, E. B., DeLany, J. P., Lang, W., Punchai, S., Gourash, W., & Jakicic, J. M. (2020). Bariatric surgery vs lifestyle intervention for diabetes treatment: 5-year outcomes from a randomized trial. *The Journal of Clinical Endocrinology and Metabolism*, 105(3), 866–876. <https://doi.org/10.1210/clinem/dgaa006>
- Cowan, N. (2010). The magical mystery four: How is working memory capacity limited, and why? *Current Directions in Psychological Science*, 19(1), 51–57. <https://doi.org/10.1177/0963721409359277>
- Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal of Psychiatry*, 150(6), 782–786. <https://doi.org/10.1192/bjp.150.6.782>
- Cunningham, C. O., Arnsten, J. H., Zhang, C., Heo, M., Bachhuber, M. A., Jost, J. J., Grossberg, R., Stein, M. R., & Sohlner, N. L. (2020). Abstinence-reinforcing contingency management improves HIV viral load suppression among HIV-infected people who use drugs: A randomized controlled

- trial. *Drug and Alcohol Dependence*, 216, Article 108230. <https://doi.org/10.1016/j.drugalcdep.2020.108230>
- *Cupples, M. E., & McKnight, A. (1994). Randomised controlled trial of health promotion in general practice for patients at high cardiovascular risk. *The BMJ*, 309(6960), 993–996. <https://doi.org/10.1136/bmj.309.6960.993>
- Dai, W., Palmer, R., Sunderrajan, A., Durantini, M., Sánchez, F., Glasman, L. R., Chen, F. X., & Albarracín, D. (2020). More behavioral recommendations produce more change: A meta-analysis of efficacy of multibehavior recommendations to reduce nonmedical substance use. *Psychology of Addictive Behaviors*, 34(7), 709–725. <https://doi.org/10.1037/adb0000586>
- *de Freitas, P. P., de Menezes, M. C., Dos Santos, L. C., Pimenta, A. M., Ferreira, A. V. M., & Lopes, A. C. S. (2020). The transtheoretical model is an effective weight management intervention: A randomized controlled trial. *BMC Public Health*, 20(1), Article 652. <https://doi.org/10.1186/s12889-020-08796-1>
- *DeBar, L. L., Ritenbaugh, C., Aickin, M., Orwoll, E., Elliot, D., Dickerson, J., Vuckovic, N., Stevens, V. J., Moe, E., & Irving, L. M. (2006). Youth: A health plan-based lifestyle intervention increases bone mineral density in adolescent girls. *Archives of Pediatrics & Adolescent Medicine*, 160(12), 1269–1276. <https://doi.org/10.1001/archpedi.160.12.1269>
- Dechartres, A., Trinquart, L., Faber, T., & Ravaud, P. (2016). Empirical evaluation of which trial characteristics are associated with treatment effect estimates. *Journal of Clinical Epidemiology*, 77, 24–37. <https://doi.org/10.1016/j.jclinepi.2016.04.005>
- Derogatis, L. R., Lipman, R. S., & Covi, L. (1973). SCL-90: An outpatient psychiatric rating scale—Preliminary report. *Psychopharmacology Bulletin*, 9(1), 13–28.
- *Desai, J. R., Vazquez-Benitez, G., Taylor, G., Johnson, S., Anderson, J., Garrett, J. E., Gilmer, T., Vue-Her, H., Rinn, S., Engel, K., Schiff, J., & O'Connor, P. J. (2020). The effects of financial incentives on diabetes prevention program attendance and weight loss among low-income patients: The we can prevent diabetes cluster-randomized controlled trial. *BMC Public Health*, 20(1), Article 1587. <https://doi.org/10.1186/s12889-020-09683-5>
- *Dévieux, J. G., Rosenberg, R., Jean-Gilles, M., Villalba, K., Attonito, J., Lerner, B., Saxena, A., & Stein, J. (2022). Effectiveness of a Cognitive Behavioral Randomized Controlled Trial for People Living with HIV who are heavy drinkers: The Holistic Health Recovery Program (HHRP) Trial in Miami. *Journal of Clinical Psychology in Medical Settings*, 29(3), 498–508. <https://doi.org/10.1007/s10880-022-09878-5>
- *Diao, H., Wang, H., Yang, L., & Li, T. (2020). The impacts of multiple obesity-related interventions on quality of life in children and adolescents: A randomized controlled trial. *Health and Quality of Life Outcomes*, 18(1), Article 213. <https://doi.org/10.1186/s12955-020-01459-0>
- *DiClemente, R. J., Wingood, G. M., Harrington, K. F., Lang, D. L., Davies, S. L., Hook, E. W., III, Oh, M. K., Crosby, R. A., Hertzberg, V. S., Gordon, A. B., Hardin, J. W., Parker, S., & Robillard, A. (2004). Efficacy of an HIV prevention intervention for African American adolescent girls: A randomized controlled trial. *JAMA: Journal of the American Medical Association*, 292(2), 171–179. <https://doi.org/10.1001/jama.292.2.171>
- Dijksterhuis, A., Aarts, H., Bargh, J. A., & Van Knippenberg, A. (2000). On the relation between associative strength and automatic behavior. *Journal of Experimental Social Psychology*, 36(5), 531–544. <https://doi.org/10.1006/jesp.2000.1427>
- *Dimitrov Ulian, M., Pinto, A. J., de Moraes Sato, P., B Benatti, F., Lopes de Campos-Ferraz, P., Coelho, D., Roble, O. J., Sabatini, F., Perez, I., Aburad, L., Vessoni, A., Fernandez Unsain, R., Macedo Rogero, M., Toporcov, T. N., de Sá-Pinto, A. L., Gualano, B., & B Scagliusi, F. (2018). Effects of a new intervention based on the health at every size approach for the management of obesity: The “Health and Wellness in Obesity” study. *PLOS ONE*, 13(7), Article e0198401. <https://doi.org/10.1371/journal.pone.0198401>
- *Donenberg, G., Emerson, E., & Kendall, A. D. (2018). HIV-risk reduction intervention for juvenile offenders on probation: The PHAT Life group randomized controlled trial. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association*, 37(4), 364–374. <https://doi.org/10.1037/hea0000582>
- Dotson, K. B., Dunn, M. E., & Bowers, C. A. (2015). Stand-alone personalized normative feedback for college student drinkers: A meta-analytic review, 2004 to 2014. *PLOS ONE*, 10(10), Article e0139518. <https://doi.org/10.1371/journal.pone.0139518>
- Doumas, D. M., Esp, S., Flay, B., & Bond, L. (2017). A randomized controlled trial testing the efficacy of a brief online alcohol intervention for high school seniors. *Journal of Studies on Alcohol and Drugs*, 78(5), 706–715. <https://doi.org/10.15288/jsad.2017.78.706>
- *Dove, J., Ferreira, M., Galbreath, M., Chandran, R., Beavers, K., Serra, M., Li, R., Jitomir, J., Shelmadine, B., Buford, T. W., Nassar, E., Bunn, J., Hudson, G. M., Parker, A., Rasmussen, C. G., Fredenburg, K., Wooddy, M., Wilson, R., & Kreide, R. (2008). Effects of a multi-component school-based intervention on health markers, body composition, physical fitness, and psychological measures in overweight and obese adolescent females. *The FASEB Journal*, 22(S2), Article 793. https://doi.org/10.1096/fasebj.22.2_supplement.793
- *Doyle, A. C., Goldschmidt, A., Huang, C., Winzelberg, A. J., Taylor, C. B., & Wilfley, D. E. (2008). Reduction of overweight and eating disorder symptoms via the Internet in adolescents: A randomized controlled trial. *Journal of Adolescent Health*, 43(2), 172–179. <https://doi.org/10.1016/j.jadohealth.2008.01.011>
- *Dracup, K., Meleis, A. I., Clark, S., Clyburn, A., Shields, L., & Staley, M. (1984). Group counseling in cardiac rehabilitation: Effect on patient compliance. *Patient Education and Counseling*, 6(4), 169–177. [https://doi.org/10.1016/0738-3991\(84\)90053-3](https://doi.org/10.1016/0738-3991(84)90053-3)
- *Duggan, C., Tapsoba, J. D. D., Shivappa, N., Harris, H. R., Hébert, J. R., Wang, C. Y., & McTiernan, A. (2021). Changes in dietary inflammatory index patterns with weight loss in women: A randomized controlled trial. *Cancer Prevention Research*, 14(1), 85–94. <https://doi.org/10.1158/1940-6207.CAPR-20-0181>
- *Duijzer, G., Haveman-Nies, A., Jansen, S. C., Beek, J. T., van Bruggen, R., Willink, M. G. J., Hiddink, G. J., & Feskens, E. J. M. (2017). Effect and maintenance of the SLIMMER diabetes prevention lifestyle intervention in Dutch primary healthcare: A randomised controlled trial. *Nutrition & Diabetes*, 7(5), Article e268. <https://doi.org/10.1038/nutd.2017.21>
- Dushay, R. A., Singer, M., Weeks, M. R., Rohena, L., & Gruber, R. (2001). Lowering HIV risk among ethnic minority drug users: Comparing culturally targeted intervention to a standard intervention. *The American Journal of Drug and Alcohol Abuse*, 27(3), 501–524. <https://doi.org/10.1081/ADA-100104515>
- *Eaton, L. A., Kalichman, S. C., Kalichman, M. O., Driffin, D. D., Baldwin, R., Zohren, L., & Conway-Washington, C. (2018). Randomised controlled trial of a sexual risk reduction intervention for STI prevention among men who have sex with men in the USA. *Sexually Transmitted Infections*, 94(1), 40–45. <https://doi.org/10.1136/sextans-2016-052835>
- *El-Bassel, N., McCrimmon, T., Mergenova, G., Chang, M., Terlikbayeva, A., Primbetova, S., Kuskulov, A., Baisarkin, B., Denebayeva, A., Kurmetova, K., & Witte, S. S. (2021). A cluster-randomized controlled trial of a combination HIV risk reduction and microfinance intervention for female sex workers who use drugs in Kazakhstan. *Journal of the International AIDS Society*, 24(5), Article e25682. <https://doi.org/10.1002/jia2.25682>
- *El-Bassel, N., White, S., Gilbert, L., Wu, E., Chang, M., Hill, J., & Steinglass, P. (2003). The efficacy of a relationship-based HIV/STD prevention program for heterosexual couples. *American Journal of Public Health*, 93(6), 963–969. <https://doi.org/10.2105/AJPH.93.6.963>
- *Elkind-Hirsch, K. E., Seidemann, E., & Harris, R. (2020). A randomized trial of dapagliflozin and metformin, alone and combined, in overweight women after gestational diabetes mellitus. *American Journal of Obstetrics*

- & *Gynecology MFM*, 2(3), Article 100139. <https://doi.org/10.1016/j.ajogmf.2020.100139>
- *Eriksen, A. K., Hansen, R. D., Borre, M., Larsen, R. G., Jensen, J. M., Overgaard, K., Borre, M., Kyrø, C., Landberg, R., Olsen, A., & Tjønneland, A. (2017). A lifestyle intervention among elderly men on active surveillance for non-aggressive prostate cancer: A randomised feasibility study with whole-grain rye and exercise. *Trials*, 18(1), Article 20. <https://doi.org/10.1186/s13063-016-1734-1>
- *Espeland, M. A., Carmichael, O., Hayden, K., Neiberg, R. H., Newman, A. B., Keller, J. N., Wadden, T. A., Rapp, S. R., Hill, J. O., Horton, E. S., Johnson, K. C., Wagenknecht, L., Wing, R. R., & Action for Health In Diabetes Brain Magnetic Resonance Imaging (Look AHEAD Brain) and Action for Health Movement and Memory Ancillary Study Research Groups. (2018). Long-term impact of weight loss intervention on changes in cognitive function: Exploratory analyses from the action for health in diabetes randomized controlled clinical trial. *The Journals of Gerontology: Series A*, 73(4), 484–491. <https://doi.org/10.1093/geronol/glx165>
- *Estabrooks, P. A., & Smith-Ray, R. L. (2008). Piloting a behavioral intervention delivered through interactive voice response telephone messages to promote weight loss in a pre-diabetic population. *Patient Education and Counseling*, 72(1), 34–41. <https://doi.org/10.1016/j.pec.2008.01.007>
- EuroQol Group. (1990). EuroQol—A new facility for the measurement of health-related quality of life. *Health Policy*, 16(3), 199–208. [https://doi.org/10.1016/0168-8510\(90\)90421-9](https://doi.org/10.1016/0168-8510(90)90421-9)
- Farnham, P. G., Holtgrave, D. R., Sansom, S. L., & Hall, H. I. (2010). Medical costs averted by HIV prevention efforts in the United States, 1991–2006. *Journal of Acquired Immune Deficiency Syndromes*, 54(5), 565–567. <https://doi.org/10.1097/QAI.0b013e3181e461b2>
- *Feaster, D. J., Mitrani, V. B., Burns, M. J., McCabe, B. E., Brincks, A. M., Rodríguez, A. E., Asthana, D., & Robbins, M. S. (2010). A randomized controlled trial of Structural Ecosystems Therapy for HIV medication adherence and substance abuse relapse prevention. *Drug and Alcohol Dependence*, 111(3), 227–234. <https://doi.org/10.1016/j.drugalcdep.2010.04.017>
- *Feldstein Ewing, S., Bryan, A. D., Dash, G. F., Lovejoy, T. I., Borsari, B., & Schmiege, S. J. (2022). Randomized controlled trial of motivational interviewing for alcohol and cannabis use within a predominantly Hispanic adolescent sample. *Experimental and Clinical Psychopharmacology*, 30(3), 287–299. <https://doi.org/10.1037/pha0000445>
- *Fernandez, A. C., Waller, R., Walton, M. A., Bonar, E. E., Ignacio, R. V., Chermack, S. T., Cunningham, R. M., Booth, B. M., Ilgen, M. A., Barry, K. L., & Blow, F. C. (2019). Alcohol use severity and age moderate the effects of brief interventions in an emergency department randomized controlled trial. *Drug and Alcohol Dependence*, 194, 386–394. <https://doi.org/10.1016/j.drugalcdep.2018.10.021>
- *Ferrante, J. M., Devine, K. A., Bator, A., Rodgers, A., Ohman-Strickland, P. A., Bandera, E. V., & Hwang, K. O. (2020). Feasibility and potential efficacy of commercial mHealth/eHealth tools for weight loss in African American breast cancer survivors: Pilot randomized controlled trial. *Translational Behavioral Medicine*, 10(4), 938–948. <https://doi.org/10.1093/tbm/iby124>
- *Ferré, R., Plana, N., Merino, J., Aragonès, G., Girona, J., Heras, M., Coll, B., Cos, R., & Masana, L. (2012). Effects of therapeutic lifestyle changes on peripheral artery tonometry in patients with abdominal obesity. *Nutrition, Metabolism, and Cardiovascular Diseases: NMCD*, 22(2), 95–102. <https://doi.org/10.1016/j.numecd.2010.04.008>
- *Firestone, R., Moorsmith, R., James, S., Urey, M., Greifinger, R., Lloyd, D., Hartenberger-Toby, L., Gausman, J., & Sanoe, M. (2016). Intensive group learning and on-site services to improve sexual and reproductive health among young adults in Liberia: A randomized evaluation of HealthyActions. *Global Health, Science and Practice*, 4(3), 435–451. <https://doi.org/10.9745/GHSP-D-16-00074>
- Firth, J., Solmi, M., Wootton, R. E., Vancampfort, D., Schuch, F. B., Hoare, E., Gilbody, S., Torous, J., Teasdale, S. B., Jackson, S. E., Smith, L., Eaton, M., Jacka, F. N., Veronese, N., Marx, W., Ashdown-Franks, G., Siskind, D., Sarris, J., Rosenbaum, S., ... Stubbs, B. (2020). A meta-review of “lifestyle psychiatry”: The role of exercise, smoking, diet and sleep in the prevention and treatment of mental disorders. *World Psychiatry*, 19(3), 360–380. <https://doi.org/10.1002/wps.20773>
- Fishbach, A., Shah, J. Y., & Kruglanski, A. W. (2004). Emotional transfer in goal systems. *Journal of Experimental Social Psychology*, 40(6), 723–738. <https://doi.org/10.1016/j.jesp.2004.04.001>
- Fisher, J. D., & Fisher, W. A. (1992). Changing AIDS-risk behavior. *Psychological Bulletin*, 111(3), 455–474. <https://doi.org/10.1037/0033-2909.111.3.455>
- Fisher, J. D., Fisher, W. A., Amico, K. R., & Harman, J. J. (2006). An information-motivation-behavioral skills model of adherence to antiretroviral therapy. *Health Psychology*, 25(4), 462–473. <https://doi.org/10.1037/0278-6133.25.4.462>
- Fleig, L., Kerschreiter, R., Schwarzer, R., Pomp, S., & Lippke, S. (2014). ‘Sticking to a healthy diet is easier for me when I exercise regularly’: Cognitive transfer between physical exercise and healthy nutrition. *Psychology & Health*, 29(12), 1361–1372. <https://doi.org/10.1080/08870446.2014.930146>
- Fleig, L., Lippke, S., Pomp, S., & Schwarzer, R. (2011). Intervention effects of exercise self-regulation on physical exercise and eating fruits and vegetables: A longitudinal study in orthopedic and cardiac rehabilitation. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 53(3), 182–187. <https://doi.org/10.1016/j.ypmed.2011.06.019>
- *Fogelholm, M., Kukkonen-Harjula, K., Nenonen, A., & Pasanen, M. (2000). Effects of walking training on weight maintenance after a very-low-energy diet in premenopausal obese women. *Archives of Internal Medicine*, 160(14), 2177–2184. <https://doi.org/10.1001/archinte.160.14.2177>
- *Frie, K., Hartmann-Boyce, J., Jebb, S. A., & Aveyard, P. (2020). Effectiveness of a self-regulation intervention for weight loss: A randomized controlled trial. *British Journal of Health Psychology*, 25(3), 652–676. <https://doi.org/10.1111/bjhp.12436>
- *Fromme, K., & Corbin, W. (2004). Prevention of heavy drinking and associated negative consequences among mandated and voluntary college students. *Journal of Consulting and Clinical Psychology*, 72(6), 1038–1049. <https://doi.org/10.1037/0022-006X.72.6.1038>
- *Frommer, M. S., Mandryk, J. A., Edye, B. V., Healey, S., Berry, G., & Ferguson, D. A. (1990). A randomised controlled trial of counseling in a workplace setting for coronary heart disease risk factor modification: Effects on blood pressure. *Asia-Pacific Journal of Public Health*, 4(1), 25–33. <https://doi.org/10.1177/101053959000400105>
- *Fu, C. P., Oczypok, E. E., Ali, H., DeLany, J. P., Reeves, V. L., Chang, R. F., & Kershaw, E. E. (2022). Effect of physical activity in a weight loss program on circulating total ANGPTL8 concentrations in northern Americans with obesity: A prospective randomized controlled trial. *Nutrition, Metabolism, and Cardiovascular Diseases: NMCD*, 32(7), 1725–1733. <https://doi.org/10.1016/j.numecd.2022.04.006>
- *Gabiola, J., Morales, D., Quizon, O., Cadiz, R. I., Feliciano, K., Ruiz, R. L., Aguatis, C. J., Mararac, T., Rojina, J., Garcia, A., Hedlin, H., Cullen, M., & Palaniappan, L. (2020). The effectiveness of lifestyle with diet and physical activity education program among prehypertensives and stage 1 hypertensives in an urban community setting (enlighten) study. *Journal of Community Health: The Publication for Health Promotion and Disease Prevention*, 45(3), 478–487. <https://doi.org/10.1007/s10900-019-00764-0>
- *Gabriele, J. (2008). *Effects of nondirective and directive support on weight loss in an e-counseling intervention* [Ph.D. dissertation, Washington University]. Dissertations & Theses: Full Text. (Publication No. AAT 3332093).
- *Gaede, P., Beck, M., Vedel, P., & Pedersen, O. (2001). Limited impact of lifestyle education in patients with Type 2 diabetes mellitus and microalbuminuria: Results from a randomized intervention study.

- Diabetic Medicine*, 18(2), 104–108. <https://doi.org/10.1046/j.1464-5491.2001.00444.x>
- *Galárraga, O., Harries, J., Maughan-Brown, B., Cooper, D., Short, S. E., Lurie, M. N., & Harrison, A. (2018). The *Empower Nudge* lottery to increase dual protection use: A proof-of-concept randomised pilot trial in South Africa. *Reproductive Health Matters*, 26(52), 67–80. <https://doi.org/10.1080/09688080.2018.1510701>
- *García Pérez de Sevilla, G., Barceló Guido, O., De la Cruz, M. P., Blanco Fernández, A., Alejo, L. B., Montero Martínez, M., & Pérez-Ruiz, M. (2021). Adherence to a lifestyle exercise and nutrition intervention in university employees during the COVID-19 pandemic: A randomized controlled trial. *International Journal of Environmental Research and Public Health*, 18(14), Article 7510. <https://doi.org/10.3390/ijerph18147510>
- *Geliebter, A., Maher, M. M., Gerace, L., Gutin, B., Heymsfield, S. B., & Hashim, S. A. (1997). Effects of strength or aerobic training on body composition, resting metabolic rate, and peak oxygen consumption in obese dieting subjects. *The American Journal of Clinical Nutrition*, 66(3), 557–563. <https://doi.org/10.1093/ajcn/66.3.557>
- *Gepner, Y., Shelef, I., Schwarzfuchs, D., Zelicha, H., Tene, L., Yaskolka Meir, A., Tsaban, G., Cohen, N., Bril, N., Rein, M., Serfaty, D., Kenigsbuch, S., Komy, O., Wolak, A., Chassidim, Y., Golan, R., Avni-Hassid, H., Bilitzky, A., Sarusi, B., ... Shai, I. (2018). Effect of distinct lifestyle interventions on mobilization of fat storage pools: CENTRAL magnetic resonance imaging randomized controlled trial. *Circulation*, 137(11), 1143–1157. <https://doi.org/10.1161/CIRCULATIONAHA.117.030501>
- *Gibson, C. A., Gupta, A., Greene, J. L., Lee, J., Mount, R. R., & Sullivan, D. K. (2020). Feasibility and acceptability of a televideo physical activity and nutrition program for recent kidney transplant recipients. *Pilot and Feasibility Studies*, 6(1), Article 126. <https://doi.org/10.1186/s40814-020-00672-4>
- Glanz, K., & Bishop, D. B. (2010). The role of behavioral science theory in development and implementation of public health interventions. *Annual Review of Public Health*, 31(1), 399–418. <https://doi.org/10.1146/annurev.publhealth.012809.103604>
- *Glasgow, R. E., Toobert, D. J., Hampson, S. E., Brown, J. E., Lewinsohn, P. M., & Donnelly, J. (1992). Improving self-care among older patients with type II diabetes: The “Sixty Something...” study. *Patient Education and Counseling*, 19(1), 61–74. [https://doi.org/10.1016/0738-3991\(92\)90102-O](https://doi.org/10.1016/0738-3991(92)90102-O)
- *Glaysher, M. A., Ward, J., Aldhwayan, M., Ruban, A., Prechtel, C. G., Fisk, H. L., Chhina, N., Al-Najim, W., Smith, C., Klimowska-Nassar, N., Johnson, N., Falaschetti, E., Goldstone, A. P., Miras, A. D., Byrne, J. P., Calder, P. C., & Teare, J. P. (2021). The effect of a duodenal-jejunal bypass liner on lipid profile and blood concentrations of long chain polyunsaturated fatty acids. *Clinical Nutrition*, 40(4), 2343–2354. <https://doi.org/10.1016/j.clnu.2020.10.026>
- *Go, V. F., Frangakis, C., Minh, N. L., Latkin, C., Ha, T. V., Mo, T. T., Sripaipan, T., Davis, W. W., Zelaya, C., Vu, P. T., Celentano, D. D., & Quan, V. M. (2015). Efficacy of a multi-level intervention to reduce injecting and sexual risk behaviors among HIV-infected people who inject drugs in Vietnam: A four-arm randomized controlled trial. *PLOS ONE*, 10(5), Article e0125909. <https://doi.org/10.1371/journal.pone.0125909>
- *Gokulakrishnan, K., Ranjani, H., Weber, M. B., Pandey, G. K., Anjana, R. M., Balasubramanyam, M., Prabhakaran, D., Tandon, N., Narayan, K. M., & Mohan, V. (2017). Effect of lifestyle improvement program on the biomarkers of adiposity, inflammation and gut hormones in overweight/obese Asian Indians with prediabetes. *Acta Diabetologica*, 54(9), 843–852. <https://doi.org/10.1007/s00592-017-1015-9>
- Gong, Q., Zhang, P., Wang, J., Ma, J., An, Y., Chen, Y., Zhang, B., Feng, X., Li, H., Chen, X., Cheng, Y. J., Gregg, E. W., Hu, Y., Bennett, P. H., Li, G., Qian, X., Zhang, L., Hui, Y., He, S., ... the Da Qing Diabetes Prevention Study Group. (2019). Morbidity and mortality after lifestyle intervention for people with impaired glucose tolerance: 30-year results of the Da Qing Diabetes Prevention Outcome Study. *The Lancet. Diabetes & Endocrinology*, 7(6), 452–461. [https://doi.org/10.1016/S2213-8587\(19\)30093-2](https://doi.org/10.1016/S2213-8587(19)30093-2)
- *Goodwin, P. J., Segal, R. J., Vallis, M., Ligiabel, J. A., Pond, G. R., Robidoux, A., Findlay, B., Gralow, J. R., Mukherjee, S. D., Levine, M., & Pritchard, K. I. (2020). The LISA randomized trial of a weight loss intervention in postmenopausal breast cancer. *NPI Breast Cancer*, 6(1), Article 6. <https://doi.org/10.1038/s41523-020-0149-z>
- *Gordon, N. F., English, C. D., Contractor, A. S., Salmon, R. D., Leighton, R. F., Franklin, B. A., & Haskell, W. L. (2002). Effectiveness of three models for comprehensive cardiovascular disease risk reduction. *The American Journal of Cardiology*, 89(11), 1263–1268. [https://doi.org/10.1016/S0002-9149\(02\)02323-8](https://doi.org/10.1016/S0002-9149(02)02323-8)
- *Gorini, G., Carreras, G., Giordano, L., Anghinoni, E., Iossa, A., Coppo, A., Talassi, F., Galavotti, M., Chellini, E., & the SPRINT Working Group. (2012). The Pap smear screening as an occasion for smoking cessation and physical activity counselling: Effectiveness of the SPRINT randomized controlled trial. *BMC Public Health*, 12(1), Article 740. <https://doi.org/10.1186/1471-2458-12-740>
- *Griffin, S. J., Simmons, R. K., Prevost, A. T., Williams, K. M., Hardeman, W., Sutton, S., Brage, S., Ekelund, U., Parker, R. A., Wareham, N. J., Kinmonth, A. L., & the ADDITION-Plus study team. (2014). Multiple behaviour change intervention and outcomes in recently diagnosed type 2 diabetes: The ADDITION-Plus randomised controlled trial. *Diabetologia*, 57(7), 1308–1319. <https://doi.org/10.1007/s00125-014-3236-6>
- *Grilo, C. M., White, M. A., Masheb, R. M., Ivezaj, V., Morgan, P. T., & Gueorguieva, R. (2020). Randomized controlled trial testing the effectiveness of adaptive “SMART” stepped-care treatment for adults with binge-eating disorder comorbid with obesity. *American Psychologist*, 75(2), 204–218. <https://doi.org/10.1037/amp0000534>
- *Grimley, D. M., & Hook, E. W., III. (2009). A 15-minute interactive, computerized condom use intervention with biological endpoints. *Sexually Transmitted Diseases*, 36(2), 73–78. <https://doi.org/10.1097/OLQ.0b013e31818eca81>
- *Groenendijk, I., Chan, R., Woo, J., Ong, S., Parikh, P., Bragt, M. C. E., & de Groot, L. C. P. G. M. (2020). A combined nutrition and exercise intervention influences serum vitamin B-12 and 25-hydroxyvitamin D and bone turnover of healthy Chinese middle-aged and older adults. *The Journal of Nutrition*, 150(8), 2112–2119. <https://doi.org/10.1093/jn/nxaa149>
- Grubbs, L. M., & Tabano, M. (2000). Use of sunscreen in health care professionals. The health belief model. *Cancer Nursing*, 23(3), 164–167. <https://doi.org/10.1097/00002820-200006000-00002>
- *Gruppo di Ricerca del Progetto Romano di Prevenzione della Cardiopatia Coronarica. (1980). Variazione dei fattori di rischio in quattro anni nel progetto romano di prevenzione della cardiopatia coronarica [Assessment of absolute global cardiovascular risk: The chart of the heart project]. *Giornale Italiano di Cardiologia*, 10, 204–215.
- *Gryczynski, J., Mitchell, S. G., Schwartz, R. P., Dusek, K., O’Grady, K. E., Cowell, A. J., Barbosa, C., Barnosky, A., & DiClemente, C. C. (2021). Computer- vs. nurse practitioner-delivered brief intervention for adolescent marijuana, alcohol, and sex risk behaviors in school-based health centers. *Drug and Alcohol Dependence*, 218, Article 108423. <https://doi.org/10.1016/j.drugalcdep.2020.108423>
- *Gryczynski, J., Nordeck, C. D., Welsh, C., Mitchell, S. G., O’Grady, K. E., & Schwartz, R. P. (2021). Preventing hospital readmission for patients with comorbid substance use disorder: A randomized trial. *Annals of Internal Medicine*, 174(7), 899–909. <https://doi.org/10.7326/M20-5475>
- *Guo, Y., Li, Y., Yu, C., Xu, H., Hong, Y. A., Wang, X., Zhang, N., Zeng, Y., Monroe-Wise, A., Li, L., Liu, C., Cai, W., & Lin, A. (2022). Long-term effects of a social media-based intervention (Run4Love) on depressive symptoms of people living with HIV: 3-year follow-up of a randomized controlled trial. *Journal of Medical Internet Research*, 24(6), Article e36809. <https://doi.org/10.2196/36809>

- *Gür, F., Can Gür, G., & Okanlı, A. (2017). The effect of the cognitive-behavioral model-based psychoeducation and exercise intervention on quality of life in alcohol use disorder. *Archives of Psychiatric Nursing*, 31(6), 541–548. <https://doi.org/10.1016/j.apnu.2017.07.005>
- *Hadley, W., Brown, L. K., Barker, D., Warren, J., Weddington, P., Fortune, T., & Juzang, I. (2016). Work it out together: Preliminary efficacy of a parent and adolescent DVD and workbook intervention on adolescent sexual and substance use attitudes and parenting behaviors. *AIDS and Behavior*, 20(9), 1961–1972. <https://doi.org/10.1007/s10461-016-1418-6>
- *Haire-Joshu, D., Schwarz, C. D., Steger-May, K., Lapka, C., Schechtman, K., Brownson, R. C., & Tabak, R. G. (2018). A randomized trial of weight change in a national home visiting program. *American Journal of Preventive Medicine*, 54(3), 341–351. <https://doi.org/10.1016/j.amepre.2017.12.012>
- Hall, D., Jarrold, C., Towse, J. N., & Zarandi, A. L. (2015). The developmental influence of primary memory capacity on working memory and academic achievement. *Developmental Psychology*, 51(8), 1131–1147. <https://doi.org/10.1037/a0039464>
- Hamilton, M. (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery & Psychiatry*, 23(1), 56–62. <https://doi.org/10.1136/jnnp.23.1.56>
- *Hanson, T., Alessi, S. M., & Petry, N. M. (2008). Contingency management reduces drug-related human immunodeficiency virus risk behaviors in cocaine-abusing methadone patients. *Addiction*, 103(7), 1187–1197. <https://doi.org/10.1111/j.1360-0443.2008.02216.x>
- *Hardcastle, S. J., Taylor, A. H., Bailey, M. P., Harley, R. A., & Hagger, M. S. (2013). Effectiveness of a motivational interviewing intervention on weight loss, physical activity and cardiovascular disease risk factors: A randomised controlled trial with a 12-month post-intervention follow-up. *The International Journal of Behavioral Nutrition and Physical Activity*, 10(1), Article 40. <https://doi.org/10.1186/1479-5868-10-40>
- Harkins, S., & Petty, R. (1981). The multiple source effect in persuasion: The effects of distraction. *Personality and Social Psychology Bulletin*, 7(4), 627–635. <https://doi.org/10.1177/014616728174019>
- *Harvey, S. M., Henderson, J. T., Thorburn, S., Beckman, L. J., Casillas, A., Mendez, L., & Cervantes, R. (2004). A randomized study of a pregnancy and disease prevention intervention for Hispanic couples. *Perspectives on Sexual and Reproductive Health*, 36(4), 162–169. <https://doi.org/10.1363/3616204>
- *Haskell, W. L., Alderman, E. L., Fair, J. M., Maron, D. J., Mackey, S. F., Superko, H. R., Williams, P. T., Johnstone, I. M., Champagne, M. A., Krauss, R. M., & Farquhar, J. W. (1994). Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinic cardiac events in men and women with coronary artery disease. *Circulation*, 89, 975–990. <https://doi.org/10.1161/01.CIR.89.3.975>
- *Haug, S., Paz Castro, R., Wenger, A., & Schaub, M. P. (2021). A mobile phone-based life-skills training program for substance use prevention among adolescents: Cluster-randomized controlled trial. *JMIR mHealth and uHealth*, 9(7), Article e26951. <https://doi.org/10.2196/26951>
- Hayes, A. F., & Rockwood, N. J. (2017). Regression-based statistical mediation and moderation analysis in clinical research: Observations, recommendations, and implementation. *Behaviour Research and Therapy*, 98, 39–57. <https://doi.org/10.1016/j.brat.2016.11.001>
- Hedges, L. V. (2007). Effect sizes in cluster-randomized designs. *Journal of Educational and Behavioral Statistics*, 32(4), 341–370. <https://doi.org/10.3102/1076998606298043>
- Hedges, L. V., & Olkin, I. (1985). *Statistical methods for meta-analysis*. Academic Press.
- *Heirich, M. A., Foote, A., Erfurt, J. C., & Konopka, B. (1993). Work-site physical fitness programs. Comparing the impact of different program designs on cardiovascular risks. *Journal of Occupational Medicine*, 35(5), 510–517.
- Henrich, J., Heine, S. J., & Norenzayan, A. (2010). The weirdest people in the world? *Behavioral and Brain Sciences*, 33(2–3), 61–83. <https://doi.org/10.1017/S0140525X0999152X>
- Henry, J. D., & Crawford, J. R. (2005). The short-form version of the Depression Anxiety Stress Scales (DASS-21): Construct validity and normative data in a large non-clinical sample. *British Journal of Clinical Psychology*, 44(2), 227–239. <https://doi.org/10.1348/014466505X29657>
- Heron, M. (2019). Deaths: Leading causes for 2017. *National Vital Statistics Reports*, 68(6), 1–77.
- *Hershberger, S. L., Wood, M. M., & Fisher, D. G. (2003). A cognitive-behavioral intervention to reduce HIV risk behaviors in crack and injection drug users. *AIDS and Behavior*, 7(3), 229–243. <https://doi.org/10.1023/A:1025487501743>
- Hershow, R. B., Reyes, H. L. M., Ha, T. V., Chander, G., Mai, N. V. T., Sripaipan, T., Frangakis, C., Dowdy, D. W., Latkin, C., Hutton, H. E., Pettifor, A., Maman, S., & Go, V. F. (2020). Longitudinal analysis of alcohol use and intimate partner violence perpetration among men with HIV in northern Vietnam. *Drug and Alcohol Dependence*, 213, Article 108098. <https://doi.org/10.1016/j.drugalcdep.2020.108098>
- *Heufelder, A. E., Saad, F., Bunck, M. C., & Gooren, L. (2009). Fifty-two-week treatment with diet and exercise plus transdermal testosterone reverses the metabolic syndrome and improves glycemic control in men with newly diagnosed type 2 diabetes and subnormal plasma testosterone. *Journal of Andrology*, 30(6), 726–733. <https://doi.org/10.2164/jandrol.108.007005>
- *Hickey, M. D., Ouma, G. B., Mattah, B., Pederson, B., DesLauriers, N. R., Mohamed, P., Obanda, J., Odhiambo, A., Njoroge, B., Otieno, L., Zoughbie, D. E., Ding, E. L., Fiorella, K. J., Bukusi, E. A., Cohen, C. R., Geng, E. H., & Salmen, C. R. (2021). The Kanyakla study: Randomized controlled trial of a microclinic social network intervention for promoting engagement and retention in HIV care in rural western Kenya. *PLOS ONE*, 16(9), Article e0255945. <https://doi.org/10.1371/journal.pone.0255945>
- *Hien, D. A., Wells, E. A., Jiang, H., Suarez-Morales, L., Campbell, A. N., Cohen, L. R., Miele, G. M., Killeen, T., Brigham, G. S., Zhang, Y., Hansen, C., Hodgkins, C., Hatch-Maillette, M., Brown, C., Kulaga, A., Kristman-Valente, A., Chu, M., Sage, R., Robinson, J. A., ... Nunes, E. V. (2009). Multisite randomized trial of behavioral interventions for women with co-occurring PTSD and substance use disorders. *Journal of Consulting and Clinical Psychology*, 77(4), 607–619. <https://doi.org/10.1037/a0016227>
- Hjermann, I., Holme, I., Velve Byre, K., & Leren, P. (1981). Effect of diet and smoking intervention on the incidence of coronary heart disease. Report from the Oslo Study Group of a randomised trial in healthy men. *Lancet*, 2(8259), 1303–1310. [https://doi.org/10.1016/S0140-6736\(81\)91338-6](https://doi.org/10.1016/S0140-6736(81)91338-6)
- Hofmann, W., Luhmann, M., Fisher, R. R., Vohs, K. D., & Baumeister, R. F. (2014). Yes, but are they happy? Effects of trait self-control on affective well-being and life satisfaction. *Journal of Personality*, 82(4), 265–277. <https://doi.org/10.1111/jopy.12050>
- *Hom, K., Branstetter, S., Zhang, J., Jarrett, T., Tompkins, N. O., Anesetti-Rothermel, A., Olfert, M., Richards, T., & Dino, G. (2013). Understanding physical activity outcomes as a function of teen smoking cessation. *Journal of Adolescent Health*, 53(1), 125–131. <https://doi.org/10.1016/j.jadohealth.2013.01.019>
- *Horvath, K. J., Lammert, S., MacLehose, R. F., Danh, T., Baker, J. V., & Carrico, A. W. (2019). A pilot study of a mobile app to support HIV antiretroviral therapy adherence among men who have sex with men who use stimulants. *AIDS and Behavior*, 23(11), 3184–3198. <https://doi.org/10.1007/s10461-019-02597-3>
- *Huang, S.-H., Weng, K.-P., Hsieh, K.-S., Ou, S. F., Lin, C. C., Chien, K. J., Liu, P. Y., & Ho, T. Y. (2007). Effects of a classroom-based weight-control intervention on cardiovascular disease in elementary-school obese children. *Acta Paediatrica Taiwanica*, 48(4), 201–206.

- Huber, P. (1967). Behavior of maximum likelihood estimates under nonstandard conditions. In L. M. Le Cam & J. Neyman (Eds.), *Proceedings of the fifth Berkeley symposium on mathematical statistics and probability* (Vol. 1, No. 1, pp. 221–233). The University of California Press.
- *Hunter, C. M., Peterson, A. L., Alvarez, L. M., Poston, W. C., Brundige, A. R., Haddock, C. K., Van Brunt, D. L., & Foreyt, J. P. (2008). Weight management using the internet a randomized controlled trial. *American Journal of Preventive Medicine*, 34(2), 119–126. <https://doi.org/10.1016/j.amepre.2007.09.026>
- *Hussien, S. (2007). *Impact of a culturally-sensitive lifestyle intervention on reducing risk factors for type 2 diabetes in Arab Canadian Muslim women* [Ph.D. dissertation, The University of Western Ontario]. Dissertations & Theses: Full Text (Publication No. AAT NR36701).
- Hyman, D. J., Pavlik, V. N., Taylor, W. C., Goodrick, G. K., & Moye, L. (2007). Simultaneous vs sequential counseling for multiple behavior change. *Archives of Internal Medicine*, 167(11), 1152–1158. <https://doi.org/10.1001/archinte.167.11.1152>
- *Ifejika, N. L., Bhadane, M., Cai, C. C., Noser, E. A., Grotta, J. C., & Savitz, S. I. (2020). Use of a smartphone-based mobile app for weight management in obese minority stroke survivors: Pilot randomized controlled trial with open blinded end point. *JMIR mHealth and uHealth*, 8(4), Article e17816. <https://doi.org/10.2196/17816>
- Irie, M., Asami, S., Nagata, S., Miyata, M., & Kasai, H. (2001). Relationships between perceived workload, stress and oxidative DNA damage. *International Archives of Occupational and Environmental Health*, 74(2), 153–157. <https://doi.org/10.1007/s004200000209>
- *Ismail, K., Bayley, A., Twist, K., Stewart, K., Ridge, K., Britneff, E., Greenough, A., Ashworth, M., Rundle, J., Cook, D. G., Whincup, P., Treasure, J., McCrone, P., Winkley, K., & Stahl, D. (2020). Reducing weight and increasing physical activity in people at high risk of cardiovascular disease: A randomised controlled trial comparing the effectiveness of enhanced motivational interviewing intervention with usual care. *Heart*, 106(6), 447–454. <https://doi.org/10.1136/heartjnl-2019-315656>
- *Iso, H., Shimamoto, T., Yokota, K., Sankai, T., Jacobs, D. R., Jr., & Komachi, Y. (1996). Community-based education classes for hypertension control. A 1.5-year randomized controlled trial. *Hypertension*, 27(4), 968–974. <https://doi.org/10.1161/01.HYP.27.4.968>
- *Jacobs, N., Clays, E., De Bacquer, D., De Backer, G., Dendale, P., Thijs, H., Bourdeaudhuij, I., & Claes, N. (2011). Effect of a tailored behavioral change program on a composite lifestyle change score: A randomized controlled trial. *Health Education Research*, 26(5), 886–895. <https://doi.org/10.1093/her/cyr046>
- *Jacobs, N., De Bourdeaudhuij, I. D., Thijs, H., & Claes, N. (2011). Effect of a cardiovascular program on health behavior and BMI in highly educated adults: A randomized controlled trial. *Patient Education and Counseling*, 85(1), 122–126. <https://doi.org/10.1016/j.pec.2010.08.024>
- *Jacot, W., Arnaud, A., Jarlier, M., Lefevre-Plesse, C., Dalivoust, P., Senesse, P., Azzedine, A., Tredan, O., Sadot-Lebouvier, S., Mas, S., Carayol, M., Bleuse, J. P., Gourgu, S., Janiszewski, C., Launay, S., D'Hondt, V., Lauridant, G., Grenier, J., Romieu, G., ... Vanlemmens, L. (2020). Brief hospital supervision of exercise and diet during adjuvant breast cancer therapy is not enough to relieve fatigue: A multicenter randomized controlled trial. *Nutrients*, 12(10), Article 3081. <https://doi.org/10.3390/nu12103081>
- James, E., Freund, M., Booth, A., Duncan, M. J., Johnson, N., Short, C. E., Wolfenden, L., Stacey, F. G., Kay-Lambkin, F., & Vandelanotte, C. (2016). Comparative efficacy of simultaneous versus sequential multiple health behavior change interventions among adults: A systematic review of randomised trials. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 89, 211–223. <https://doi.org/10.1016/j.ypmed.2016.06.012>
- *Jatuporn, S., Sangwatanaroj, S., Saengsiri, A. O., Rattanaprucks, S., Srimahachota, S., Uthayachalerm, W., Kuanoon, W., Panpakdee, O., Tangkijvanich, P., & Tosukhowong, P. (2003). Short-term effects of an intensive lifestyle modification program on lipid peroxidation and antioxidant systems in patients with coronary artery disease. *Clinical Hemorheology and Microcirculation*, 29(3–4), 429–436.
- *Jelalian, E., Evans, E. W., Rancourt, D., Ranzenhofer, L., Taylor, N., Hart, C., Seifer, R., Klinepiper, K., & Foster, G. D. (2020). JOIN for ME: Testing a scalable weight control intervention for adolescents. *Childhood Obesity*, 16(3), 192–203. <https://doi.org/10.1089/chi.2019.0053>
- *Jemmott, J. B., III, Zhang, J., Jemmott, L. S., Icard, L. D., Ngwane, Z., Makiwane, M., & O'Leary, A. (2019). Intervention increases physical activity and healthful diet among South African adolescents over 54 months: A randomized controlled trial. *Journal of Adolescent Health*, 65(1), 139–146. <https://doi.org/10.1016/j.jadohealth.2019.01.027>
- *Jemmott, L. S., Jemmott, J. B., III, Icard, L. D., & Hsu, J. (2020). Effects of church-based parent–child abstinence-only interventions on adolescents' sexual behaviors. *Journal of Adolescent Health*, 66(1), 107–114. <https://doi.org/10.1016/j.jadohealth.2019.07.021>
- *Jemmott, L. S., Jemmott, J. B., III, & O'Leary, A. (2007). Effects on sexual risk behavior and STD rate of brief HIV/STD prevention interventions for African American women in primary care settings. *American Journal of Public Health*, 97(6), 1034–1040. <https://doi.org/10.2105/AJPH.2003.020271>
- *Jensen, K., Nielsen, C., Ekstrøm, C. T., & Roessler, K. K. (2019). Physical exercise in the treatment of alcohol use disorder (AUD) patients affects their drinking habits: A randomized controlled trial. *Scandinavian Journal of Public Health*, 47(4), 462–468. <https://doi.org/10.1177/1403494818759842>
- *Jiskoot, G., Timman, R., Beerthuis, A., Dietz de Loos, A., Busschbach, J., & Laven, J. (2020). Weight reduction through a cognitive behavioral therapy lifestyle intervention in PCOS: The primary outcome of a randomized controlled trial. *Obesity*, 28(11), 2134–2141. <https://doi.org/10.1002/oby.22980>
- Joffe, M. M., & Rosenbaum, P. R. (1999). Invited commentary: Propensity scores. *American Journal of Epidemiology*, 150(4), 327–333. <https://doi.org/10.1093/oxfordjournals.aje.a010011>
- *Johnson, M. O., Dilworth, S. E., Taylor, J. M., & Neillands, T. B. (2011). Improving coping skills for self-management of treatment side effects can reduce antiretroviral medication nonadherence among people living with HIV. *Annals of Behavioral Medicine*, 41(1), 83–91. <https://doi.org/10.1007/s12160-010-9230-4>
- *Jolly, K., Sidhu, M. S., Hewitt, C. A., Coventry, P. A., Daley, A., Jordan, R., Heneghan, C., Singh, S., Ives, N., Adab, P., Jowett, S., Varghese, J., Nunan, D., Ahmed, K., Dowson, L., & Fitzmaurice, D. (2018). Self management of patients with mild COPD in primary care: Randomised controlled trial. *BMJ: British Medical Journal*, 361, Article k2241. <https://doi.org/10.1136/bmj.k2241>
- *Jospe, M. R., Roy, M., Brown, R. C., Haszard, J. J., Meredith-Jones, K., Fangupo, L. J., Osborne, H., Fleming, E. A., & Taylor, R. W. (2020). Intermittent fasting, Paleolithic, or Mediterranean diets in the real world: Exploratory secondary analyses of a weight-loss trial that included choice of diet and exercise. *The American Journal of Clinical Nutrition*, 111(3), 503–514. <https://doi.org/10.1093/ajcn/nqz330>
- Kalichman, S. C., Simbayi, L. C., Cain, D., Carey, K. B., Carey, M. P., Eaton, L., Harel, O., Mehlomakhulu, V., & Mwaba, K. (2014). Randomized community-level HIV prevention intervention trial for men who drink in South African alcohol-serving venues. *European Journal of Public Health*, 24(5), 833–839. <https://doi.org/10.1093/eurpub/ckt172>
- *Kalichman, S. C., Simbayi, L. C., Cloete, A., Clayford, M., Arnolds, W., Mxoli, M., Smith, G., Cherry, C., Shefer, T., Crawford, M., & Kalichman, M. O. (2009). Integrated gender-based violence and HIV Risk reduction intervention for South African men: Results of a quasi-experimental field trial. *Prevention Science*, 10(3), 260–269. <https://doi.org/10.1007/s11121-009-0129-x>

- *Kallio, V., Hämäläinen, H., Hakila, J., & Luurila, O. J. (1979). Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet*, 314(8152), 1091–1094. [https://doi.org/10.1016/S0140-6736\(79\)92502-9](https://doi.org/10.1016/S0140-6736(79)92502-9)
- *Kaner, E., Bland, M., Cassidy, P., Coulton, S., Dale, V., Deluca, P., Gilvarry, E., Godfrey, C., Heather, N., Myles, J., Newbury-Birch, D., Oyefeso, A., Parrott, S., Perryman, K., Phillips, T., Shepherd, J., & Drummond, C. (2013). Effectiveness of screening and brief alcohol intervention in primary care (SIPS trial): Pragmatic cluster randomised controlled trial. *BMJ: British Medical Journal*, 346, Article e8501. <https://doi.org/10.1136/bmj.e8501>
- Karlsson, P., & Bergmark, A. (2015). Compared with what? An analysis of control-group types in Cochrane and Campbell reviews of psychosocial treatment efficacy with substance use disorders. *Addiction*, 110(3), 420–428. <https://doi.org/10.1111/add.12799>
- *Karmali, S., Battram, D. S., Burke, S. M., Cramp, A., Johnson, A. M., Mantler, T., Morrow, D., Ng, V., Pearson, E. S., Petrella, R. J., Tucker, P., & Irwin, J. D. (2020). Perspectives and impact of a parent–child intervention on dietary intake and physical activity behaviours, parental motivation, and parental body composition: A randomized controlled trial. *International Journal of Environmental Research and Public Health*, 17(18), Article 6822. <https://doi.org/10.3390/ijerph17186822>
- *Karno, M. P., Rawson, R., Rogers, B., Spear, S., Grella, C., Mooney, L. J., Saitz, R., Kagan, B., & Glasner, S. (2021). Effect of screening, brief intervention and referral to treatment for unhealthy alcohol and other drug use in mental health treatment settings: A randomized controlled trial. *Addiction*, 116(1), 159–169. <https://doi.org/10.1111/add.15114>
- *Katz, D. A., Graber, M., Lounsbury, P., Vander Weg, M. W., Phillips, E. K., Clair, C., Horwitz, P. A., Cai, X., & Christensen, A. J. (2017). Multiple risk factor counseling to promote heart-healthy lifestyles in the chest pain observation unit: Pilot randomized controlled trial. *Academic Emergency Medicine*, 24(8), 968–982. <https://doi.org/10.1111/ace.13231>
- *Kawashima, M., Sano, K., Takechi, S., & Tsubota, K. (2018). Impact of lifestyle intervention on dry eye disease in office workers: A randomized controlled trial. *Journal of Occupational Health*, 60(4), 281–288. <https://doi.org/10.1539/joh.2017-0191-OA>
- *Kazemi, M., Pierson, R. A., McBairty, L. E., Chilibeck, P. D., Zello, G. A., & Chizen, D. R. (2020). A randomized controlled trial of a lifestyle intervention with longitudinal follow-up on ovarian dysmorphology in women with polycystic ovary syndrome. *Clinical Endocrinology*, 92(6), 525–535. <https://doi.org/10.1111/cen.14179>
- *Kehle-Forbes, S. M., Chen, S., Polusny, M. A., Lynch, K. G., Koffel, E., Ingram, E., Foa, E. B., Van Horn, D. H. A., Drapkin, M. L., Yusco, D. A., & Oslin, D. W. (2019). A randomized controlled trial evaluating integrated versus phased application of evidence-based psychotherapies for military veterans with comorbid PTSD and substance use disorders. *Drug and Alcohol Dependence*, 205, Article 107647. <https://doi.org/10.1016/j.drugalcdep.2019.107647>
- *Kelishadi, R., Sarrafzadegan, N., Sadri, G., Pashmi, R., Mohammadifard, N., Tavasoli, A., Amani, A., Rabiei, K., Khosravi, A., & Bahonar, A. (2011). Short-term results of a community-based program on promoting healthy lifestyle for prevention and control of chronic diseases in a developing country setting: Isfahan healthy heart program. *Asia-Pacific Journal of Public Health*, 23(4), 518–533. <https://doi.org/10.1177/1010539509348241>
- Keller, P. A., Harlam, B., Loewenstein, G., & Volpp, K. G. (2011). Enhanced active choice: A new method to motivate behavior change. *Journal of Consumer Psychology*, 21(4), 376–383. <https://doi.org/10.1016/j.jcps.2011.06.003>
- *Kempf, K., Altpeter, B., Berger, J., Reuß, O., Fuchs, M., Schneider, M., Gärtner, B., Niedermeier, K., & Martin, S. (2017). Efficacy of the telemedical lifestyle intervention program TeLiPro in advanced stages of type 2 diabetes: A randomized controlled trial. *Diabetes Care*, 40(7), 863–871. <https://doi.org/10.2337/dc17-0303>
- *Kershaw, T. S., Magriples, U., Westdahl, C., Rising, S. S., & Ickovics, J. (2009). Pregnancy as a window of opportunity for HIV prevention: Effects of an HIV intervention delivered within prenatal care. *American Journal of Public Health*, 99(11), 2079–2086. <https://doi.org/10.2105/AJPH.2008.154476>
- *Keyserling, T. C., Samuel Hodge, C. D., Jilcott, S. B., Johnston, L. F., Garcia, B. A., Gizlice, Z., Gross, M. D., Saviñon, C. E., Bangdiwala, S. I., Will, J. C., Farris, R. P., Trost, S., & Ammerman, A. S. (2008). Randomized trial of a clinic-based, community-supported, lifestyle intervention to improve physical activity and diet: The North Carolina enhanced WISEWOMAN project. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 46(6), 499–510. <https://doi.org/10.1016/j.ypmed.2008.02.011>
- *Keyserling, T. C., Samuel-Hodge, C. D., Ammerman, A. S., Ainsworth, B. E., Henríquez-Roldán, C. F., Elasy, T. A., Skelly, A. H., Johnston, L. F., & Bangdiwala, S. I. (2002). A randomized trial of an intervention to improve self-care behaviors of African-American women with type 2 diabetes: Impact on physical activity. *Diabetes Care*, 25(9), 1576–1583. <https://doi.org/10.2337/diacare.25.9.1576>
- *Kharmats, A. Y., Wang, C., Fuentes, L., Hu, L., Kline, T., Welding, K., & Cheskin, L. J. (2022). Monday-focused tailored rapid interactive mobile messaging for weight management 2 (MTRIMM2): Results from a randomized controlled trial. *mHealth*, 8, Article 1. <https://doi.org/10.21037/mhealth-21-3>
- Kiernan, M., Brown, S. D., Schoffman, D. E., Lee, K., King, A. C., Taylor, C. B., Schleicher, N. C., & Perri, M. G. (2013). Promoting healthy weight with “stability skills first”: A randomized trial. *Journal of Consulting and Clinical Psychology*, 81(2), 336–346. <https://doi.org/10.1037/a0030544>
- Kiesel, A., Steinhäuser, M., Wendt, M., Falkenstein, M., Jost, K., Philipp, A. M., & Koch, I. (2010). Control and interference in task switching—A review. *Psychological Bulletin*, 136(5), 849–874. <https://doi.org/10.1037/a0019842>
- *Killen, J. D., Robinson, T. N., Telch, M. J., Saylor, K. E., Maron, D. J., Rich, T., & Bryson, S. (1989). The Stanford adolescent heart health program. *Health Education Quarterly*, 16(2), 263–283. <https://doi.org/10.1177/109019818901600210>
- King, K., Meader, N., Wright, K., Graham, H., Power, C., Petticrew, M., White, M., & Sowden, A. J. (2015). Characteristics of interventions targeting multiple lifestyle risk behaviours in adult populations: A systematic scoping review. *PLOS ONE*, 10(1), Article e0117015. <https://doi.org/10.1371/journal.pone.0117015>
- *Kinnunen, T., Leeman, R. F., Korhonen, T., Quiles, Z. N., Terwal, D. M., Garvey, A. J., & Hartley, H. L. (2008). Exercise as an adjunct to nicotine gum in treating tobacco dependence among women. *Nicotine & Tobacco Research*, 10(4), 689–703. <https://doi.org/10.1080/14622200801979043>
- Kjelsberg, M. O., & the Multiple Risk Factor Intervention Trial Research Group. (1982). Multiple risk factor intervention trial. Risk factor changes and mortality results. *JAMA*, 248(12), 1465–1477. <https://doi.org/10.1001/jama.1982.03330120023025>
- Knäuper, B., Rabiau, M., Cohen, O., & Patriciu, N. (2004). Compensatory health beliefs: Scale development and psychometric properties. *Psychology & Health*, 19(5), 607–624. <https://doi.org/10.1080/0887044042000196737>
- *Knowlden, A. P., & Conrad, E. (2018). Two-year outcomes of the enabling mothers to prevent pediatric obesity through web-based education and reciprocal determinism (EMPOWER) randomized control trial. *Health Education & Behavior*, 45(2), 262–276. <https://doi.org/10.1177/1090198117732604>
- *Koblin, B., Chesney, M., Coates, T., & the EXPLORE Study Team. (2004). Effects of a behavioural intervention to reduce acquisition of HIV infection among men who have sex with men: The EXPLORE randomised controlled study. *The Lancet*, 364(9428), 41–50. [https://doi.org/10.1016/S0140-6736\(04\)16588-4](https://doi.org/10.1016/S0140-6736(04)16588-4)

- *Köse, S., & Yıldız, S. (2021). Motivational support programme to enhance health and well-being and promote weight loss in overweight and obese adolescents: A randomized controlled trial in Turkey. *International Journal of Nursing Practice*, 27(1), Article e12878. <https://doi.org/10.1111/ijn.12878>
- *Krasnoff, J. B., Vintro, A. Q., Ascher, N. L., Bass, N. M., Paul, S. M., Dodd, M. J., & Painter, P. L. (2006). A randomized trial of exercise and dietary counseling after liver transplantation. *American Journal of Transplantation*, 6(8), 1896–1905. <https://doi.org/10.1111/j.1600-6143.2006.01391.x>
- Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16(9), 606–613. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x>
- Kypri, K., & McAnally, H. M. (2005). Randomized controlled trial of a web-based primary care intervention for multiple health risk behaviors. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 41(3–4), 761–766. <https://doi.org/10.1016/j.ypmed.2005.07.010>
- *La Torre, G., Cocchiara, R. A., Sordo, E. L., Chiarini, M., Siliquini, R., Firenze, A., Maurici, M., Agati, L., Saulle, R., & Mannocci, A. (2018). Counseling intervention to improve quality of life in patients with pre-existing acute myocardial infarction (AMI) or chronic obstructive pulmonary disease (COPD): A pilot study. *Journal of Preventive Medicine and Hygiene*, 59(2), E153–E158.
- LaRowe, L. R., Dunsiger, S. I., & Williams, D. M. (2023). Acute exercise-induced changes in motivation and behavioral expectation for quitting smoking as predictors of smoking behavior in women. *Psychology of Addictive Behaviors*, 37(3), 475–482. <https://doi.org/10.1037/adb0000901>
- *Latkin, C. A., Donnell, D., Metzger, D., Sherman, S., Aramrattana, A., Davis-Vogel, A., Quan, V. M., Gandham, S., Vongchak, T., Perdue, T., & Celentano, D. D. (2009). The efficacy of a network intervention to reduce HIV risk behaviors among drug users and risk partners in Chiang Mai, Thailand and Philadelphia, USA. *Social Science & Medicine*, 68(4), 740–748. <https://doi.org/10.1016/j.socscimed.2008.11.019>
- *Leahey, T. M., Huedo-Medina, T. B., Grenga, A., Gay, L., Fernandes, D., Denmat, Z., Doyle, C., Areny-Joval, R., & Wing, R. R. (2020). Patient-provided e-support in reduced intensity obesity treatment: The INSPIRE randomized controlled trial. *Health Psychology*, 39(12), 1037–1047. <https://doi.org/10.1037/hea0000996>
- *Lear, S. A., Spinelli, J. J., Linden, W., Brozic, A., Kiess, M., Frohlich, J. J., & Ignaszewski, A. (2006). The Extensive Lifestyle Management Intervention (ELMI) after cardiac rehabilitation: A 4-year randomized controlled trial. *American Heart Journal*, 152(2), 333–339. <https://doi.org/10.1016/j.ahj.2005.12.023>
- *Lee, D. C., Walker, D. D., Hughes, J. R., Brunette, M. F., Scherer, E., Stanger, C., Etter, J. F., Auty, S., & Budney, A. J. (2019). Sequential and simultaneous treatment approaches to cannabis use disorder and tobacco use. *Journal of Substance Abuse Treatment*, 98, 39–46. <https://doi.org/10.1016/j.jsat.2018.12.005>
- Lee, J. S., Keil, M., & Wong, K. F. E. (2015). The effect of goal difficulty on escalation of commitment. *Journal of Behavioral Decision Making*, 28(2), 114–129. <https://doi.org/10.1002/bdm.1835>
- *Leighton, R. F., Repka, F. J., Birk, T. J., Lynch, D. J., Bingle, J. F., Gohara, A. F., Saffran, J., Weaver, M. T., Brewster, P. M., & Walsh, M. E. (1990). The Toledo exercise and diet study. Results at 26 weeks. *Archives of Internal Medicine*, 150(5), 1016–1020. <https://doi.org/10.1001/archinte.1990.00390170066015>
- *Letourneau, E. J., McCart, M. R., Sheidow, A. J., & Mauro, P. M. (2017). First evaluation of a contingency management intervention addressing adolescent substance use and sexual risk behaviors: Risk reduction therapy for adolescents. *Journal of Substance Abuse Treatment*, 72, 56–65. <https://doi.org/10.1016/j.jsat.2016.08.019>
- *Lewis, C. F., Rivera, A. V., Crawford, N. D., DeCuir, J., & Amesty, S. (2015). Pharmacy-randomized intervention delivering HIV prevention services during the syringe sale to people who inject drugs in New York City. *Drug and Alcohol Dependence*, 153, 72–77. <https://doi.org/10.1016/j.drugalcdep.2015.06.006>
- Light, R. J., & Pillemer, D. B. (1984). *Summing up: The science of reviewing research*. Harvard University Press. <https://doi.org/10.4159/9780674040243>
- *Lillis, J., Dunsiger, S., Thomas, J. G., Ross, K. M., & Wing, R. R. (2021). Novel behavioral interventions to improve long-term weight loss: A randomized trial of acceptance and commitment therapy or self-regulation for weight loss maintenance. *Journal of Behavioral Medicine*, 44(4), 527–540. <https://doi.org/10.1007/s10865-021-00215-z>
- *Lin, H. H., Tsai, Y. F., Lin, P. J., & Tsay, P. K. (2010). Effects of a therapeutic lifestyle-change programme on cardiac risk factors after coronary artery bypass graft. *Journal of Clinical Nursing*, 19(1–2), 60–68. <https://doi.org/10.1111/j.1365-2702.2009.02980.x>
- *Lindström, J., Eriksson, J. G., Valle, T. T., Aunola, S., Cepaitis, Z., Hakumäki, M., Hämäläinen, H., Ilanne-Parikka, P., Keinänen-Kiukaanniemi, S., Laakso, M., Louheranta, A., Mannelin, M., Martikkala, V., Moltchanov, V., Rastas, M., Salminen, V., Sundvall, J., Uusitupa, M., & Tuomilehto, J. (2003). Prevention of diabetes mellitus in subjects with impaired glucose tolerance in the Finnish Diabetes Prevention Study: Results from a randomized clinical trial. *Journal of the American Society of Nephrology*, 14(Suppl_2), S108–S113. <https://doi.org/10.1097/01.ASN.0000070157.96264.13>
- *Lindström, J., Louheranta, A., Mannelin, M., Rastas, M., Salminen, V., Eriksson, J., Uusitupa, M., Tuomilehto, J., & the Finnish Diabetes Prevention Study Group. (2003). The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care*, 26(12), 3230–3236. <https://doi.org/10.2337/diaca.re.26.12.3230>
- Lippke, S., Nigg, C. R., & Maddock, J. E. (2012). Health-promoting and health-risk behaviors: Theory-driven analyses of multiple health behavior change in three international samples. *International Journal of Behavioral Medicine*, 19(1), 1–13. <https://doi.org/10.1007/s12529-010-9135-4>
- *Lloyd-Richardson, E. E., Stanton, C. A., Papandonatos, G. D., Shadel, W. G., Stein, M., Tashima, K., Flanagan, T., Morrow, K., Neighbors, C., & Niaura, R. (2009). Motivation and patch treatment for HIV+ smokers: A randomized controlled trial. *Addiction*, 104(11), 1891–1900. <https://doi.org/10.1111/j.1360-0443.2009.02623.x>
- *López-Padrós, C., Salord, N., Alves, C., Vilarrasa, N., Gasa, M., Planas, R., Montserrat, M., Virgili, M. N., Rodríguez, C., Pérez-Ramos, S., López-Cadena, E., Ramos, M. I., Dorca, J., & Monasterio, C. (2020). Effectiveness of an intensive weight-loss program for severe OSA in patients undergoing CPAP treatment: A randomized controlled trial. *Journal of Clinical Sleep Medicine*, 16(4), 503–514. <https://doi.org/10.5664/jcsm.8252>
- *Lovell, K., Wearden, A., Bradshaw, T., Tomenson, B., Pedley, R., Davies, L. M., Husain, N., Woodham, A., Escott, D., Swarbrick, C. M., Femi-Ajao, O., Warburton, J., & Marshall, M. (2014). An exploratory randomized controlled study of a healthy living intervention in early intervention services for psychosis: The INTERvention to encourage ACTivity, improve diet, and reduce weight gain (INTERACT) study. *The Journal of Clinical Psychiatry*, 75(5), 498–505. <https://doi.org/10.4088/JCP.13m08503>
- *Lugada, E., Levin, J., Abang, B., Mermin, J., Mugalanzi, E., Namara, G., Gupta, S., Grosskurth, H., Jaffar, S., Coutinho, A., & Bunnell, R. (2010). Comparison of home and clinic-based HIV testing among household members of persons taking antiretroviral therapy in Uganda: Results from a randomized trial. *Journal of Acquired Immune Deficiency Syndromes*, 55(2), 245–252. <https://doi.org/10.1097/QAI.0b013e3181e9e069>
- *Lutalo, T., Kigozi, G., Kimera, E., Serwadda, D., Wawer, M. J., Zabin, L. S., & Gray, R. H. (2010). A randomized community trial of enhanced family planning outreach in Rakai, Uganda. *Studies in Family Planning*, 41(1), 55–60. <https://doi.org/10.1111/j.1728-4465.2010.00224.x>
- *Lv, J., Liu, Q. M., Ren, Y. J., He, P. P., Wang, S. F., Gao, F., Li, L. M., & the Community Interventions for Health (CIH) Collaboration. (2014). A community-based multilevel intervention for smoking, physical activity

- and diet: Short-term findings from the Community Interventions for Health programme in Hangzhou, China. *Journal of Epidemiology and Community Health*, 68(4), 333–339. <https://doi.org/10.1136/jech-2013-203356>
- *Machado, M. P., Fidalgo, T. M., Brasiliano, S., Hochgraf, P. B., & Noto, A. R. (2020). The contribution of mindfulness to outpatient substance use disorder treatment in Brazil: A preliminary study. *Brazilian Journal of Psychiatry*, 42(5), 527–531. <https://doi.org/10.1590/1516-4446-2019-0725>
- *Maddison, R., Roberts, V., McRobbie, H., Bullen, C., Prapavessis, H., Glover, M., Jiang, Y., Brown, P., Leung, W., Taylor, S., & Tsai, M. (2014). Exercise counseling to enhance smoking cessation outcomes: The Fit2Quit randomized controlled trial. *Annals of Behavioral Medicine: A Publication of the Society of Behavioral Medicine*, 48(2), 194–204. <https://doi.org/10.1007/s12160-014-9588-9>
- *Manios, Y., Moschonis, G., Koutsikas, K., Papoutsou, S., Petraki, I., Bellou, E., Naoumi, A., Kostea, S., & Tanagra, S. (2009). Changes in body composition following a dietary and lifestyle intervention trial: The postmenopausal health study. *Maturitas*, 62(1), 58–65. <https://doi.org/10.1016/j.maturitas.2008.11.005>
- Marcolina, S. T. (2007). The effect of exercise on depression. *Alternative Therapies in Womens Health*, 9(8), 57–62.
- *Marcus, B. H., Albrecht, A. E., Niaura, R. S., Taylor, E. R., Simkin, L. R., Feder, S. I., Abrams, D. B., & Thompson, P. D. (1995). Exercise enhances the maintenance of smoking cessation in women. *Addictive Behaviors*, 20(1), 87–92. [https://doi.org/10.1016/0306-4603\(94\)00048-4](https://doi.org/10.1016/0306-4603(94)00048-4)
- *Marsch, L. A., Guarino, H., Grabinski, M. J., Syckes, C., Dillingham, E. T., Xie, H., & Crosier, B. S. (2015). Comparative effectiveness of web-based vs. educator-delivered HIV prevention for adolescent substance users: A randomized, controlled trial. *Journal of Substance Abuse Treatment*, 59, 30–37. <https://doi.org/10.1016/j.jsat.2015.07.003>
- *Marsiglia, F. F., Ayers, S. L., Han, S., & Weide, A. (2019). The role of culture of origin on the effectiveness of a parents-involved intervention to prevent substance use among Latino middle school youth: Results of a cluster randomized controlled trial. *Prevention Science*, 20(5), 643–654. <https://doi.org/10.1007/s11121-018-0968-4>
- Martin, C. K., Johnson, W. D., Myers, C. A., Apolzan, J. W., Earnest, C. P., Thomas, D. M., Rood, J. C., Johannsen, N. M., Tudor-Locke, C., Harris, M., Hsia, D. S., & Church, T. S. (2019). Effect of different doses of supervised exercise on food intake, metabolism, and non-exercise physical activity: The E-MECHANIC randomized controlled trial. *The American Journal of Clinical Nutrition*, 110(3), 583–592. <https://doi.org/10.1093/ajcn/nqz054>
- *Martin, S. S., O'Connell, D. J., Inciardi, J. A., Surratt, H. L., & Beard, R. A. (2003). HIV/AIDS among probationers: An assessment of risk and results from a brief intervention. *Journal of Psychoactive Drugs*, 35(4), 435–443. <https://doi.org/10.1080/02791072.2003.10400490>
- Mata, J., Silva, M. N., Vieira, P. N., Carraça, E. V., Andrade, A. M., Coutinho, S. R., Sardinha, L. B., & Teixeira, P. J. (2009). Motivational “spill-over” during weight control: Increased self-determination and exercise intrinsic motivation predict eating self-regulation. *Health Psychology*, 28(6), 709–716. <https://doi.org/10.1037/a0016764>
- *Mausbach, B. T., Semple, S. J., Strathdee, S. A., Zians, J., & Patterson, T. L. (2007). Efficacy of a behavioral intervention for increasing safer sex behaviors in HIV-negative, heterosexual methamphetamine users: Results from the Fast-Lane Study. *Annals of Behavioral Medicine: A Publication of the Society of Behavioral Medicine*, 34(3), 263–274. <https://doi.org/10.1007/BF02874551>
- McCaffrey, D. F., & Bell, R. M. (2003). Bias reduction in standard errors for linear and generalized linear models with multi-stage samples. *Proceedings of Statistics Canada Symposium*, 28(12), 1–10.
- McCloskey, K., & Johnson, B. T. (2019). Habits, quick and easy: Perceived complexity moderates the associations of contextual stability and rewards with behavioral automaticity. *Frontiers in Psychology*, 10, Article 1556. <https://doi.org/10.3389/fpsyg.2019.01556>
- *McCusker, J., Bigelow, C., Frost, R., Garfield, F., Hindin, R., Vickers-Lahti, M., & Lewis, B. (1997). The effects of planned duration of residential drug abuse treatment on recovery and HIV risk behavior. *American Journal of Public Health*, 87(10), 1637–1644. <https://doi.org/10.2105/AJPH.87.10.1637>
- McDonald, J., McDonald, P., Hughes, C., & Albarracín, D. (2017). Recalling and intending to enact health recommendations: Optimal number of prescribed behaviors in multibehavior messages. *Clinical Psychological Science*, 5(5), 858–865. <https://doi.org/10.1177/2167702617704453>
- McEachan, R. R. C., Conner, M., Taylor, N. J., & Lawton, R. J. (2011). Prospective prediction of health-related behaviours with the theory of planned behaviour: A meta-analysis. *Health Psychology Review*, 5(2), 97–144. <https://doi.org/10.1080/17437199.2010.521684>
- *McGillion, M. H., Watt-Watson, J., Stevens, B., Lefort, S. M., Coyte, P., & Graham, A. (2008). Randomized controlled trial of a psychoeducation program for the self-management of chronic cardiac pain. *Journal of Pain and Symptom Management*, 36(2), 126–140. <https://doi.org/10.1016/j.jpainnsymman.2007.09.015>
- McHorney, C. A., Ware, J. E., & Raczek, A. E. (1993). The MOS 36-item short-form health survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Medical Care*, 31(3), 247–263. <https://doi.org/10.1097/00005650-199303000-00006>
- McLellan, A. T., Kushner, H., Metzger, D., Peters, R., Smith, I., Grissom, G., Pettinati, H., & Argeriou, M. (1992). The fifth edition of the Addiction Severity Index. *Journal of Substance Abuse Treatment*, 9(3), 199–213. [https://doi.org/10.1016/0740-5472\(92\)90062-S](https://doi.org/10.1016/0740-5472(92)90062-S)
- McNair, D. M., Lorr, M., & Droppleman, L. F. (1971). *EITS Manual for the Profile of Mood States* (Issue 27). Educational and Industrial Testing Service.
- *Meacham, M. C., Ramo, D. E., Prochaska, J. J., Maier, L. J., Delucchi, K. L., Kaur, M., & Satre, D. D. (2021). A Facebook intervention to address cigarette smoking and heavy episodic drinking: A pilot randomized controlled trial. *Journal of Substance Abuse Treatment*, 122, Article 108211. <https://doi.org/10.1016/j.jsat.2020.108211>
- *Meade, C. S., Drabkin, A. S., Hansen, N. B., Wilson, P. A., Kochman, A., & Sikkema, K. J. (2010). Reductions in alcohol and cocaine use following a group coping intervention for HIV-positive adults with childhood sexual abuse histories. *Addiction*, 105(11), 1942–1951. <https://doi.org/10.1111/j.1360-0443.2010.03075.x>
- Meader, N., King, K., Wright, K., Graham, H. M., Petticrew, M., Power, C., White, M., & Sowden, A. J. (2017). Multiple risk behavior interventions: Meta-analyses of RCTs. *American Journal of Preventive Medicine*, 53(1), e19–e30. <https://doi.org/10.1016/j.amepre.2017.01.032>
- *Melanson, K. J., Dell'Olio, J., Carpenter, M. R., & Angelopoulos, T. J. (2004). Changes in multiple health outcomes at 12 and 24 weeks resulting from 12 weeks of exercise counseling with or without dietary counseling in obese adults. *Nutrition*, 20(10), 849–856. <https://doi.org/10.1016/j.nut.2004.06.004>
- *Melnik, B. M., Jacobson, D., Kelly, S., Belyea, M., Shaibi, G., Small, L., O'Haver, J., & Marsiglia, F. F. (2013). Promoting healthy lifestyles in high school adolescents: A randomized controlled trial. *American Journal of Preventive Medicine*, 45(4), 407–415. <https://doi.org/10.1016/j.amepre.2013.05.013>
- Melsen, W. G., Bootsma, M. C. J., Rovers, M. M., & Bonten, M. J. M. (2014). The effects of clinical and statistical heterogeneity on the predictive values of results from meta-analyses. *Clinical Microbiology and Infection*, 20(2), 123–129. <https://doi.org/10.1111/1469-0691.12494>
- *Mendez-Ruiz, M. D., Villegas-Pantoja, M. A., Alarcón-Luna, N. S., Villegas, N., Cianelli, R., & Peragallo-Montano, N. (2020). Prevention of alcohol consumption and transmission of human immunodeficiency virus: Randomized clinical trial. *Revista Latino-Americana de Enfermagem*, 28, Article e3262. <https://doi.org/10.1590/1518-8345.3393.3262>

- *Menotti, A., Farchi, G., Buongiorno, A. M., Capocaccia, R., Conti, S., Di Carlo, G. G., Dima, F., Giampaoli, S., Mariotti, S., Morisi, G., Seccareccia, F., Verdecchia, A., Ricci, G., Urbinati, G. C., Angelico, F., Antonini, R., Bucci, A., Clemente, P., Conti, R., ... the Research Group of the Rome Project of Coronary Heart Disease Prevention. (1986). Eight-Year follow-up results from the Rome Project of Coronary Heart Disease Prevention. *Preventive Medicine*, 15(2), 176–191. [https://doi.org/10.1016/0091-7435\(86\)90087-3](https://doi.org/10.1016/0091-7435(86)90087-3)
- *Miettinen, T. A., Huttunen, J. K., Naukkarinen, V., Strandberg, T., Mattila, S., Kumlin, T., & Sarna, S. (1985). Multifactorial primary prevention of cardiovascular diseases in middle-aged men: Risk factor changes, incidence, and mortality. *Journal of the American Medical Association*, 254(15), 2097–2102. <https://doi.org/10.1001/jama.1985.03360150073027>
- *Miller, M. K., Catley, D., Adams, A., Staggs, V. S., Dowd, M. D., Stancil, S. L., Miller, E., Satterwhite, C. L., Bauermeister, J., & Goggin, K. (2021). Brief motivational intervention to improve adolescent sexual health service uptake: A pilot randomized controlled trial in the emergency department. *The Journal of Pediatrics*, 237, 250–257.e2. <https://doi.org/10.1016/j.jpeds.2021.06.007>
- *Mimiaga, M. J., Pantalone, D. W., Biello, K. B., Hughto, J. M. W., Frank, J., O'Cleirigh, C., Reisner, S. L., Restar, A., Mayer, K. H., & Safren, S. A. (2019). An initial randomized controlled trial of behavioral activation for treatment of concurrent crystal methamphetamine dependence and sexual risk for HIV acquisition among men who have sex with men. *AIDS Care*, 31(9), 1083–1095. <https://doi.org/10.1080/09540121.2019.1595518>
- *Mittag, O., China, C., Hoberg, E., Juers, E., Kolenda, K. D., Richardt, G., Maurischat, C., & Raspe, H. (2006). Outcomes of cardiac rehabilitation with versus without a follow-up intervention rendered by telephone (Luebeck follow-up trial): Overall and gender-specific effects. *International Journal of Rehabilitation Research. Internationale Zeitschrift Fur Rehabilitationsforschung. Revue Internationale de Recherches de Readaptation*, 29(4), 295–302. <https://doi.org/10.1097/MRR.0b013e328010ba9a>
- Mokdad, A. H., Ballesteros, K., Echko, M., Glenn, S., Olsen, H. E., Mullany, E., Lee, A., Khan, A. R., Ahmadi, A., Ferrari, A. J., Kasaiean, A., Werdecker, A., Carter, A., Zipkin, B., Sartorius, B., Serdar, B., Sykes, B. L., Troeger, C., Fitzmaurice, C., ... the US Burden of Disease Collaborators. (2018). The state of US health, 1990–2016: Burden of diseases, injuries, and risk factors among US states. *JAMA: Journal of the American Medical Association*, 319(14), 1444–1472. <https://doi.org/10.1001/jama.2018.0158>
- Montgomery, S. A., & Asberg, M. (1979). A new depression scale designed to be sensitive to change. *The British Journal of Psychiatry*, 134(4), 382–389. <https://doi.org/10.1192/bjp.134.4.382>
- *Monti, P. M., Mastroleo, N. R., Barnett, N. P., Colby, S. M., Kahler, C. W., & Operario, D. (2016). Brief motivational intervention to reduce alcohol and HIV/sexual risk behavior in emergency department patients: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*, 84(7), 580–591. <https://doi.org/10.1037/ccp0000097>
- *Morais, L. D. C., Paravidino, V. B., Mediano, M. F. F., Benthroldo, R., Gonçalves, T. R., Sgambato, M. R., de Souza, B. D. S. N., Junior, E. V., Marques, E. S., Pereira, R. A., Sichieri, R., & Cunha, D. B. (2021). Effectiveness of a school-based randomized controlled trial aimed at increasing physical activity time in adolescents. *European Journal of Public Health*, 31(2), 367–372. <https://doi.org/10.1093/eupub/ckab025>
- *Moravcová, K., Karbanová, M., Bretschneider, M. P., Sovová, M., Ožana, J., & Sovová, E. (2022). Comparing digital therapeutic intervention with an intensive obesity management program: Randomized controlled trial. *Nutrients*, 14(10), Article 2005. <https://doi.org/10.3390/nu14102005>
- *Morey, M. C., Snyder, D. C., Sloane, R., Cohen, H. J., Peterson, B., Hartman, T. J., Miller, P., Mitchell, D. C., & Demark-Wahnefried, W. (2009). Effects of home-based diet and exercise on functional outcomes among older, overweight long-term cancer survivors: RENEW: A randomized controlled trial. *JAMA*, 301(18), 1883–1891. <https://doi.org/10.1001/jama.2009.643>
- Morris, S. B. (2008). Estimating effect sizes from pretest–posttest–control group designs. *Organizational Research Methods*, 11(2), 364–386. <https://doi.org/10.1177/1094428106291059>
- Mouttapa, M., Watson, D. W., McCuller, W. J., Reiber, C., & Tsai, W. (2009). Substance use and HIV prevention for youth in correctional facilities. *Journal of Correctional Education*, 60(4), 289–315.
- *Murphy, D. A., Chen, X., Naar-King, S., Parsons, J. T., & the Adolescent Trials Network. (2012). Alcohol and marijuana use outcomes in the Healthy Choices motivational interviewing intervention for HIV-positive youth. *AIDS Patient Care and STDs*, 26(2), 95–100. <https://doi.org/10.1089/apc.2011.0157>
- Murray, C. J., Aravkin, A. Y., Zheng, P., Abbafati, C., Abbas, K. M., Abbasi-Kangevari, M., Abd-Allah, F., Abdelalim, A., Abdollahi, M., Abdollahpour, I., Abegaz, K. H., Abolhassani, H., Aboyans, V., Abreu, L. G., Abrigo, M. R. M., Abualhasan, A., Abu-Raddad, L. J., Abushouk, A. I., Adabi, M., ... the GBD 2019 Risk Factors Collaborators. (2020). Global burden of 87 risk factors in 204 countries and territories, 1990–2019: A systematic analysis for the Global Burden of Disease Study 2019. *Lancet*, 396(10258), 1223–1249. [https://doi.org/10.1016/S0140-6736\(20\)30752-2](https://doi.org/10.1016/S0140-6736(20)30752-2)
- Murthy, V. H. (2017). Surgeon general's report on alcohol, drugs, and health. *JAMA*, 317(2), 133–134. <https://doi.org/10.1001/jama.2016.18215>
- *Muto, T., & Yamauchi, K. (2001). Evaluation of a multicomponent workplace health promotion program conducted in Japan for improving employees' cardiovascular disease risk factors. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 33(6), 571–577. <https://doi.org/10.1006/pmed.2001.0923>
- *Nader, P. R., Stone, E. J., Lytle, L. A., Perry, C. L., Osganian, S. K., Kelder, S., Webber, L. S., Elder, J. P., Montgomery, D., Feldman, H. A., Wu, M., Johnson, C., Parcel, G. S., & Luepker, R. V. (1999). Three-year maintenance of improved diet and physical activity: The CATCH cohort. Child and Adolescent Trial for Cardiovascular Health. *Archives of Pediatrics & Adolescent Medicine*, 153(7), 695–704. <https://doi.org/10.1001/archpedi.153.7.695>
- Ngandu, T., Lehtisalo, J., Solomon, A., Levälähti, E., Ahtiluoto, S., Antikainen, R., Bäckman, L., Hänninen, T., Jula, A., Laatikainen, T., Lindström, J., Mangialasche, F., Pajananen, T., Pajala, S., Peltonen, M., Rauramaa, R., Stigsdotter-Neely, A., Strandberg, T., Tuomilehto, J., ... Kivipelto, M. (2015). A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): A randomised controlled trial. *The Lancet*, 385(9984), 2255–2263. [https://doi.org/10.1016/S0140-6736\(15\)60461-5](https://doi.org/10.1016/S0140-6736(15)60461-5)
- *Niaura, R., Marcus, B., Albrecht, A., Thompson, P., & Abrams, D. (1998). Exercise, smoking cessation, and short-term changes in serum lipids in women: A preliminary investigation. *Medicine and Science in Sports and Exercise*, 30(9), 1414–1418. <https://doi.org/10.1097/0000-5768-199809000-00010>
- *Nickel, N. C., Doupe, M., Enns, J. E., Brownell, M., Sarkar, J., Chateau, D., Burland, E., Chartier, M., Katz, A., Crockett, L., Azad, M. B., McGavock, J. M., & Santos, R. (2021). Differential effects of a school-based obesity prevention program: A cluster randomized trial. *Maternal and Child Nutrition*, 17(1), Article e13009. <https://doi.org/10.1111/mcn.13009>
- Nigg, C., Amrein, M., Rackow, P., Scholz, U., & Inauen, J. (2021). Compensation and transfer effects of eating behavior change in daily life: Evidence from a randomized controlled trial. *Appetite*, 162, Article 105170. <https://doi.org/10.1016/j.appet.2021.105170>
- Nigg, C. R., Lee, H. R., Hubbard, A. E., & Min-Sun, K. (2009). Gateway health behaviors in college students: Investigating transfer and compensation effects. *Journal of American College Health*, 58(1), 39–44. <https://doi.org/10.3200/JACH.58.1.39-44>

- Nigg, C. R., & Long, C. R. (2012). A systematic review of single health behavior change interventions vs. multiple health behavior change interventions among older adults. *Translational Behavioral Medicine*, 2(2), 163–179. <https://doi.org/10.1007/s13142-012-0130-y>
- *Nilsson, P. M., Klasson, E. B., & Nyberg, P. (2001). Life-style intervention at the worksite—Reduction of cardiovascular risk factors in a randomized study. *Scandinavian Journal of Work, Environment & Health*, 27(1), 57–62. <https://doi.org/10.5271/sjweh.587>
- *Njuguna, N., Ngure, K., Mugo, N., Sambu, C., Siano, C., Gakuo, S., Irungu, E., Baeten, J., & Heffron, R. (2016). The effect of human immunodeficiency virus prevention and reproductive health text messages on human immunodeficiency virus testing among young women in rural Kenya: A pilot study. *Sexually Transmitted Diseases*, 43(6), 353–359. <https://doi.org/10.1097/OLQ.0000000000000450>
- Noar, S. M., Chabot, M., & Zimmerman, R. S. (2008). Applying health behavior theory to multiple behavior change: Considerations and approaches. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 46(3), 275–280. <https://doi.org/10.1016/j.ypmed.2007.08.001>
- Nooijen, C. F. J., Del Pozo-Cruz, B., Nyberg, G., Sanders, T., Galanti, M. R., & Forsell, Y. (2018). Are changes in occupational physical activity level compensated by changes in exercise behavior? *European Journal of Public Health*, 28(5), 940–943. <https://doi.org/10.1093/eurpub/cky007>
- *Nöstlinger, C., Platteau, T., Bogner, J., Buyze, J., Dec-Pietrowska, J., Dias, S., Newbury-Helps, J., Kocsis, A., Mueller, M., Rojas, D., Stanekova, D., van Lankveld, J., Colebunders, R., & the Eurosupport Study Group. (2016). Implementation and operational research: Computer-assisted intervention for safer sex in HIV-positive men having sex with men: Findings of a European randomized multi-center trial. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 71(3), e63–e72. <https://doi.org/10.1097/QAI.0000000000000882>
- Odutayo, A., Emdin, C. A., Hsiao, A. J., Shakir, M., Copsey, B., Dutton, S., Chiochia, V., Schluskel, M., Dutton, P., Roberts, C., Altman, D. G., & Hopewell, S. (2017). Association between trial registration and positive study findings: Cross sectional study (Epidemiological Study of Randomized Trials-ESORT). *The BMJ*, 356, Article j917. <https://doi.org/10.1136/bmj.j917>
- *Oenema, A., Brug, J., Dijkstra, A., de Weerd, I., & de Vries, H. (2008). Efficacy and use of an internet-delivered computer-tailored lifestyle intervention, targeting saturated fat intake, physical activity and smoking cessation: A randomized controlled trial. *Annals of Behavioral Medicine*, 35(2), 125–135. <https://doi.org/10.1007/s12160-008-9023-1>
- Office of Disease Prevention and Health Promotion. (2018). *Behavioral risk factor surveillance survey*. <https://www.healthypeople.gov/2020/data-source/behavioral-risk-factor-surveillance-system>
- *Ögmundsdóttir Michelsen, H., Sjölin, I., Bäck, M., Gonzalez Garcia, M., Olsson, A., Sandberg, C., Schiopu, A., & Leósdóttir, M. (2022). Effect of a lifestyle-focused web-based application on risk factor management in patients who have had a myocardial infarction: Randomized controlled trial. *Journal of Medical Internet Research*, 24(3), Article e25224. <https://doi.org/10.2196/25224>
- *Oh, E. G., Hyun, S. S., Kim, S. H., Bang, S. Y., Chu, S. H., Jeon, J. Y., & Kang, M. S. (2008). A randomized controlled trial of therapeutic lifestyle modification in rural women with metabolic syndrome: A pilot study. *Metabolism: Clinical and Experimental*, 57(2), 255–261. <https://doi.org/10.1016/j.metabol.2007.09.009>
- *Olvera, N., Bush, J. A., Sharma, S., Knox, B., Scherer, R., & Butte, N. (2010). BOUNCE: A community-based mother–daughter healthy lifestyle intervention for low-income Latino families. *Obesity*, 18(S1), S102–S104. <https://doi.org/10.1038/oby.2009.439>
- *Ornish, D., Brown, S. E., Billings, J. H., Scherwitz, L. W., Armstrong, W. T., Ports, T. A., McLanahan, S. M., Kirkeeide, R. L., Brand, R. J., & Gould, K. L. (1990). Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet*, 336(8708), 129–133. [https://doi.org/10.1016/0140-6736\(90\)91656-U](https://doi.org/10.1016/0140-6736(90)91656-U)
- *Otterstad, J. E., & the Vestfold Heartcare Study Group. (2003). Influence on lifestyle measures and five-year coronary risk by a comprehensive lifestyle intervention programme in patients with coronary heart disease. *European Journal of Cardiovascular Prevention and Rehabilitation*, 10(6), 429–437. <https://doi.org/10.1097/01.hjr.0000107024.38316.6a>
- *Owolabi, E. O., Goon, D. T., & Ajayi, A. I. (2020). Impact of mobile phone text messaging intervention on adherence among patients with diabetes in a rural setting: A randomized controlled trial. *Medicine*, 99(12), Article e18953. <https://doi.org/10.1097/MD.00000000000018953>
- Özerbil, O., Okudan, N., Gökbil, H., & Levendoğlu, F. (2006). Comparison of the effects of two antidepressants on exercise performance of the female patients with fibromyalgia. *Clinical Rheumatology*, 25(4), 495–497. <https://doi.org/10.1007/s10067-005-0076-2>
- Page, M. J., Higgins, J. P., Clayton, G., Sterne, J. A., Hróbjartsson, A., & Savović, J. (2016). Empirical evidence of study design biases in randomized trials: Systematic review of meta-epidemiological studies. *PLOS ONE*, 11(7), Article e0159267. <https://doi.org/10.1371/journal.pone.0159267>
- *Palavras, M. A., Hay, P., Mannan, H., da Luz, F. Q., Sainsbury, A., Touyz, S., & Claudino, A. M. (2021). Integrated weight loss and cognitive behavioural therapy (CBT) for the treatment of recurrent binge eating and high body mass index: A randomized controlled trial. *Eating and Weight Disorders*, 26(1), 249–262. <https://doi.org/10.1007/s40519-020-00846-2>
- *Parcesepe, A. M., L Engle, K. L., Martin, S. L., Green, S., Sinkele, W., Suchindran, C., Speizer, I. S., Mwarogo, P., & Kingola, N. (2016). The impact of an alcohol harm reduction intervention on interpersonal violence and engagement in sex work among female sex workers in Mombasa, Kenya: Results from a randomized controlled trial. *Drug and Alcohol Dependence*, 161, 21–28. <https://doi.org/10.1016/j.drugalcdep.2015.12.037>
- *Parsons, J. T., Golub, S. A., Rosof, E., & Holder, C. (2007). Motivational interviewing and cognitive-behavioral intervention to improve HIV medication adherence among hazardous drinkers: A randomized controlled trial. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 46(4), 443–450. <https://doi.org/10.1097/QAI.0b013e318158a461>
- *Parsons, J. T., Lelutiu-Weinberger, C., Botsko, M., & Golub, S. A. (2014). A randomized controlled trial utilizing motivational interviewing to reduce HIV risk and drug use in young gay and bisexual men. *Journal of Consulting and Clinical Psychology*, 82(1), 9–18. <https://doi.org/10.1037/a0035311>
- Parsons, T. J., Power, C., & Manor, O. (2005). Physical activity, television viewing and body mass index: A cross-sectional analysis from childhood to adulthood in the 1958 British cohort. *International Journal of Obesity*, 29(10), 1212–1221. <https://doi.org/10.1038/sj.ijo.0802932>
- *Patel, R. M., Misra, R., Raj, S., & Balasubramanyam, A. (2017). Effectiveness of a group-based culturally tailored lifestyle intervention program on changes in risk factors for type 2 diabetes among Asian Indians in the United States. *Journal of Diabetes Research*, 2017, Article 2751980. <https://doi.org/10.1155/2017/2751980>
- *Paul, L., van der Heiden, C., van Hoeken, D., Deen, M., Vlijm, A., Klaassen, R. A., Biter, L. U., & Hoek, H. W. (2021). Cognitive behavioral therapy versus usual care before bariatric surgery: One-year follow-up results of a randomized controlled trial. *Obesity Surgery*, 31(3), 970–979. <https://doi.org/10.1007/s11695-020-05081-3>
- Pavlov, I. P. (1927). *Conditioned reflexes: An investigation of the physiological activity of the cerebral cortex*. Oxford University Press.
- *Pedley, C. F., Case, L. D., Blackwell, C. S., Katula, J. A., & Vitolins, M. Z. (2018). The 24-month metabolic benefits of the healthy living partnerships to prevent diabetes: A community-based translational study. *Diabetes & Metabolic Syndrome*, 12(3), 215–220. <https://doi.org/10.1016/j.dsx.2017.09.011>

- *Peltzer, K., Simbayi, L., Banyini, M., & Kekana, Q. (2012). HIV risk reduction intervention among medically circumcised young men in South Africa: A randomized controlled trial. *International Journal of Behavioral Medicine*, 19(3), 336–341. <https://doi.org/10.1007/s12529-011-9171-8>
- *Penberthy, J. K., Hook, J. N., Hettema, J., Farrell-Carnahan, L., & Ingersoll, K. (2013). Depressive symptoms moderate treatment response to brief intervention for prevention of alcohol exposed pregnancy. *Journal of Substance Abuse Treatment*, 45(4), 335–342. <https://doi.org/10.1016/j.jsat.2013.05.002>
- *Peragallo, N., Gonzalez-Guarda, R. M., McCabe, B. E., & Cianelli, R. (2012). The efficacy of an HIV risk reduction intervention for Hispanic women. *AIDS and Behavior*, 16(5), 1316–1326. <https://doi.org/10.1007/s10461-011-0052-6>
- Pintrich, P. R. (2000). An achievement goal theory perspective on issues in motivation terminology, theory, and research. *Contemporary Educational Psychology*, 25(1), 92–104. <https://doi.org/10.1006/ceps.1999.1017>
- *Pogosova, N., Yufereva, Y., Sokolova, O., Yusubova, A., Suvorov, A., & Saner, H. (2021). Telemedicine Intervention to Improve Long-Term Risk Factor Control and Body Composition in Persons with High Cardiovascular Risk: Results from a Randomized Trial: Telehealth strategies may offer an advantage over standard institutional based interventions for improvement of cardiovascular risk in high-risk patients long-term. *Global Heart*, 16(1), Article 21. <https://doi.org/10.5334/gh.825>
- *Policarpo, S., Machado, M. V., & Cortez-Pinto, H. (2021). Telemedicine as a tool for dietary intervention in NAFLD-HIV patients during the COVID-19 lockdown: A randomized controlled trial. *Clinical Nutrition ESPEN*, 43, 329–334. <https://doi.org/10.1016/j.clnesp.2021.03.031>
- Prochaska, J. J., & Prochaska, J. O. (2011). A review of multiple health behavior change interventions for primary prevention. *American Journal of Lifestyle Medicine*, 5(3), 208–221. <https://doi.org/10.1177/1559827610391883>
- Prochaska, J. J., Spring, B., & Nigg, C. R. (2008). Multiple health behavior change research: An introduction and overview. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 46(3), 181–188. <https://doi.org/10.1016/j.ypmed.2008.02.001>
- Protogerou, C., McHugh, R. K., & Johnson, B. T. (2020). How best to reduce unhealthy risk-taking behaviours? A meta-review of evidence syntheses of interventions using self-regulation principles. *Health Psychology Review*, 14(1), 86–115. <https://doi.org/10.1080/17437199.2019.1707104>
- *Ptomey, L. T., Washburn, R. A., Goetz, J. R., Sullivan, D. K., Gibson, C. A., Mayo, M. S., Krebill, R., Gorczyca, A. M., Montgomery, R. N., Honas, J. J., Helsel, B. C., & Donnelly, J. E. (2021). Weight loss interventions for adolescents with intellectual disabilities: An RCT. *Pediatrics*, 148(3), Article e2021050261. <https://doi.org/10.1542/peds.2021-050261>
- *Purcell, D. W., Latka, M. H., Metsch, L. R., Latkin, C. A., Gómez, C. A., Mizuno, Y., Arnsten, J. H., Wilkinson, J. D., Knight, K. R., Knowlton, A. R., Santibanez, S., Tobin, K. E., Rose, C. D., Valverde, E. E., Gourevitch, M. N., Eldred, L., Borkowf, C. B., & the INSPIRE Study Team. (2007). Results from a randomized controlled trial of a peer-mentoring intervention to reduce HIV transmission and increase access to care and adherence to HIV medications among HIV-seropositive injection drug users. *JAIDS Journal of Acquired Immune Deficiency Syndromes*, 46(Suppl. 2), S35–S47. <https://doi.org/10.1097/QAI.0b013e31815767c4>
- Pustejovsky, J. E. (2021). *clubSandwich: Cluster-Robust (Sandwich) variance estimators with small-sample corrections*. R package.
- Pustejovsky, J. E., & Rodgers, M. A. (2019). Testing for funnel plot asymmetry of standardized mean differences. *Research Synthesis Methods*, 10(1), 57–71. <https://doi.org/10.1002/jrsm.1332>
- Pustejovsky, J. E., & Tipton, E. (2018). Small-sample methods for cluster-robust variance estimation and hypothesis testing in fixed effects models. *Journal of Business & Economic Statistics*, 36(4), 672–683. <https://doi.org/10.1080/07350015.2016.1247004>
- Pustejovsky, J. E., & Tipton, E. (2022). Meta-analysis with robust variance estimation: Expanding the range of working models. *Prevention Science*, 23(3), 425–438. <https://doi.org/10.1007/s1121-021-01246-3>
- Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1(3), 385–401. <https://doi.org/10.1177/014662167700100306>
- *Ratliff, J. C., Palmese, L. B., Tonizzo, K. M., Chwastiak, L., & Tek, C. (2012). Contingency management for the treatment of antipsychotic-induced weight gain: A randomized controlled pilot study. *Obesity Facts*, 5(6), 919–927. <https://doi.org/10.1159/000345975>
- *Reback, C. J., Peck, J. A., Dierst-Davies, R., Nuno, M., Kamien, J. B., & Amass, L. (2010). Contingency management among homeless, out-of-treatment men who have sex with men. *Journal of Substance Abuse Treatment*, 39(3), 255–263. <https://doi.org/10.1016/j.jsat.2010.06.007>
- *Reis, L. O., Zani, E. L., Saad, R. D., Chaim, E. A., de Oliveira, L. C., & Fregonesi, A. (2012). Bariatric surgery does not interfere with sperm quality—A preliminary long-term study. *Reproductive Sciences*, 19(10), 1057–1062. <https://doi.org/10.1177/1933719112440747>
- Revelle, W. (2015). *Package “psych”—Procedures for psychological, psychometric and personality research* [Software]. <https://personality-project.org/r/psych-manual.pdf>
- Rhodes, R. E., Janssen, I., Bredin, S. S. D., Warburton, D. E. R., & Bauman, A. (2017). Physical activity: Health impact, prevalence, correlates and interventions. *Psychology & Health*, 32(8), 942–975. <https://doi.org/10.1080/08870446.2017.1325486>
- *Rhodes, S. D., Alonzo, J., Mann, L., Song, E. Y., Tanner, A. E., Arellano, J. E., Rodriguez-Celedon, R., Garcia, M., Freeman, A., Reboussin, B. A., & Painter, T. M. (2017). Small-group randomized controlled trial to increase condom use and HIV testing among Hispanic/Latino gay, bisexual, and other men who have sex with men. *American Journal of Public Health*, 107(6), 969–976. <https://doi.org/10.2105/AJPH.2017.303814>
- Rich, A., Brandes, K., Mullan, B., & Hagger, M. S. (2015). Theory of planned behavior and adherence in chronic illness: A meta-analysis. *Journal of Behavioral Medicine*, 38(4), 673–688. <https://doi.org/10.1007/s10865-015-9644-3>
- *Rinott, E., Youngster, I., Yaskolka Meir, A., Tsaban, G., Zelicha, H., Kaplan, A., Knights, D., Tuohy, K., Fava, F., Scholz, M. U., Ziv, O., Rubin, E., Tirosh, A., Rudich, A., Blüher, M., Stumvoll, M., Ceglarek, U., Clement, K., Koren, O., ... Shai, I. (2021). Effects of diet-modulated autologous fecal microbiota transplantation on weight regain. *Gastroenterology*, 160(1), 158–173.e10. <https://doi.org/10.1053/j.gastro.2020.08.041>
- Ritchwood, T. D., Ford, H., DeCoster, J., Sutton, M., & Lochman, J. E. (2015). Risky sexual behavior and substance use among adolescents: A meta-analysis. *Children and Youth Services Review*, 52, 74–88. <https://doi.org/10.1016/j.childyouth.2015.03.005>
- Roberts, B. W., Luo, J., Briley, D. A., Chow, P. I., Su, R., & Hill, P. L. (2017). A systematic review of personality trait change through intervention. *Psychological Bulletin*, 143(2), 117–141. <https://doi.org/10.1037/bul0000088>
- Robinson, K. (2013). Early disparities in mathematics gains among poor and non-poor children: Examining the role of behavioral engagement in learning. *The Elementary School Journal*, 114(1), 22–47. <https://doi.org/10.1086/670737>
- *Robles, R. R., Reyes, J. C., Colón, H. M., Sahai, H., Marrero, C. A., Matos, T. D., Calderón, J. M., & Shepard, E. W. (2004). Effects of combined counseling and case management to reduce HIV risk behaviors among Hispanic drug injectors in Puerto Rico: A randomized controlled study. *Journal of Substance Abuse Treatment*, 27(2), 145–152. <https://doi.org/10.1016/j.jsat.2004.06.004>
- Rodgers, M. A., & Pustejovsky, J. E. (2021). Evaluating meta-analytic methods to detect selective reporting in the presence of dependent effect sizes. *Psychological Methods*, 26(2), 141–160. <https://doi.org/10.1037/me0000300>

- *Rodríguez Cristóbal, J. J., Alonso-Villaverde Grote, C., Travé Mercadé, P., Pérez Santos, J. M., Peña Sendra, E., Muñoz Lloret, A., Fernández Pérez, C., Bleda Fernández, D., & the EFAP group. (2012). Randomised clinical trial of an intensive intervention in the primary care setting of patients with high plasma fibrinogen in the primary prevention of cardiovascular disease. *BMC Research Notes*, 5(1), Article 126. <https://doi.org/10.1186/1756-0500-5-126>
- Rogers, R. D., & Monsell, S. (1995). Costs of a predictable switch between simple cognitive tasks. *Journal of Experimental Psychology: General*, 124(2), 207–231. <https://doi.org/10.1037/0096-3445.124.2.207>
- *Rongkavilit, C., Naar-King, S., Wang, B., Panthong, A., Bunupuradah, T., Parsons, J. T., Phonphithak, S., Koken, J. A., Saengcharnchai, P., & Phanuphak, P. (2013). Motivational interviewing targeting risk behaviors for youth living with HIV in Thailand. *AIDS and Behavior*, 17(6), 2063–2074. <https://doi.org/10.1007/s10461-013-0407-2>
- Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48(2), 1–36. <https://doi.org/10.18637/jss.v048.i02>
- *Rotheram-Borus, M. J., Lee, M. B., Murphy, D. A., Futterman, D., Duan, N., Birnbaum, J. M., Lightfoot, M., & the Teens Linked to Care Consortium. (2001). Efficacy of a preventive intervention for youths living with HIV. *American Journal of Public Health*, 91(3), 400–405. <https://doi.org/10.2105/AJPH.91.3.400>
- *Rowe, C. L., Alberga, L., Dakof, G. A., Henderson, C. E., Ungaro, R., & Liddle, H. A. (2016). Family-based HIV and sexually transmitted infection risk reduction for drug-involved young offenders: 42-month outcomes. *Family Process*, 55(2), 305–320. <https://doi.org/10.1111/famp.12206>
- *Ruiz-Vozmediano, J., Löhnchen, S., Jurado, L., Recio, R., Rodríguez-Carrillo, A., López, M., Mustieles, V., Expósito, M., Arroyo-Morales, M., & Fernández, M. F. (2020). Influence of a multidisciplinary program of diet, exercise, and mindfulness on the quality of life of stage IIA-IIB breast cancer survivors. *Integrative Cancer Therapies*, 19, 1–11. <https://doi.org/10.1177/1534735420924757>
- *Sacco, W. P., Malone, J. I., Morrison, A. D., Friedman, A., & Wells, K. (2009). Effect of a brief, regular telephone intervention by paraprofessionals for type 2 diabetes. *Journal of Behavioral Medicine*, 32(4), 349–359. <https://doi.org/10.1007/s10865-009-9209-4>
- *Safabakhsh, L., Arbabisarjou, A., Jahantigh, M., Nazemzadeh, M., Rigi, S. N., & Nosrati, S. (2015). The effect of health promoting programs on patient's life style after coronary artery bypass graft-hospitalized in Shiraz hospitals. *Global Journal of Health Science*, 8(5), 154–159. <https://doi.org/10.5539/gjhs.v8n5p154>
- *Sakane, N., Kotani, K., Takahashi, K., Sano, Y., Tsuzaki, K., Okazaki, K., Sato, J., Suzuki, S., Morita, S., Oshima, Y., Izumi, K., Kato, M., Ishizuka, N., Noda, M., & Kuzuya, H. (2015). Effects of telephone-delivered lifestyle support on the development of diabetes in participants at high risk of type 2 diabetes: J-DOIT1, a pragmatic cluster randomised trial. *BMJ Open*, 5(8), Article e007316. <https://doi.org/10.1136/bmjopen-2014-007316>
- *Sakthivel, R. (2022). Impact of adherence counselling with hatha yoga on HIV-infected adolescents: A randomized controlled trial. HIV & AIDS Review. *International Journal of HIV-Related Problems*, 21(3), 215–221. <https://doi.org/10.5114/hivar.2022.117284>
- Salamone, J. D., Pardo, M., Yohn, S. E., López-Cruz, L., SanMiguel, N., & Correa, M. (2015). Mesolimbic dopamine and the regulation of motivated behavior. In E. Simpson & P. Balsam (Eds.), *Current topics in behavioral neurosciences* (Vol. 27, pp. 231–257). Springer. https://doi.org/10.1007/7854_2015_383
- Salas-Salvadó, J., Díaz-López, A., Ruiz-Canela, M., Basora, J., Fitó, M., Corella, D., Serra-Majem, L., Wärnberg, J., Romaguera, D., Estruch, R., Vidal, J., Martínez, J. A., Arós, F., Vázquez, C., Ros, E., Vioque, J., López-Miranda, J., Bueno-Cavanillas, A., Tur, J. A., ... the PREDIMED-Plus investigators. (2019). Effect of a lifestyle intervention program with energy-restricted Mediterranean diet and exercise on weight loss and cardiovascular risk factors: One-year results of the PREDIMED-Plus trial. *Diabetes Care*, 42(5), 777–788. <https://doi.org/10.2337/dc18-0836>
- *Samet, J. H., Blokhina, E., Cheng, D. M., Walley, A. Y., Lioznov, D., Gnatienco, N., Quinn, E. K., Briden, C., Chaisson, C. E., Toussova, O., Gifford, A. L., Raj, A., & Krupitsky, E. (2019). A strengths-based case management intervention to link HIV-positive people who inject drugs in Russia to HIV care. *Aids*, 33(9), 1467–1476. <https://doi.org/10.1097/QAD.0000000000002230>
- *Samet, J. H., Raj, A., Cheng, D. M., Blokhina, E., Briden, C., Chaisson, C. E., Walley, A. Y., Palfai, T. P., Quinn, E. K., Zvartau, E., Lioznov, D., & Krupitsky, E. (2015). HERMITAGE—A randomized controlled trial to reduce sexually transmitted infections and HIV risk behaviors among HIV-infected Russian drinkers. *Addiction*, 110(1), 80–90. <https://doi.org/10.1111/add.12716>
- *Sanchez, Z. M., Valente, J. Y., Galvão, P. P., Gubert, F. A., Melo, M. H. S., Caetano, S. C., Mari, J. J., & Cogo-Moreira, H. (2021). A cluster randomized controlled trial evaluating the effectiveness of the school-based drug prevention program #Tamojunto2.0. *Addiction*, 116(6), 1580–1592. <https://doi.org/10.1111/add.15358>
- *Santa-Maria, C. A., Coughlin, J. W., Sharma, D., Armanios, M., Blackford, A. L., Schreyer, C., Dalcin, A., Carpenter, A., Jerome, G. J., Armstrong, D. K., Chaudhry, M., Cohen, G. I., Connolly, R. M., Fetting, J., Miller, R. S., Smith, K. L., Snyder, C., Wolfe, A., Wolff, A. C., ... Stearns, V. (2020). The effects of a remote-based weight loss program on adipocytokines, metabolic markers, and telomere length in breast cancer survivors: The POWER-remote trial. *Clinical Cancer Research*, 26(12), 3024–3034. <https://doi.org/10.1158/1078-0432.CCR-19-2935>
- *Santschi, V., Wuerzner, G., Pais, B., Chiolerio, A., Schaller, P., Cloutier, L., Paradis, G., & Burnier, M. (2021). Team-based care for improving hypertension management: A pragmatic randomized controlled trial. *Frontiers in Cardiovascular Medicine*, 8, Article 760662. <https://doi.org/10.3389/fcvm.2021.760662>
- *Sarrafzadegan, N., Kelishadi, R., Esmailzadeh, A., Mohammadifard, N., Rabiei, K., Roohafza, H., Azadbakht, L., Bahonar, A., Sadri, G., Amani, A., Heidari, S., & Malekafzali, H. (2009). Do lifestyle interventions work in developing countries? Findings from the Isfahan Healthy Heart Program in the Islamic Republic of Iran. *Bulletin of the World Health Organization*, 87(1), 39–50. <https://doi.org/10.2471/BLT.07.049841>
- *Satre, D. D., Leibowitz, A., Sterling, S. A., Lu, Y., Travis, A., & Weisner, C. (2016). A randomized clinical trial of Motivational Interviewing to reduce alcohol and drug use among patients with depression. *Journal of Consulting and Clinical Psychology*, 84(7), 571–579. <https://doi.org/10.1037/ccp0000096>
- *Savoye, M., Shaw, M., Dziura, J., Tamborlane, W. V., Rose, P., Guandalini, C., Goldberg-Gell, R., Burgert, T. S., Cali, A. M., Weiss, R., & Caprio, S. (2007). Effects of a weight management program on body composition and metabolic parameters in overweight children: A randomized controlled trial. *JAMA: Journal of the American Medical Association*, 297(24), 2697–2704. <https://doi.org/10.1001/jama.297.24.2697>
- *Sawyer, A. T., Wheeler, J., Jennelle, P., Pepe, J., & Robinson, P. S. (2020). A randomized controlled trial of a motivational interviewing intervention to improve whole-person lifestyle. *Journal of Primary Care & Community Health*, 11, 1–7. <https://doi.org/10.1177/2150132720922714>
- *Schäfer, I., Lotzin, A., Hiller, P., Sehner, S., Driessen, M., Hillemacher, T., Schäfer, M., Scherbaum, N., Schneider, B., & Grundmann, J. (2019). A multisite randomized controlled trial of Seeking Safety vs. Relapse Prevention Training for women with co-occurring posttraumatic stress disorder and substance use disorders. *European Journal of Psychotraumatology*, 10(1), Article 1577092. <https://doi.org/10.1080/2008198.2019.1577092>
- *Sharma, A. K., Baig, V. N., Ahuja, J., Sharma, S., Panwar, R. B., Katoch, V. M., & Gupta, R. (2021). Efficacy of IVRS-based mHealth intervention in reducing cardiovascular risk in metabolic syndrome: A cluster randomized

- trial. *Diabetes & Metabolic Syndrome*, 15(5), Article 102182. <https://doi.org/10.1016/j.dsx.2021.06.019>
- *Shaul, L., de Waal, M., Blankers, M., Koeter, M. W. J., Schippers, G. M., & Goudriaan, A. E. (2020). Effectiveness of a brief motivation enhancing intervention on treatment initiation, treatment retention and abstinence: Results from a multi-site cluster-randomized trial. *Journal of Substance Abuse Treatment*, 110, 28–36. <https://doi.org/10.1016/j.jsat.2019.12.002>
- *Sherwood, N. E., Crain, A. L., Seburg, E. M., Butryn, M. L., Forman, E. M., Crane, M. M., Levy, R. L., Kunin-Batson, A. S., & Jeffery, R. W. (2022). BestFIT sequential multiple assignment randomized trial results: A SMART approach to developing individualized weight loss treatment sequences. *Annals of Behavioral Medicine*, 56(3), 291–304. <https://doi.org/10.1093/abnm/kaab061>
- *Shibayama, T., Kobayashi, K., Takano, A., Kadowaki, T., & Kazuma, K. (2007). Effectiveness of lifestyle counseling by certified expert nurse of Japan for non-insulin-treated diabetic outpatients: A 1-year randomized controlled trial. *Diabetes Research and Clinical Practice*, 76(2), 265–268. <https://doi.org/10.1016/j.diabres.2006.09.017>
- Shrout, P. E., & Bolger, N. (2002). Mediation in experimental and nonexperimental studies: New procedures and recommendations. *Psychological Methods*, 7(4), 422–445. <https://doi.org/10.1037/1082-989X.7.4.422>
- *Siddiqui, F., Kurbasic, A., Lindblad, U., Nilsson, P. M., & Bennet, L. (2017). Effects of a culturally adapted lifestyle intervention on cardio-metabolic outcomes: A randomized controlled trial in Iraqi immigrants to Sweden at high risk for Type 2 diabetes. *Metabolism: Clinical and Experimental*, 66, 1–13. <https://doi.org/10.1016/j.metabol.2016.10.001>
- *Simkin-Silverman, L., Wing, R. R., Hansen, D. H., Klem, M. L., Pasagian-Macaulay, A., Meilahn, E. N., & Kuller, L. H. (1995). Prevention of cardiovascular risk factor elevations, in healthy premenopausal women. *Preventive Medicine*, 24(5), 509–517. <https://doi.org/10.1006/pmed.1995.1081>
- *Simkin-Silverman, L. R., Wing, R. R., Boraz, M. A., Meilahn, E. N., & Kuller, L. H. (1998). Maintenance of cardiovascular risk factor changes among middle-aged women in a lifestyle intervention trial. *Women's Health*, 4, 255–271. <https://europepmc.org/article/med/9787651>
- Simpson, E. H. (1951). The interpretation of interaction in contingency tables. *Journal of the Royal Statistical Society. Series B. Methodological*, 13(2), 238–241. <https://doi.org/10.1111/j.2517-6161.1951.tb00088.x>
- *Singh, R. B., Sharma, V. K., Rastogi, S. S., & Singh, N. K. (1992). In patients with mild hypertension, does exercise and a gradual rather than abrupt increase in fatty acid and salt intake cause less rise in cardiovascular risk factors? *Clinical Nutrition*, 11(5), 309–314. [https://doi.org/10.1016/0261-5614\(92\)90009-F](https://doi.org/10.1016/0261-5614(92)90009-F)
- *Skjåkødegård, H. F., Conlon, R. P. K., Hystad, S. W., Roelants, M., Olsson, S. J. G., Frisk, B., Wilfley, D. E., Danielsen, Y. S., & Juliusson, P. B. (2022). Family-based treatment of children with severe obesity in a public healthcare setting: Results from a randomized controlled trial. *Clinical Obesity*, 12(3), Article e12513. <https://doi.org/10.1111/cob.12513>
- Smedley, B. D., & Syme, S. L. (2001). Committee on capitalizing on social science and behavioral research to improve the public's health. Promoting health: Intervention strategies from social and behavioral research. *American Journal of Health Promotion*, 15(3), 149–166. <https://doi.org/10.4278/0890-1171-15.3.149>
- *Smith, L. (2007). *Support for Healthy Eating and Exercise (SHEE)* [Ph.D. dissertation, University of Colorado]. Dissertations & Theses: Full Text (Publication No. AAT 3315760).
- Smith, L. E., Amlôt, R., Weinman, J., Yiend, J., & Rubin, G. J. (2017). A systematic review of factors affecting vaccine uptake in young children. *Vaccine*, 35(45), 6059–6069. <https://doi.org/10.1016/j.vaccine.2017.09.046>
- *Soares, T. S., Piovesan, C. H., Gustavo, A. S., Macagnan, F. E., Bodanese, L. C., & Feoli, A. M. (2014). Alimentary habits, physical activity, and Framingham global risk score in metabolic syndrome. *Arquivos Brasileiros de Cardiologia*, 102(4), 374–382. <https://doi.org/10.5935/abc.20140029>
- Sobel, M. E. (1982). Asymptotic confidence intervals for indirect effects in structural equation models. *Sociological Methodology*, 13, 290–312. <https://doi.org/10.2307/270723>
- Somaa, F., Asghar, A., & Hamid, P. F. (2021). Academic performance and emotional intelligence with age and gender as moderators: A meta-analysis. *Developmental Neuropsychology*, 46(8), 537–554. <https://doi.org/10.1080/87565641.2021.1999455>
- *Song, K. J. (2003). The effects of self-efficacy promoting cardiac rehabilitation program on self-efficacy, health behavior, and quality of life. *Taehan Kanho Hakhoe Chi*, 33(4), 510–518. <https://doi.org/10.4040/jkan.2003.33.4.510>
- Spearing, M. K., Post, R. M., Leverich, G. S., Brandt, D., & Nolen, W. (1997). Modification of the Clinical Global Impressions (CGI) Scale for use in bipolar illness (BP): The CGI-BP. *Psychiatry Research*, 73(3), 159–171. [https://doi.org/10.1016/S0165-1781\(97\)00123-6](https://doi.org/10.1016/S0165-1781(97)00123-6)
- *Spring, B., Doran, N., Pagoto, S., Schneider, K., Pingitore, R., & Hedeker, D. (2004). Randomized controlled trial for behavioral smoking and weight control treatment: Effect of concurrent versus sequential intervention. *Journal of Consulting and Clinical Psychology*, 72(5), 785–796. <https://doi.org/10.1037/0022-006X.72.5.785>
- Spring, B., Howe, D., Berendsen, M., McFadden, H. G., Hitchcock, K., Rademaker, A. W., & Hitsman, B. (2009). Behavioral intervention to promote smoking cessation and prevent weight gain: A systematic review and meta-analysis. *Addiction*, 104(9), 1472–1486. <https://doi.org/10.1111/j.1360-0443.2009.02610.x>
- *St Lawrence, J., Eldridge, G. D., Shelby, M. C., Little, C. E., Brasfield, T. L., & O'Bannon, R. E., III. (1997). HIV risk reduction for incarcerated women: A comparison of brief interventions based on two theoretical models. *Journal of Consulting and Clinical Psychology*, 65(3), 504–509. <https://doi.org/10.1037/0022-006X.65.3.504>
- *St Lawrence, J. S., Jefferson, K. W., Alleyne, E., & Brasfield, T. L. (1995). Comparison of education versus behavioral skills training interventions in lowering sexual HIV-risk behavior of substance-dependent adolescents. *Journal of Consulting and Clinical Psychology*, 63(1), 154–157. <https://doi.org/10.1037/0022-006X.63.1.154>
- *Stasinaki, A., Büchter, D., Shih, C. I., Heldt, K., Güsewell, S., Brogle, B., Farpour-Lambert, N., Kowatsch, T., & l'Allemand, D. (2021). Effects of a novel mobile health intervention compared to a multi-component behaviour changing program on body mass index, physical capacities and stress parameters in adolescents with obesity: A randomized controlled trial. *BMC Pediatrics*, 21(1), Article 308. <https://doi.org/10.1186/s12887-021-02781-2>
- Stedry, A. C., & Kay, E. (1966). The effects of goal difficulty on performance: A field experiment. *Behavioral Science*, 11(6), 459–470. <https://doi.org/10.1002/bs.3830110605>
- *Stephenson, R., Sullivan, S. P., Mitchell, J. W., Johnson, B. A., & Sullivan, P. S. (2022). Efficacy of a telehealth delivered couples' HIV counseling and testing (CHTC) intervention to improve formation and adherence to safer sexual agreements among male couples in the US: Results from a randomized control trial. *AIDS and Behavior*, 26(8), 2813–2824. <https://doi.org/10.1007/s10461-022-03619-3>
- *Stephenson, R., Vwalika, B., Greenberg, L., Ahmed, Y., Vwalika, C., Chomba, E., Kilembe, W., Tichacek, A., & Allen, S. (2011). A randomized controlled trial to promote long-term contraceptive use among HIV-serodiscordant and concordant positive couples in Zambia. *Journal of Women's Health*, 20(4), 567–574. <https://doi.org/10.1089/jwh.2010.2113>
- *Stewart, K. E., Wright, P. B., Montgomery, B. E. E., Cornell, C., Gullette, D., Pulley, L., Ounpraseuth, S., Thostenson, J., & Booth, B. (2017). Reducing risky sex among rural African American cocaine users: A controlled trial. *Journal of Health Care for the Poor and Underserved*, 28(1), 528–547. <https://doi.org/10.1353/hpu.2017.0038>

- *Stier-Jarmer, M., Frisch, D., Neuy, S., & Schuh, A. (2021). A 3-week naturopathic intervention improves HbA_{1c}, weight, and quality of life among overweight and obese adults with type 2 diabetes: 6-month results from a randomized trial. *Alternative Therapies in Health and Medicine*, 27(S1), 61–71.
- *Strandberg, T. E., Salomaa, V. V., Naukkarinen, V. A., Vanhanen, H. T., Sarna, S. J., & Miettinen, T. A. (1991). Long-term mortality after 5-year multi-factorial primary prevention of cardiovascular diseases. *Journal of the American Medical Association*, 266(9), 1225–1229. <https://doi.org/10.1001/jama.1991.03470090059032>
- *Sturgeon, K. M., Dean, L. T., Heroux, M., Kane, J., Bauer, T., Palmer, E., Long, J., Lynch, S., Jacobs, L., Sarwer, D. B., Leonard, M. B., & Schmitz, K. (2017). Commercially available lifestyle modification program: Randomized controlled trial addressing heart and bone health in BRCA1/2+ breast cancer survivors after risk-reducing salpingo-oophorectomy. *Journal of Cancer Survivorship: Research and Practice*, 11(2), 246–255. <https://doi.org/10.1007/s11764-016-0582-z>
- Sunderrajan, A., White, B., Durantini, M., Sanchez, F., Glasman, L., & Albarracín, D. (2021). Complex solutions for a complex problem: A meta-analysis of the efficacy of multiple-behavior interventions on change in outcomes related to HIV. *Health Psychology*, 40(9), 642–653. <https://doi.org/10.1037/hea0001088>
- *Svetkey, L. P., Pollak, K. I., Yancy, W. S., Jr., Dolor, R. J., Batch, B. C., Samsa, G., Matchar, D. B., & Lin, P. H. (2009). Hypertension improvement project: Randomized trial of quality improvement for physicians and lifestyle modification for patients. *Hypertension*, 54(6), 1226–1233. <https://doi.org/10.1161/HYPERTENSIONAHA.109.134874>
- Sweeting, M. J., Sutton, A. J., & Lambert, P. C. (2004). What to add to nothing? Use and avoidance of continuity corrections in meta-analysis of sparse data. *Statistics in Medicine*, 23(9), 1351–1375. <https://doi.org/10.1002/sim.1761>
- *Sylvia, L. G., Pegg, S. L., Dufour, S. C., Janos, J. A., Bernstein, E. E., Chang, W. C., Hall, N. E., Ellard, K. K., Nierenberg, A. A., & Deckersbach, T. (2019). Pilot study of a lifestyle intervention for bipolar disorder: Nutrition exercise wellness treatment (NEW Tx). *Journal of Affective Disorders*, 250, 278–283. <https://doi.org/10.1016/j.jad.2019.03.033>
- *Taheri, S., Zaghoul, H., Chagoury, O., Elhadad, S., Ahmed, S. H., El Khatib, N., Amona, R. A., El Nahas, K., Suleiman, N., Alnaama, A., Al-Hamaq, A., Charlson, M., Wells, M. T., Al-Abdulla, S., & Abou-Samra, A. B. (2020). Effect of intensive lifestyle intervention on bodyweight and glycaemia in early type 2 diabetes (DIADEM-I): An open-label, parallel-group, randomised controlled trial. *The Lancet. Diabetes & Endocrinology*, 8(6), 477–489. [https://doi.org/10.1016/S2213-8587\(20\)30117-0](https://doi.org/10.1016/S2213-8587(20)30117-0)
- *Talavera, G. A., Castañeda, S. F., Mendoza, P. M., Lopez-Gurrola, M., Roesch, S., Pichardo, M. S., Garcia, M. L., Muñoz, F., & Gallo, L. C. (2021). Latinos understanding the need for adherence in diabetes (LUNA-D): A randomized controlled trial of an integrated team-based care intervention among Latinos with diabetes. *Translational Behavioral Medicine*, 11(9), 1665–1675. <https://doi.org/10.1093/tbm/ibab052>
- *Talvi, A. I., Järvisalo, J. O., & Knuts, L. R. (1999). A health promotion programme for oil refinery employees: Changes of health promotion needs observed at three years. *Occupational Medicine*, 49(2), 93–101. <https://doi.org/10.1093/occmed/49.2.93>
- Tanner-Smith, E. E., & Tipton, E. (2014). Robust variance estimation with dependent effect sizes: Practical considerations including a software tutorial in Stata and SPSS. *Research Synthesis Methods*, 5(1), 13–30. <https://doi.org/10.1002/jrsm.1091>
- Tanner-Smith, E. E., Tipton, E., & Polanin, J. R. (2016). Handling complex meta-analytic data structures using robust variance estimates: A tutorial in R. *Journal of Developmental and Life-Course Criminology*, 2(1), 85–112. <https://doi.org/10.1007/s40865-016-0026-5>
- *Tapper, K., Shaw, C., Ilsley, J., Hill, A. J., Bond, F. W., & Moore, L. (2009). Exploratory randomised controlled trial of a mindfulness-based weight loss intervention for women. *Appetite*, 52(2), 396–404. <https://doi.org/10.1016/j.appet.2008.11.012>
- *Tate, D. F., Jackvony, E. H., & Wing, R. R. (2003). Effects of Internet behavioral counseling on weight loss in adults at risk for type 2 diabetes: A randomized trial. *JAMA: Journal of the American Medical Association*, 289(14), 1833–1836. <https://doi.org/10.1001/jama.289.14.1833>
- *Temple, J. R., Baumler, E., Wood, L., Thiel, M., Peskin, M., & Torres, E. (2021). A dating violence prevention program for middle school youth: A cluster randomized trial. *Pediatrics*, 148(5), Article e2021052880. <https://doi.org/10.1542/peds.2021-052880>
- *Thato, R., Jenkins, R. A., & Dusitsin, N. (2008). Effects of the culturally-sensitive comprehensive sex education programme among Thai secondary school students. *Journal of Advanced Nursing*, 62(4), 457–469. <https://doi.org/10.1111/j.1365-2648.2008.04609.x>
- Thoits, P. A. (2010). Stress and health: Major findings and policy implications. *Journal of Health and Social Behavior*, 51(1_Suppl), S41–S53. <https://doi.org/10.1177/0022146510383499>
- *Thomas, E. A., Zaman, A., Sloggett, K. J., Steinke, S., Grau, L., Catenacci, V. A., Cornier, M. A., & Rynders, C. A. (2022). Early time-restricted eating compared with daily caloric restriction: A randomized trial in adults with obesity. *Obesity*, 30(5), 1027–1038. <https://doi.org/10.1002/oby.23420>
- *Thompson, D., Baranowski, T., Cullen, K., Watson, K., Liu, Y., Canada, A., Bhatt, R., & Zakeri, I. (2008). Food, fun, and fitness internet program for girls: Pilot evaluation of an e-Health youth obesity prevention program examining predictors of obesity. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 47(5), 494–497. <https://doi.org/10.1016/j.ypmed.2008.07.014>
- *Thuita, A. W., Kiage, B. N., Onyango, A. N., & Makokha, A. O. (2020). Effect of a nutrition education programme on the metabolic syndrome in type 2 diabetes mellitus patients at a level 5 Hospital in Kenya: “a randomized controlled trial.” *BMC Nutrition*, 6(1), Article 30. <https://doi.org/10.1186/s40795-020-00355-6>
- *Thurman, T. R., Kidman, R., Carton, T. W., & Chiroro, P. (2016). Psychological and behavioral interventions to reduce HIV risk: Evidence from a randomized control trial among orphaned and vulnerable adolescents in South Africa. *AIDS Care*, 28(Suppl 1), 8–15. <https://doi.org/10.1080/09540121.2016.1146213>
- *Tingey, L., Chambers, R., Littlepage, S., Lee, A., Slimp, A., Melgar, L., Cwik, M., Gaydos, C., Rompalo, A., & Rosenstock, S. (2021). “Empowering our people”: Impact of a culturally adapted evidence-based intervention on sexually transmitted infection risks among native Americans with binge substance use. *Sexually Transmitted Diseases*, 48(9), 622–628. <https://doi.org/10.1097/OLQ.0000000000001418>
- *Tobin, K., Davey-Rothwell, M. A., Nonyane, B. A. S., Knowlton, A., Wissow, L., & Latkin, C. A. (2017). RCT of an integrated CBT-HIV intervention on depressive symptoms and HIV risk. *PLOS ONE*, 12(12), Article e0187180. <https://doi.org/10.1371/journal.pone.0187180>
- *Toobert, D. J., Glasgow, R. E., & Radcliffe, J. L. (2000). Physiologic and related behavioral outcomes from the Women’s Lifestyle Heart Trial. *Annals of Behavioral Medicine*, 22(1), 1–9. <https://doi.org/10.1007/BF02895162>
- *Toobert, D. J., Glasgow, R. E., Strycker, L. A., Barrera, M., Jr., Ritzwoller, D. P., & Weidner, G. (2007). Long-term effects of the Mediterranean lifestyle program: A randomized clinical trial for postmenopausal women with type 2 diabetes. *The International Journal of Behavioral Nutrition and Physical Activity*, 4(1), Article 1. <https://doi.org/10.1186/1479-5868-4-1>
- *Toobert, D. J., Strycker, L. A., Barrera, M., Jr., Osuna, D., King, D. K., & Glasgow, R. E. (2011). Outcomes from a multiple risk factor diabetes self-management trial for Latinas: ¡Viva Bien! *Annals of Behavioral Medicine*, 41(3), 310–323. <https://doi.org/10.1007/s12160-010-9256-7>
- Topa, G., & Moriano, L. (2010). Theory of planned behavior and smoking: Meta-analysis and SEM model. *Substance Abuse and Rehabilitation*, 1, 23–33. <https://doi.org/10.2147/SAR.S15168>

- *Towfighi, A., Cheng, E. M., Hill, V. A., Barry, F., Lee, M., Valle, N. P., Mittman, B., Ayala-Rivera, M., Moreno, L., Espinosa, A., Dombish, H., Wang, D., Ochoa, D., Chu, A., Atkins, M., & Vickrey, B. G. (2020). Results of a pilot trial of a lifestyle intervention for stroke survivors: Healthy eating and lifestyle after stroke. *Journal of Stroke and Cerebrovascular Diseases*, 29(12), Article 105323. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2020.105323>
- *Trento, M., Passera, P., Tomalino, M., Pagnozzi, F., Pomero, F., Vaccari, P., Bajardi, M., Molinatti, G. M., & Porta, M. (1998). Therapeutic group education in the follow-up of patients with non-insulin treated, non-insulin dependent diabetes mellitus. *Diabetes, Nutrition & Metabolism*, 11, 212–216.
- *Tripette, J., Gando, Y., Murakami, H., Kawakami, R., Tanisawa, K., Ohno, H., Konishi, K., Tanimoto, M., Tanaka, N., Kawano, H., Yamamoto, K., Morishita, A., Iemitsu, M., Sanada, K., Miyatake, N., & Miyachi, M. (2021). Effect of a 1-year intervention comprising brief counselling sessions and low-dose physical activity recommendations in Japanese adults, and retention of the effect at 2 years: A randomized trial. *BMC Sports Science, Medicine & Rehabilitation*, 13(1), Article 133. <https://doi.org/10.1186/s13102-021-00360-7>
- *Tucker, J. S., D'Amico, E. J., Ewing, B. A., Miles, J. N., & Pedersen, E. R. (2017). A group-based motivational interviewing brief intervention to reduce substance use and sexual risk behavior among homeless young adults. *Journal of Substance Abuse Treatment*, 76, 20–27. <https://doi.org/10.1016/j.jsat.2017.02.008>
- *Tudor-Smith, C., Nutbeam, D., Moore, L., & Catford, J. (1998). Effects of the Heartbeat Wales programme over five years on behavioural risks for cardiovascular disease: Quasi-experimental comparison of results from Wales and a matched reference area. *The BMJ*, 316(7134), 818–822. <https://doi.org/10.1136/bmj.316.7134.818>
- *Tzilos Wernette, G., Plegue, M., Kahler, C. W., Sen, A., & Zlotnick, C. (2018). A pilot randomized controlled trial of a computer-delivered brief intervention for substance use and risky sex during pregnancy. *Journal of Women's Health*, 27(1), 83–92. <https://doi.org/10.1089/jwh.2017.6408>
- *Uemura, K., Yamada, M., & Okamoto, H. (2021). The effectiveness of an active learning program in promoting a healthy lifestyle among older adults with low health literacy: A randomized controlled trial. *Gerontology*, 67(1), 25–35. <https://doi.org/10.1159/000511357>
- *Ussher, M., Lewis, S., Aveyard, P., Manyonda, I., West, R., Lewis, B., Marcus, B., Riaz, M., Taylor, A., Daley, A., & Coleman, T. (2015). Physical activity for smoking cessation in pregnancy: Randomised controlled trial. *The BMJ*, 350, Article h2145. <https://doi.org/10.1136/bmj.h2145>
- *Ussher, M., West, R., McEwen, A., Taylor, A., & Steptoe, A. (2003). Efficacy of exercise counselling as an aid for smoking cessation: A randomized controlled trial. *Addiction*, 98(4), 523–532. <https://doi.org/10.1046/j.1360-0443.2003.00346.x>
- *Ussher, M., West, R., McEwen, A., Taylor, A., & Steptoe, A. (2007). Randomized controlled trial of physical activity counseling as an aid to smoking cessation: 12 month follow-up. *Addictive Behaviors*, 32(12), 3060–3064. <https://doi.org/10.1016/j.addbeh.2007.04.009>
- *Usta, E., & Aygin, D. (2020). Prospective randomized trial on effects of structured training and counseling on depression, body image, and quality of life. *Bariatric Surgical Practice and Patient Care*, 15(2), 55–62. <https://doi.org/10.1089/bari.2019.0028>
- *Vale, M. J., Jelinek, M. V., Best, J. D., Dart, A. M., Grigg, L. E., Hare, D. L., Ho, B. P., Newman, R. W., McNeil, J. J., & the COACH Study Group. (2003). Coaching patients On Achieving Cardiovascular Health (COACH): A multicenter randomized trial in patients with coronary heart disease. *Archives of Internal Medicine*, 163(22), 2775–2783. <https://doi.org/10.1001/archinte.163.22.2775>
- *Valente, J. Y., Cogo-Moreira, H., & Sanchez, Z. M. (2020). Decision-making skills as a mediator of the #Tamojuntto school-based prevention program: Indirect effects for drug use and school violence of a cluster-randomized trial. *Drug and Alcohol Dependence*, 206, Article 107718. <https://doi.org/10.1016/j.drugalcdep.2019.107718>
- *Vamvakis, A., Gkaliagkousi, E., Lazaridis, A., Grammatikopoulou, M. G., Triantafyllou, A., Nikolaidou, B., Koletsos, N., Anyfanti, P., Tzimos, C., Zebekakis, P., & Douma, S. (2020). Impact of intensive lifestyle treatment (diet plus exercise) on endothelial and vascular function, arterial stiffness and blood pressure in stage 1 hypertension: Results of the HINTreat randomized controlled trial. *Nutrients*, 12(5), Article 1326. <https://doi.org/10.3390/nu12051326>
- *van Elderen, T., Maes, S., Seegers, G., Kragten, H., & Relik-Van Wely, L. (1994). Effects of a post-hospitalization group health education program for patients with coronary heart disease. *Psychology and Health*, 9(4), 317–330. <https://doi.org/10.1080/08870449408407490>
- *Van Stappen, V., Cardon, G., De Craemer, M., Mavrogianni, C., Usheva, N., Kivelä, J., Wikström, K., De Miquel-Etayo, P., González-Gil, E. M., Radó, A. S., Nánási, A., Iotova, V., Manios, Y., & Brondeel, R. (2021). The effect of a cluster-randomized controlled trial on lifestyle behaviors among families at risk for developing type 2 diabetes across Europe: The Feel4Diabetes-study. *The International Journal of Behavioral Nutrition and Physical Activity*, 18(1), Article 86. <https://doi.org/10.1186/s12966-021-01153-4>
- *van Zuilen, A., Blankestijn, P., van Buren, M., ten Dam, M., Kaasjager, K., Ligtenberg, G., Sijpkens, Y., Sluiter, H., van de Ven, P., Vervoort, G., Vleming, L., Bots, M., & Wetzels, J. (2011). Nurse practitioners improve quality of care in chronic kidney disease: Two-year results of a randomised study. *The Netherlands Journal of Medicine*, 69(11), 517–526.
- *Vanninen, E., Uusitupa, M., Siitonen, O., Laitinen, J., & Länsimies, E. (1992). Habitual physical activity, aerobic capacity and metabolic control in patients with newly-diagnosed type 2 (non-insulin-dependent) diabetes mellitus: Effect of 1-year diet and exercise intervention. *Diabetologia*, 35(4), 340–346. <https://doi.org/10.1007/BF00401201>
- *Vaughn, A. E., Hennink-Kaminski, H., Moore, R., Burney, R., Chittams, J. L., Parker, P., Luecking, C. T., Hales, D., & Ward, D. S. (2021). Evaluating a child care-based social marketing approach for improving children's diet and physical activity: Results from the Healthy Me, Healthy We cluster-randomized controlled trial. *Translational Behavioral Medicine*, 11(3), 775–784. <https://doi.org/10.1093/tbm/ibaa113>
- *Vaz, C. L., Carnes, N., Pousti, B., Zhao, H., & Williams, K. J. (2021). A randomized controlled trial of an innovative, user-friendly, interactive smartphone app-based lifestyle intervention for weight loss. *Obesity Science & Practice*, 7(5), 555–568. <https://doi.org/10.1002/osp4.503>
- Venditti, E. M., Wylie-Rosett, J., Delahanty, L. M., Mele, L., Hoskin, M. A., Edelstein, S. L., & the Diabetes Prevention Program Research Group. (2014). Short and long-term lifestyle coaching approaches used to address diverse participant barriers to weight loss and physical activity adherence. *The International Journal of Behavioral Nutrition and Physical Activity*, 11(1), Article 16. <https://doi.org/10.1186/1479-5868-11-16>
- *Verduci, E., Banderali, G., Di Profio, E., Vizzuso, S., Zuccotti, G., & Radaelli, G. (2021). Effect of individual- versus collective-based nutritional-lifestyle intervention on the atherogenic index of plasma in children with obesity: A randomized trial. *Nutrition & Metabolism*, 18(1), Article 11. <https://doi.org/10.1186/s12986-020-00537-w>
- Verplanken, B., & Orbell, S. (2003). Reflections on past behavior: A self-report index of habit strength. *Journal of Applied Social Psychology*, 33(6), 1313–1330. <https://doi.org/10.1111/j.1559-1816.2003.tb01951.x>
- Vevea, J. L., & Hedges, L. V. (1995). A general linear model for estimating effect size in the presence of publication bias. *Psychometrika*, 60(3), 419–435. <https://doi.org/10.1007/BF02294384>
- *Vickers, K. S., Patten, C. A., Lewis, B. A., Clark, M. M., Ussher, M., Ebbert, J. O., Croghan, I. T., Decker, P. A., Hathaway, J., Marcus, B. H., & Hurt, R. D. (2009). Feasibility of an exercise counseling intervention for depressed women smokers. *Nicotine & Tobacco Research*, 11(8), 985–995. <https://doi.org/10.1093/ntr/ntp101>

- Viechtbauer, W. (2019). *Package "metafor"* [Computer software]. Comprehensive R Archive Network.
- *Viesters, L., Verhagen, E. A. L. M., Bongers, P. M., & van der Beek, A. J. (2018). Effectiveness of a worksite intervention for male construction workers on dietary and physical activity behaviors, body mass index, and health outcomes: Results of randomized controlled trial. *American Journal of Health Promotion*, 32(3), 795–805. <https://doi.org/10.1177/0890117117694450>
- *von Gruenigen, V. E., Courneya, K. S., Gibbons, H. E., Kavanagh, M. B., Waggoner, S. E., & Lerner, E. (2008). Feasibility and effectiveness of a lifestyle intervention program in obese endometrial cancer patients: A randomized trial. *Gynecologic Oncology*, 109(1), 19–26. <https://doi.org/10.1016/j.ygyno.2007.12.026>
- Vos, T., Lim, S. S., Abbafati, C., Abbas, K. M., Abbasi, M., Abbasifard, M., Abbasi-Kangevari, M., Abbastabar, H., Abd-Allah, F., Abdelalim, A., Abdollahi, M., Abdollahpour, I., Abolhassani, H., Aboyans, V., Abrams, E. M., Abreu, L. G., Guimarães, L., Abrego, M. R. M., Abu-Raddad, L. J. ... the GBD 2019 Diseases and Injuries Collaborators. (2020). Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: A systematic analysis for the Global Burden of Disease Study 2019. *Lancet*, 396(10258), 1204–1222. [https://doi.org/10.1016/S0140-6736\(20\)30925-9](https://doi.org/10.1016/S0140-6736(20)30925-9)
- *Wadden, T. A., Berkowitz, R. I., Womble, L. G., Sarwer, D. B., Phelan, S., Cato, R. K., Hesson, L. A., Osei, S. Y., Kaplan, R., & Stunkard, A. J. (2005). Randomized trial of lifestyle modification and pharmacotherapy for obesity. *The New England Journal of Medicine*, 353(20), 2111–2120. <https://doi.org/10.1056/NEJMoa050156>
- *Wadden, T. A., Tronieri, J. S., Sugimoto, D., Lund, M. T., Auerbach, P., Jensen, C., & Rubino, D. (2020). Liraglutide 3.0 mg and intensive behavioral therapy (IBT) for obesity in primary care: The SCALE IBT randomized controlled trial. *Obesity*, 28(3), 529–536. <https://doi.org/10.1002/oby.22726>
- Walker, S. N., & Hill-Polerecky, D. M. (1996). *Psychometric evaluation of the health-promoting lifestyle profile II*. University of Nebraska Medical Center.
- *Wandera, B., Tumwesigye, N. M., Nankabirwa, J. I., Mafigiri, D. K., Parkes-Ratanshi, R. M., Kapiga, S., Hahn, J., & Sethi, A. K. (2017). Efficacy of a single, brief alcohol reduction intervention among men and women living with HIV/AIDS and using alcohol in Kampala, Uganda: A randomized trial. [JIAPAC]. *Journal of the International Association of Providers of AIDS Care*, 16(3), 276–285. <https://doi.org/10.1177/2325957416649669>
- *Wang, H., Zhou, J., Huang, L., Li, X., Fennie, K. P., & Williams, A. B. (2010). Effects of nurse-delivered home visits combined with telephone calls on medication adherence and quality of life in HIV-infected heroin users in Hunan of China. *Journal of Clinical Nursing*, 19(3–4), 380–388. <https://doi.org/10.1111/j.1365-2702.2009.03048.x>
- *Wang, X., Liu, D., Du, M., Hao, R., Zheng, H., & Yan, C. (2020). The role of text messaging intervention in Inner Mongolia among patients with type 2 diabetes mellitus: A randomized controlled trial. *BMC Medical Informatics and Decision Making*, 20(1), Article 90. <https://doi.org/10.1186/s12911-020-01129-7>
- Ware, J. E., Jr., & Sherbourne, C. D. (1992). The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Medical Care*, 30(6), 473–483. <https://doi.org/10.1097/00005650-199206000-00002>
- *Watanabe, M., Okayama, A., Shimamoto, K., & Ueshima, H. (2007). Short-term effectiveness of an individual counseling program for impaired fasting glucose and mild type 2 diabetes in Japan: A multicenter randomized control trial. *Asia Pacific Journal of Clinical Nutrition*, 16(3), 489–497.
- *Watkins, L. L., Sherwood, A., Feinglos, M., Hinderliter, A., Babyak, M., Gullette, E., Waugh, R., & Blumenthal, J. A. (2003). Effects of exercise and weight loss on cardiac risk factors associated with syndrome X. *Archives of Internal Medicine*, 163(16), 1889–1895. <https://doi.org/10.1001/archinte.163.16.1889>
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, 54(6), 1063–1070. <https://doi.org/10.1037/0022-3514.54.6.1063>
- *Webb, D. R., Khunti, K., Gray, L. J., Srinivasan, B. T., Farooqi, A., Wareham, N., Griffin, S. C., Davies, M. J., & the ADDITION-Leicester study group. (2012). Intensive multifactorial intervention improves modelled coronary heart disease risk in screen-detected Type 2 diabetes mellitus: A cluster randomized controlled trial. *Diabetic Medicine*, 29(4), 531–540. <https://doi.org/10.1111/j.1464-5491.2011.03441.x>
- *Wechsberg, W. M., Lusenro, W. K., Lam, W. K., Parry, C. D., & Morojele, N. K. (2006). Substance use, sexual risk, and violence: HIV prevention intervention with sex workers in Pretoria. *AIDS and Behavior*, 10(2), 131–137. <https://doi.org/10.1007/s10461-005-9036-8>
- *Wechsberg, W. M., Zule, W. A., El-Bassel, N., Doherty, I. A., Minnis, A. M., Novak, S. D., Myers, B., & Carney, T. (2016). The male factor: Outcomes from a cluster randomized field experiment with a couples-based HIV prevention intervention in a South African township. *Drug and Alcohol Dependence*, 161, 307–315. <https://doi.org/10.1016/j.drugalcdep.2016.02.017>
- *Weiland, A., Reiband, N., Schäffeler, N., Zurstiege, G., Giel, K. E., Zipfel, S., & Mack, I. (2022). A serious game for the prevention of obesity in school children—impact of parent’s involvement: A randomized controlled trial. *Life*, 12(6), Article 779. <https://doi.org/10.3390/life12060779>
- Weingarten, E., Chen, Q., McAdams, M., Yi, J., Hepler, J., & Albarracín, D. (2016). From primed concepts to action: A meta-analysis of the behavioral effects of incidentally presented words. *Psychological Bulletin*, 142(5), 472–497. <https://doi.org/10.1037/bul0000030>
- *Werner, T., Boutagy, N., Osterberg, K., Rivero, J., & Davy, K. (2016). Singular and combined effects of nebulivol and lifestyle modification on large artery stiffness in hypertensive adults. *Therapeutic Advances in Cardiovascular Diseases*, 7(6), 285–292. <https://doi.org/10.1177/1753944713513062>
- *West, J. S., Guelfi, K. J., Dimmock, J. A., & Jackson, B. (2018). Testing the feasibility and preliminary efficacy of an 8-week exercise and compensatory eating intervention. *Nutrients*, 10(7), Article 923. <https://doi.org/10.3390/nu10070923>
- White, H. (1980). A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica*, 48(4), 817–838. <https://doi.org/10.2307/1912934>
- *White, N., Carnahan, J., Nugent, C. A., Iwaoka, T., & Dodson, M. A. (1986). Management of obese patients with diabetes mellitus: Comparison of advice education with group management. *Diabetes Care*, 9(5), 490–496. <https://doi.org/10.2337/diacare.9.5.490>
- *Wijesuriya, M., Fountoulakis, N., Guess, N., Banneheka, S., Vasantharajah, L., Gulliford, M., Viberti, G., Gnudi, L., & Karalliedde, J. (2017). A pragmatic lifestyle modification programme reduces the incidence of predictors of cardio-metabolic disease and dysglycaemia in a young healthy urban South Asian population: A randomised controlled trial. *BMC Medicine*, 15(1), Article 146. <https://doi.org/10.1186/s12916-017-0905-6>
- *Wilhelmsen, L., Berglund, G., Elmfeldt, D., Tibblin, G., Wedel, H., Pennert, K., Vedin, A., Wilhelmsson, C., & Werkö, L. (1986). The multifactor primary prevention trial in Göteborg, Sweden. *European Heart Journal*, 7(4), 279–288. <https://doi.org/10.1093/oxfordjournals.eurheartj.a062065>
- *Wilkinson, S. A., & McIntyre, H. D. (2012). Evaluation of the ‘healthy start to pregnancy’ early antenatal health promotion workshop: A randomized controlled trial. *BMC Pregnancy and Childbirth*, 12(1), Article 131. <https://doi.org/10.1186/1471-2393-12-131>
- *Williams, A., Wiggers, J., O’Brien, K. M., Wolfenden, L., Yoong, S. L., Hodder, R. K., Lee, H., Robson, E. K., McAuley, J. H., Haskins, R.,

- Kamper, S. J., Rissel, C., & Williams, C. M. (2018). Effectiveness of a healthy lifestyle intervention for chronic low back pain: A randomised controlled trial. *Pain*, 159(6), 1137–1146. <https://doi.org/10.1097/j.pain.0000000000001198>
- *Williams, S. P., Myles, R. L., Sperling, C. C., & Carey, D. (2018). An intervention for reducing the sexual risk of men released from jails. *Journal of Correctional Health Care*, 24(1), 71–83. <https://doi.org/10.1177/1078345817745537>
- *Williamson, D. A., Martin, P. D., White, M. A., Newton, R., Walden, H., York-Crowe, E., Alfonso, A., Gordon, S., & Ryan, D. (2005). Efficacy of an internet-based behavioral weight loss program for overweight adolescent African-American girls. *Eating and Weight Disorders*, 10(3), 193–203. <https://doi.org/10.1007/BF03327547>
- *Wilson, D. B., Jones, R. M., McClish, D., Westerberg, A. L., & Danish, S. (2012). Fruit and vegetable intake among rural youth following a school-based randomized controlled trial. *Preventive Medicine: An International Journal Devoted to Practice and Theory*, 54(2), 150–156. <https://doi.org/10.1016/j.ypmed.2011.11.005>
- Wilson, K., Senay, I., Durantini, M., Sánchez, F., Hennessy, M., Spring, B., & Albarracín, D. (2015). When it comes to lifestyle recommendations, more is sometimes less: A meta-analysis of theoretical assumptions underlying the effectiveness of interventions promoting multiple behavior domain change. *Psychological Bulletin*, 141(2), 474–509. <https://doi.org/10.1037/a0038295>
- *Wilson, T. E., Gousse, Y., Joseph, M. A., Browne, R. C., Camilien, B., McFarlane, D., Mitchell, S., Brown, H., Urraca, N., Romeo, D., Johnson, S., Salifu, M., Stewart, M., Vavagiakis, P., & Fraser, M. (2019). HIV prevention for black heterosexual men: The barbershop talk with brothers cluster randomized trial. *American Journal of Public Health*, 109(8), 1131–1137. <https://doi.org/10.2105/AJPH.2019.305121>
- *Winston, G., Sifat, M., Phillips, E., Dietz, W., Wikner, E., Barrow, M., Khurana, K., & Charlson, M. (2019). Engaging children to support parental weight loss: A randomized trial. *Health Education & Behavior*, 46(5), 755–762. <https://doi.org/10.1177/1090198119853005>
- *Wister, A., Loewen, N., Kennedy-Symonds, H., McGowan, B., McCoy, B., & Singer, J. (2007). One-year follow-up of a therapeutic lifestyle intervention targeting cardiovascular disease risk. *Canadian Medical Association Journal*, 177(8), 859–865. <https://doi.org/10.1503/cmaj.061059>
- *Wolitski, R. J., Gómez, C. A., Parsons, J. T., & the SUMIT Study Group. (2005). Effects of a peer-led behavioral intervention to reduce HIV transmission and promote serostatus disclosure among HIV-seropositive gay and bisexual men. *Aids*, 19(Suppl. 1), S99–S109. <https://doi.org/10.1097/01.aids.0000167356.94664.59>
- Wolitski, R. J., Janssen, R. S., Onorato, I. M., Purcell, D. W., & Crepaz, N. (2005). An overview of prevention with people living with HIV. In S. C. Kalichman (Ed.), *Positive Prevention* (pp. 1–28). Springer. https://doi.org/10.1007/0-306-48700-4_1
- Wood, W. (2017). Habit in personality and social psychology. *Personality and Social Psychology Review*, 21(4), 389–403. <https://doi.org/10.1177/1088868317720362>
- Wood, W., & Neal, D. T. (2007). A new look at habits and the habit–goal interface. *Psychological Review*, 114(4), 843–863. <https://doi.org/10.1037/0033-295X.114.4.843>
- World Population Review. (2023). *Western countries 2023*. <https://worldpopulationreview.com/country-rankings/western-countries>
- Worley, M. J., Trim, R. S., Roesch, S. C., Mrnak-Meyer, J., Tate, S. R., & Brown, S. A. (2012). Comorbid depression and substance use disorder: Longitudinal associations between symptoms in a controlled trial. *Journal of Substance Abuse Treatment*, 43(3), 291–302. <https://doi.org/10.1016/j.jsat.2011.12.010>
- Wu, P., Wilson, K., Dimoulas, P., & Mills, E. J. (2006). Effectiveness of smoking cessation therapies: A systematic review and meta-analysis. *BMC Public Health*, 6(1), Article 300. <https://doi.org/10.1186/1471-2458-6-300>
- *Xu, H., Li, Y., Shang, X., Du, S., Zhang, Q., Liu, A., & Ma, G. (2020). Effect of comprehensive interventions including nutrition education and physical activity on high blood pressure among children: Evidence from school-based cluster randomized control trial in China. *International Journal of Environmental Research and Public Health*, 17(23), Article 8944. <https://doi.org/10.3390/ijerph17238944>
- *Xu, X., Chen, S., Chen, J., Chen, Z., Fu, L., Song, D., Zhao, M., & Jiang, H. (2021). Feasibility and preliminary efficacy of a community-based addiction rehabilitation electronic system in substance use disorder: Pilot randomized controlled trial. *JMIR mHealth and uHealth*, 9(4), Article e21087. <https://doi.org/10.2196/21087>
- *Yang, X., Tian, H., Zhang, F., Zhang, C., Li, Y., Leng, J., Wang, L., Liu, G., Dong, L., Yu, Z., Hu, G., & Chan, J. C. (2014). A randomised translational trial of lifestyle intervention using a 3-tier shared care approach on pregnancy outcomes in Chinese women with gestational diabetes mellitus but without diabetes. *Journal of Translational Medicine*, 12(1), Article 290. <https://doi.org/10.1186/s12967-014-0290-2>
- *Yeary, K. H. K., Cornell, C. E., Moore, P. C., Gauss, C. H., Prewitt, T. E., & Turner, J. (2020). The WORD: Outcomes of a behavioral weight loss maintenance effectiveness trial in rural black adults of faith. *Obesity*, 28(3), 510–520. <https://doi.org/10.1002/oby.22717>
- *Yonkers, K. A., Forray, A., Howell, H. B., Gotman, N., Kershaw, T., Rounsaville, B. J., & Carroll, K. M. (2012). Motivational enhancement therapy coupled with cognitive behavioral therapy versus brief advice: A randomized trial for treatment of hazardous substance use in pregnancy and after delivery. *General Hospital Psychiatry*, 34(5), 439–449. <https://doi.org/10.1016/j.genhosppsych.2012.06.002>
- *Zule, W. A., Costenbader, E. C., Coomes, C. M., & Wechsberg, W. M. (2009). Effects of an HCV risk reduction motivational intervention on alcohol use, injection and sexual risk behaviors. *American Journal of Public Health*, 99(S1), S180–S186. <https://doi.org/10.2105/AJPH.2007.126854>

Received June 26, 2022

Revision received December 8, 2023

Accepted December 29, 2023 ■